arXiv:0711.1937v1 [math.NT] 13 Nov 2007

EXPONENTIAL SUMS AND RANK OF DOUBLE PERSYMMETRIC MATRICES OVER $\mathbb{F}_2$

JORGEN CHERLY

Résumé. Soit $\mathbb{K}^2$ le $\mathbb{K}$ - espace vectoriel de dimension 2 où $\mathbb{K}$ dénote le corps des séries de Laurent formelles $\mathbb{F}_2((T^{-1}))$. Nous calculons en particulier des sommes exponentielles (dans $\mathbb{K}^2$) de la forme
$\sum_{\deg Y\leq k-1}\sum_{\deg Z\leq s-1}E(tYZ)\sum_{\deg U\leq s+m-1}E(\eta YU)$ où (t,η) est dans la boule unité de $\mathbb{K}^2$.
Nous démontrons qu'elles dépendent uniquement du rang de matrices doubles persymétriques avec des entrées dans $\mathbb{F}_2$, c'est-à-dire des matrices de la forme $\left[\frac{A}{B}\right]$ où A est une matrice $s\times k$ persymétrique et B une matrice $(s+m)\times k$ persymétrique (une matrice $[\alpha_{i,j}]$ est persymétrique si $\alpha_{i,j}=\alpha_{r,s}$ pour i+j = r+s). En outre, nous établissons plusieurs formules concernant des propriétés de rang de partitions de matrices doubles persymétriques, ce qui nous conduit à une formule récurrente du nombre $\Gamma_i^{\left[\substack{s\\ s+m}\right]\times k}$ des matrices de rang i de la forme $\left[\frac{A}{B}\right]$. Nous déduisons de cette formule récurrente que si $0\leq i\leq \inf(s-1,k-1)$, le nombre $\Gamma_i^{\left[\substack{s\\ s+m}\right]\times k}$ dépend uniquement de i. D'autre part, si $i\geq s+1, k\geq i$, $\Gamma_i^{\left[\substack{s\\ s+m}\right]\times k}$ peut être calculé à partir du nombre $\Gamma_{s'+1}^{\left[\substack{s'\\ s'+m'}\right]\times k'}$ de matrices de rang (s'+1) de la forme $\left[\frac{A'}{B'}\right]$ où A' est une matrice $s'\times k'$ persymétrique et B' une matrice $(s'+m')\times k'$ persymétrique, où s', m' et k' dépendent de i, s, m et k. La preuve de ce résultat est basée sur une formule (donnée dans [4]) du nombre de matrices de rang i de la forme $\left[\frac{A}{b_{-}}\right]$ où A est persymétrique et b_{-} une matrice ligne avec entrées dans $\mathbb{F}_2$. Nous montrons également que le nombre R de représentations dans $\mathbb{F}_2[T]$ des équations polynomiales

$$\begin{cases} YZ+Y_1Z_1+\ldots+Y_{q-1}Z_{q-1}=0\\ YU+Y_1U_1+\ldots+Y_{q-1}U_{q-1}=0\end{cases}$$

associées aux sommes exponentielles
$\sum_{degY\leq k-1}\sum_{degZ\leq s-1}E(tYZ)\sum_{\deg U\leq s+m-1}E(\eta YU)$ est donné par une intégrale sur la boule unité de $\mathbb{K}^2$ et est une combinaison linéaire de $\Gamma_i^{\left[\substack{s\\ s+m}\right]\times k}$ pour $i\geq 0$. Nous pouvons alors calculer explicitement le nombre R.

Abstract. Let $\mathbb{K}^2$ be the 2-dimensional vectorspace over $\mathbb{K}$ where $\mathbb{K}$ denotes the field of Laurent Series $\mathbb{F}_2((T^{-1}))$. We compute in particular exponential sums, (in $\mathbb{K}^2$) of the form $\sum_{\deg Y\leq k-1}\sum_{\deg Z\leq s-1}E(tYZ)\sum_{\deg U\leq s+m-1}E(\eta YU)$ where (t,η) is in the unit interval of $\mathbb{K}^2$. We show that they only depend on the rank of some associated double persymmetric matrices with entries in $\mathbb{F}_2$, that is matrices of the form $\left[\frac{A}{B}\right]$ where A is a $s\times k$ persymmetric matrix and B a $(s+m)\times k$ persymmetric matrix. (A matrix $[\alpha_{i,j}]$ is persymmetric if $\alpha_{i,j}=\alpha_{r,s}\ for\ i+j=r+s$). Besides, we establish several formulas concerning rank properties of partitions of double persymmetric matrices, which leads to a recurrent formula for the number $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}$ of rank i matrices of the form $\left[\frac{A}{B}\right]$. We deduce from the recurrent formula that if $0\leq i\leq\inf(s-1,k-1)$ then $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}$ depends only on i. On the other hand, if $i\geq s+1, k\geq i$, $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}$ can be computed from the number $\Gamma_{s'+1}^{\left[\substack{s'\\ s'+m'}\right]\times k'}$ of rank (s'+1) matrices of the form $\left[\frac{A'}{B'}\right]$ where A' is a $s'\times k'$ persymmetric matrix and B' a $(s'+m')\times k'$ persymmetric matrix, where s', m' and k' depend on i, s, m and k. The proof of this result is based on a formula (given in [4]) of the number of rank i matrices of the form $\left[\frac{A}{b_{-}}\right]$ where A is persymmetric and b_{-} a one-row matrix with entries in $\mathbb{F}_2$. We also prove that the number R of representations in $\mathbb{F}_2[T]$ of the polynomial equations

$$\begin{cases} YZ+Y_1Z_1+\ldots+Y_{q-1}Z_{q-1}=0\\ YU+Y_1U_1+\ldots+Y_{q-1}U_{q-1}=0\end{cases}$$

associated to the exponential sums $\sum_{degY\leq k-1}\sum_{degZ\leq s-1}E(tYZ)\sum_{\deg U\leq s+m-1}E(\eta YU)$ is given by an integral over the unit interval of $\mathbb{K}^2$, and is a linear combination of the $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}\ for\ i\geq 0$. We can then compute explicitly the number R.

Contents

1. **NOTATION** 5

1.1. **ANALYSIS ON $\mathbb{K}$** 5

1.2. **ANALYSIS ON THE TWO-DIMENSIONAL $\mathbb{K}$-VECTORSPACE** 6

2. **INTRODUCTION (REFER TO SECTION 1 AND SECTION 3)** 18

2.1. **SKETCH OF THE PROOF** 18

2.2. **ORGANIZATION OF THE PAPER** 22

3. **STATEMENT OF RESULTS** 41

4. **EXPONENTIAL SUMS FORMULAS ON $\mathbb{K}^2$** 46

5. **A RECURRENT FORMULA FOR THE NUMBER $\Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}$ OF RANK i MATRICES OF THE FORM $\left[\frac{A}{B}\right]$, WHERE A IS A $s\times k$ PERSYMMETRIC MATRIX AND B a $(s+m)\times k$ PERSYMMETRIC MATRIX WITH ENTRIES IN $\mathbb{F}_2$** 54

6. **RANK PROPERTIES OF A PARTITION OF DOUBLE PERSYMMETRIC MATRICES** 66

7. **RANK PROPERTIES OF SUBMATRICES OF DOUBLE PERSYMMETRIC MATRICES** 78

8. **STUDY OF THE REMAINDER $\Delta_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}$ IN THE RECURRENT FORMULA** 86

9. **A RECURRENT FORMULA FOR THE DIFFERENCE $\Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times (k+1)} - \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}, 1\leq i\leq 2s+m,\ k\geq i+1$** 91

10. **COMPUTATION OF THE REMAINDER R(j,s,m,k) IN THE RECURRENT FORMULA FOR THE DIFFERENCE $\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times (k+1)} - \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}\quad for\ 0\leq j\leq s+m,\ k>s+j$** 96

11. **COMPUTATION OF $\Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}\ for\ i\leq s-1,\ i\leq k-1,\ m\geq 0$** 99

12. **COMPUTATION OF $\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times (k+1)} - \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}\ for\ 0\leq j\leq s+m,\ k\geq s+j+1,\ m\geq 0$** 106

13. **COMPUTATION OF $\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}\ for\ j\in\{0,1\},\ k\geq s+j,\ m\geq 0$** 115

14. **COMPUTATION OF $\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}\ for\ j\in\{2,3\},\ k\geq s+j,\ j\leq m$** 125

15. **A REDUCTION FORMULA FOR $\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}$ in the case $1\leq j\leq m,\ k\geq s+j$** 133

16. **A REDUCTION FORMULA FOR $\Gamma_{s+j+1}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times k}$ IN THE CASE $1\leq j\leq s-1,\ k\geq s+j+1$** 140

17. **A REDUCTION FORMULA FOR $\Gamma_{s+m+1+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}$ IN THE CASE $1\leq j\leq s-1,\ k\geq s+m+1+j$** 149

18. **PROOF OF THEOREMS** $3.1, 3.2, \ldots, 3.14, 3.15$ 158

18.1. **PROOF OF THEOREM 3.1** 158

18.2. **PROOF OF THEOREM 3.2** 158

18.3. **PROOF OF THEOREM 3.3** 158

18.4. **PROOF OF THEOREM 3.4** 158

18.5. **PROOF OF THEOREM 3.5** 159

18.6. **PROOF OF THEOREM 3.6** 159
18.7. **PROOF OF THEOREM 3.7** 159
18.8. **PROOF OF THEOREM 3.8** 159
18.9. **PROOF OF THEOREM 3.9** 159
18.10. **PROOF OF THEOREM 3.10** 159
18.11. **PROOF OF THEOREM 3.11** 159
18.12. **PROOF OF THEOREM 3.12** 159
18.13. **PROOF OF THEOREM 3.13** 159
18.14. **PROOF OF THEOREM 3.14** 160
18.15. **PROOF OF THEOREM 3.15** 160
References 160

1. NOTATION

1.1. **ANALYSIS ON $\mathbb{K}$.** We denote by $\mathbb{F}_2\big((\frac{1}{T})\big) = \mathbb{K}$ the completion of the field $\mathbb{F}_2(T)$, the field of rational fonctions over the finite field $\mathbb{F}_2$, for the infinity valuation $\mathfrak{v} = \mathfrak{v}_\infty$ defined by $\mathfrak{v}\big(\frac{A}{B}\big) = degB - degA$ for each pair (A,B) of non-zero polynomials. Then every element non-zero t in $\mathbb{F}_2\big((\frac{1}{T})\big)$ can be expanded in a unique way in a convergent Laurent series $t = \sum_{j=-\infty}^{-\mathfrak{v}(t)} t_j T^j \ where \ t_j \in \mathbb{F}_2$.
We associate to the infinity valuation $\mathfrak{v} = \mathfrak{v}_\infty$ the absolute value $|\cdot|_\infty$ defined by

$$|t|_\infty = |t| = 2^{-\mathfrak{v}(t)}.$$

We denote E the Character of the additive locally compact group $\mathbb{F}_2\big((\frac{1}{T})\big)$ defined by

$$E\Big(\sum_{j=-\infty}^{-\mathfrak{v}(t)} t_j T^j\Big) = \begin{cases} 1 & \text{if } t_{-1} = 0, \\ -1 & \text{if } t_{-1} = 1. \end{cases}$$

We denote $\mathbb{P}$ the valuation ideal in $\mathbb{K}$, also denoted the unit interval of $\mathbb{K}$, i.e. the open ball of radius 1 about 0 or, alternatively, the set of all Laurent series

$$\sum_{i\geq 1} \alpha_i T^{-i} \quad (\alpha_i \in \mathbb{F}_2)$$

and, for every rational integer j, we denote by $\mathbb{P}_j$ the ideal $\{t \in \mathbb{K} | \ \mathfrak{v}(t) > j\}$. The sets $\mathbb{P}_j$ are compact subgroups of the additive locally compact group $\mathbb{K}$.
All $t \in \mathbb{F}_2\Big(\Big(\frac{1}{T}\Big)\Big)$ may be written in a unique way as $t = [t] + \{t\}$, $[t] \in \mathbb{F}_2[T]$, $\{t\} \in \mathbb{P}(= \mathbb{P}_0)$.
We denote by dt the Haar measure on $\mathbb{K}$ chosen so that

$$\int_{\mathbb{P}} dt = 1.$$

Definition 1.1. We introduce the following definitions in $\mathbb{K}$:

- Let s, m and k denote rational integers such that $s \geq 2, \ m \geq 0 \ and \ k \geq 1$.
- A matrix $D = [\alpha_{i,j}]$ is said to be persymmetric if $\alpha_{i,j} = \alpha_{r,s}$ whenever i+j = r+s.
- Set $t = \sum_{i\geq 1} \alpha_i T^{-i} \in \mathbb{P}$, we denote by $D_{s\times k}(t)$ the following $s \times k$ persymmetric matrix

$$\left(\begin{array}{cccccc} \alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1} \end{array}\right).$$

- Set $t=\sum_{i\geq 1}\alpha_i T^{-i}\in\mathbb{P}$, we denote by $D^{j}_{s\times(k-j+1)}(t)$ the following $s\times(k-j+1)$ persymmetric matrix
$$\begin{pmatrix}
\alpha_j & \alpha_{j+1} & \alpha_{j+2} & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_{j+1} & \alpha_{j+2} & \alpha_{j+3} & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{j+s-2} & \alpha_{j+s-1} & \alpha_{j+s} & \ldots & \alpha_{k+s-3} & \alpha_{k+s-2} \\
\alpha_{j+s-1} & \alpha_{j+s} & \alpha_{j+s+1} & \ldots & \alpha_{k+s-2} & \alpha_{k+s-1}
\end{pmatrix}.$$
- Set $\eta=\sum_{i\geq 1}\beta_i T^{-i}\in\mathbb{P}$, we denote by $D_{(s+m)\times k}(\eta)$ the following $(s+m)\times k$ persymmetric matrix
$$\begin{pmatrix}
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{pmatrix}.$$

1.2. ANALYSIS ON THE TWO-DIMENSIONAL $\mathbb{K}$-VECTORSPACE.

Let $\mathbb{K}\times\mathbb{K}=\mathbb{K}^2$ be the 2-dimensional vector space over $\mathbb{K}$. Let $(t,\eta)\in\mathbb{K}^2$ and $|(t,\eta)|=sup\{|t|,|\eta|\}=2^{-inf(\mathfrak{v}(t),\mathfrak{v}(\eta))}$.
It is easy to see that $(t,\eta)\longrightarrow|(t,\eta)|$ is an ultrametric valuation on $\mathbb{K}^2$, that is, $(t,\eta)\longrightarrow|(t,\eta)|$ is a norm and $|((t,\eta)+(t',\eta'))|\leq max\{|(t,\eta)|,|(t',\eta')|\}$.
We denote by $d(t,\eta)=dtd\eta$ the Haar measure on $\mathbb{K}^2$ chosen so that the measure on the unit interval of $\mathbb{K}^2$ is equal to one, thus
$$\iint_{\mathbb{P}\times\mathbb{P}}d(t,\eta)=\int_{\mathbb{P}}dt\int_{\mathbb{P}}d\eta=1\cdot 1=1.$$

Let $(t,\eta)=\left(\sum_{i=-\infty}^{-\mathfrak{v}(t)}t_iT^i,\sum_{i=-\infty}^{-\mathfrak{v}(\eta)}\eta_iT^i\right)\in\mathbb{K}^2$, we denote χ the Character on $(\mathbb{K}^2,+)$ defined by
$$\chi\Big(\sum_{i=-\infty}^{-\mathfrak{v}(t)}t_iT^i,\sum_{i=-\infty}^{-\mathfrak{v}(\eta)}\eta_iT^i\Big)=E\Big(\sum_{i=-\infty}^{-\mathfrak{v}(t)}t_iT^i\Big)\cdot E\Big(\sum_{i=-\infty}^{-\mathfrak{v}(\eta)}\eta_iT^i\Big)=\begin{cases}1 & \text{if } t_{-1}+\eta_{-1}=0,\\ -1 & \text{if } t_{-1}+\eta_{-1}=1.\end{cases}$$

Definition 1.2. We introduce the following definitions in the two-dimensional $\mathbb{K}$ -vectorspace.

- Let k, s and m denote rational integers such that $k\geq 1,\ s\geq 2\ and\ m\geq 0$.
- We denote by $\mathbb{P}/\mathbb{P}_i\times\mathbb{P}/\mathbb{P}_j$ a complete set of coset representatives of $\mathbb{P}_i\times\mathbb{P}_j$ in $\mathbb{P}\times\mathbb{P}$, for instance $\mathbb{P}/\mathbb{P}_{s+k-1}\times\mathbb{P}/\mathbb{P}_{s+m+k-1}$ denotes a complete set of coset representatives of $\mathbb{P}_{s+k-1}\times\mathbb{P}_{s+m+k-1}$ in $\mathbb{P}\times\mathbb{P}$.
- *Set* $(t,\eta)=(\sum_{i\geq 1}\alpha_iT^{-i},\sum_{i\geq 1}\beta_iT^{-i})\in\mathbb{P}\times\mathbb{P}$.

  We denote by $D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta)$ any $(2s+m)\times k$ matrix, such that after a rearrangement of the rows, if necessary, we can obtain the following double persymmetric matrix $\left[\frac{D_{s\times k}(t)}{D_{(s+m)\times k}(\eta)}\right]$

$$\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1} \\
\hline
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{array}\right).$$

We recall that the rank of a matrix does not change under elementary

row operations.

- Let j be a rational integer such that $1 \leq j \leq k-1$, *set* $(t,\eta) = (\sum_{i\geq 1}\alpha_i T^{-i}, \sum_{i\geq 1}\beta_i T^{-i}) \in \mathbb{P}\times\mathbb{P}$.

  We denote by $D_j^{\left[\substack{s\\ s+m}\right]\times(k-j+1)}(t,\eta)$ any $(2s+m)\times(k-j+1)$ matrix such that after a rearrangement of the rows, if necessary, we can obtain the following double persymmetric matrix $\left[\frac{D^j_{s\times(k-j+1)}(t)}{D^j_{(s+m)\times(k-j+1)}(\eta)}\right]$

$$\left(\begin{array}{cccccc}
\alpha_j & \alpha_{j+1} & \alpha_{j+2} & \cdots & \alpha_{k-1} & \alpha_k \\
\alpha_{j+1} & \alpha_{j+2} & \alpha_{j+3} & \cdots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{j+s-2} & \alpha_{j+s-1} & \alpha_{j+s} & \cdots & \alpha_{k+s-3} & \alpha_{k+s-2} \\
\alpha_{j+s-1} & \alpha_{j+s} & \alpha_{j+s+1} & \cdots & \alpha_{k+s-2} & \alpha_{k+s-1} \\
\hline
\beta_j & \beta_{j+1} & \beta_{j+2} & \cdots & \beta_{k-1} & \beta_k \\
\beta_{j+1} & \beta_{j+2} & \beta_{j+3} & \cdots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{j+s-2} & \beta_{j+s-1} & \beta_{j+s} & \cdots & \beta_{k+s-3} & \beta_{k+s-2} \\
\beta_{j+s-1} & \beta_{j+s} & \beta_{j+s+1} & \cdots & \beta_{k+s-2} & \beta_{k+s-1} \\
\beta_{j+s} & \beta_{j+s+1} & \beta_{j+s+2} & \cdots & \beta_{k+s-1} & \beta_{k+s} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{j+s+m-3} & \beta_{j+s+m-2} & \beta_{j+s+m-1} & \cdots & \beta_{k+s+m-4} & \beta_{k+s+m-3} \\
\beta_{j+s+m-2} & \beta_{j+s+m-1} & \beta_{j+s+m} & \cdots & \beta_{k+s+m-3} & \beta_{k+s+m-2} \\
\beta_{j+s+m-1} & \beta_{j+s+m} & \beta_{j+s+m+1} & \cdots & \beta_{k+s+m-2} & \beta_{k+s+m-1}
\end{array}\right).$$

For instance we can denote by $D_j^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k-j+1)}(t,\eta)$ the following matrix

$$\begin{pmatrix}
\alpha_j & \alpha_{j+1} & \alpha_{j+2} & \ldots & \alpha_{k-1} & \alpha_k \\
\beta_j & \beta_{j+1} & \beta_{j+2} & \ldots & \beta_{k-1} & \beta_k \\
\alpha_{j+1} & \alpha_{j+2} & \alpha_{j+3} & \ldots & \alpha_k & \alpha_{k+1} \\
\beta_{j+1} & \beta_{j+2} & \beta_{j+3} & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{j+s-2} & \alpha_{j+s-1} & \alpha_{j+s} & \ldots & \alpha_{k+s-3} & \alpha_{k+s-2} \\
\beta_{j+s-2} & \beta_{j+s-1} & \beta_{j+s} & \ldots & \beta_{k+s-3} & \beta_{k+s-2} \\
\beta_{j+s-1} & \beta_{j+s} & \beta_{j+s+1} & \ldots & \beta_{k+s-2} & \beta_{k+s-1} \\
\beta_{j+s} & \beta_{j+s+1} & \beta_{j+s+2} & \ldots & \beta_{k+s-1} & \beta_{k+s} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{j+s+m-3} & \beta_{j+s+m-2} & \beta_{j+s+m-1} & \ldots & \beta_{k+s+m-4} & \beta_{k+s+m-3} \\
\beta_{j+s+m-2} & \beta_{j+s+m-1} & \beta_{j+s+m} & \ldots & \beta_{k+s+m-3} & \beta_{s+m+k-2} \\
\alpha_{j+s-1} & \alpha_{j+s} & \alpha_{j+s+1} & \ldots & \alpha_{k+s-2} & \alpha_{k+s-1} \\
\beta_{j+s+m-1} & \beta_{j+s+m} & \beta_{j+s+m+1} & \ldots & \beta_{k+s+m-2} & \beta_{k+s+m-1}
\end{pmatrix}.$$

If j = 1 we denote $D_1^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}(t,\eta)$ by $D^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}(t,\eta)$.

- We denote by $D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ \alpha_{s-}+\beta_{s+m-} \end{smallmatrix}\right]\times k}(t,\eta)$ the following $(2s+m-1)\times k$ matrix, where the submatrix formed by the first (2s +m -2) rows is equal to the matrix $D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix}\right]\times k}(t,\eta)$, and the last row form a $1\times k$ persymmetric matrix of the form $(\alpha_s+\beta_{s+m}, \alpha_{s+1}+\beta_{s+m+1}, \ldots, \alpha_{s+k-1}+\beta_{s+k+m-1})$

$$\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\alpha_s+\beta_{s+m} & \alpha_{s+1}+\beta_{s+m+1} & \alpha_{s+2}+\beta_{s+m+2} & \ldots & \alpha_{k+s-2}+\beta_{k+s+m-2} & \alpha_{k+s-1}+\beta_{k+s+m-1}
\end{array}\right).$$

- We denote by ker D the nullspace of the matrix D and r(D) the rank of the matrix D.

- To simplify the notations concerning the exponential sums used in the proofs, we introduce the following definitions.

- Let $g_{s,k,m}(t,\eta)$ be the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto \sum_{degY\leq k-1}\sum_{degZ\leq s-1}E(tYZ)\sum_{degU\leq s+m-1}E(\eta YU)\in\mathbb{Z}.$$

- Let $g(t,\eta)$ be the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto \sum_{degY\leq k-1}\sum_{degZ\leq s-1}E(tYZ)\sum_{degU\leq s+m-1}E(\eta YU)\in\mathbb{Z}.$$

- Let $g_1(t,\eta)$ be the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto \sum_{degY\leq k-1}\sum_{degZ=s-1}E(tYZ)\sum_{degU\leq s+m-1}E(\eta YU)\in\mathbb{Z}.$$

- Let $g_2(t,\eta)$ be the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto \sum_{degY\leq k-1}\sum_{degZ\leq s-1}E(tYZ)\sum_{degU=s+m-1}E(\eta YU)\in\mathbb{Z}.$$

- Let $f_1(t,\eta)$ be the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto \sum_{degY\leq k-1}\sum_{degZ=s-1}E(tYZ)\sum_{degU\leq s+m-2}E(\eta YU)\in\mathbb{Z}.$$

- Let $f_2(t,\eta)$ be the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto \sum_{degY\leq k-1}\sum_{degZ\leq s-2}E(tYZ)\sum_{degU=s+m-1}E(\eta YU)\in\mathbb{Z}.$$

- Let $h(t,\eta)$ be the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto \sum_{degY\leq k-1}\sum_{degZ=s-1}E(tYZ)\sum_{degU=s+m-1}E(\eta YU)\in\mathbb{Z}.$$

- Let $v(t,\eta)$ be the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto \sum_{degY\leq k-1}\sum_{degZ\leq s-2}E(tYZ)\sum_{degU\leq s+m-2}E(\eta YU)\in\mathbb{Z}.$$

- Let $\psi(t,\eta)$ be the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto\sum_{degY=k-1}\sum_{degZ\leq s-1}E(tYZ)\sum_{degU\leq s+m-1}E(\eta YU)\in\mathbb{Z}.$$

- Let $\phi(t,\eta)$ be the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto\sum_{degY=k-1}\sum_{degZ\leq s-1}E(tYZ)\sum_{degU=s+m-2}E(\eta YU)\in\mathbb{Z}.$$

- Let $\phi_1(t,\eta)$ be the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto\sum_{degY=k-1}\sum_{degZ\leq s-1}E(tYZ)\sum_{degU\leq s+m-2}E(\eta YU)\in\mathbb{Z}.$$

- Let $\phi_2(t,\eta)$ be the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto\sum_{degY\leq k-2}\sum_{degZ\leq s-1}E(tYZ)\sum_{degU=s+m-1}E(\eta YU)\in\mathbb{Z}.$$

- Let $\theta_1(t,\eta)$ be the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto\sum_{degY=k-1}\sum_{degZ=s-1}E(tYZ)\sum_{degU\leq s+m-2}E(\eta YU)\in\mathbb{Z}.$$

- Let $\theta_2(t,\eta)$ be the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto\sum_{degY\leq k-2}\sum_{degZ\leq s-2}E(tYZ)\sum_{degU=s+m-1}E(\eta YU)\in\mathbb{Z}.$$

- Let $\theta_3(t,\eta)$ be the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto\sum_{degY\leq k-2}\sum_{degZ=s-1}E(tYZ)\sum_{degU\leq s+m-2}E(\eta YU)\in\mathbb{Z}.$$

- Consider the following partition of the matrix $D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta)$

$$\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \cdots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \cdots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \cdots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \cdots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \cdots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \cdots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \cdots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \cdots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \cdots & \alpha_{s+k-2} & \alpha_{s+k-1}
\end{array}\right).$$

We define

$$\sigma_{i,i}^{\left[\substack{s-1\\ s+m \\ \hline \alpha_{s-}}\right]\times k}$$

to be the cardinality of the following set

$$\left\{(t,\eta)\in \mathbb{P}/\mathbb{P}_{k+s-1}\times \mathbb{P}/\mathbb{P}_{k+s+m-1} \mid r(D^{\left[\substack{s-1\\ s+m}\right]\times k}(t,\eta)) = r(D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta)) = i\right\}.$$

- Consider the following partition of the matrix $D^{\left[\substack{s\\ s+m-1}\right]\times k}(t,\eta)$

$$\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \cdots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \cdots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \cdots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \cdots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \cdots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \cdots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \cdots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \cdots & \alpha_{s+k-2} & \alpha_{s+k-1}
\end{array}\right).$$

We define

$$\sigma_{i,i}^{\left[\substack{s-1\\ s+m-1\\ \overline{\alpha_{s-}}}\right]\times k}$$

to be the cardinality of the following set

$$\left\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-2}\mid r(D^{\left[\substack{s-1\\ s-1+m}\right]\times k}(t,\eta))=r(D^{\left[\substack{s\\ s+m-1}\right]\times k}(t,\eta))=i\right\}.$$

- We consider the following partition of the matrix $D^{\left[\substack{s-1\\ s+m-1\\ \alpha_{s-}+\beta_{s+m-}}\right]\times k}(t,\eta)$

$$\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k\\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2}\\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k\\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2}\\
\hline
\alpha_s+\beta_{s+m} & \alpha_{s+1}+\beta_{s+m+1} & \alpha_{s+2}+\beta_{s+m+2} & \ldots & \alpha_{k+s-2}+\beta_{k+s+m-2} & \alpha_{k+s-1}+\beta_{k+s+m-1}
\end{array}\right).$$

We define

$$\sigma_{i,i}^{\left[\substack{s-1\\ \frac{s+m-1}{\alpha_{s-}+\beta_{s+m-}}}\right]\times k}$$

to be the cardinality of the following set

$$\left\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\mid r(D^{\left[\substack{s-1\\ s-1+m}\right]\times k}(t,\eta))=r(D^{\left[\substack{s-1\\ s+m-1\\ \alpha_{s-}+\beta_{s+m-}}\right]\times k}(t,\eta))=i\right\}.$$

- Consider the following partition of the matrix $D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta)$

$$\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\
\hline
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{array}\right).$$

We define

$$\sigma_{i,i,i}^{\left[\begin{smallmatrix} \frac{s-1}{s+m-1} \\ \frac{\alpha_{s-}}{\beta_{s+m-}} \end{smallmatrix}\right]\times k}$$

to be the cardinality of the following set

$$\left\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\mid r(D^{\left[\overset{s-1}{s-1+m}\right]\times k}(t,\eta)) = r(D^{\left[\overset{s}{s+m-1}\right]\times k}(t,\eta)) = r(D^{\left[\overset{s-1}{s+m}\right]\times k}(t,\eta)) = i\right\}$$

$$=\left\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\mid r(D^{\left[\overset{s-1}{s-1+m}\right]\times k}(t,\eta)) = r(D^{\left[\overset{s}{s+m-1}\right]\times k}(t,\eta)) = r(D^{\left[\overset{s}{s+m}\right]\times k}(t,\eta)) = i\right\}.$$

- Let $(j_1,j_2,j_3)\in\mathbb{N}^3$ we define

$$\sigma_{j_1,j_2,j_3}^{\left[\begin{smallmatrix} \frac{s-1}{s+m-1} \\ \frac{\alpha_{s-}}{\beta_{s+m-}} \end{smallmatrix}\right]\times k}$$

to be the cardinality of the following set

$$\Big\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\mid r(D^{\left[\overset{s-1}{s+m-1}\right]\times k}(t,\eta)) = j_1,\quad r(D^{\left[\overset{s}{s+m-1}\right]\times k}(t,\eta)) = j_2,$$
$$r(D^{\left[\overset{s}{s+m}\right]\times k}(t,\eta)) = j_3\Big\}.$$

- Consider the following partition of the matrix $D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta)$

$$\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1}
\end{array}\right).$$

Let $(j_1,j_2,j_3)\in\mathbb{N}^3$ we define

$$\sigma_{j_1,j_2,j_3}^{\left[\substack{s-1\\ \frac{s+m-1}{\beta_{s+m-}}\\ \alpha_{s-}}\right]\times k}$$

to be the cardinality of the following set

$$\Big\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\mid r(D^{\left[\substack{s-1\\ s+m-1}\right]\times k}(t,\eta))=j_1,\quad r(D^{\left[\substack{s-1\\ s+m}\right]\times k}(t,\eta))=j_2,$$
$$r(D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta))=j_3\Big\}.$$

- Consider the following partition of the matrix $D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta)$

$$\left(\begin{array}{ccccc|c}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1} \\
\hline
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{array}\right).$$

Let $(j_1, j_2, j_3, j_4, j_5, j_6) \in \mathbb{N}^6$, we define

$$\#\left(\begin{array}{c|c} j_1 & j_2 \\ \hline j_3 & j_4 \\ \hline j_5 & j_6 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}$$

to be the cardinality of the following set

$$\Big\{(t,\eta) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \mid r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t,\eta)) = j_1, \quad r(D^{\left[\substack{s-1\\ s+m-1}\right]\times k}(t,\eta)) = j_2,$$
$$r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-1)}(t,\eta)) = j_3, \quad r(D^{\left[\substack{s\\ s+m-1}\right]\times k}(t,\eta)) = j_4,$$
$$r(D^{\left[\substack{s\\ s+m}\right]\times(k-1)}(t,\eta)) = j_5, \quad r(D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta)) = j_6\Big\}.$$

- Consider the following partition of the matrix $D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta)$

$$\left(\begin{array}{ccccc|c}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1}
\end{array}\right).$$

Let $(j_1, j_2, j_3, j_4, j_5, j_6) \in \mathbb{N}^6$, we define

$$\#\left(\begin{array}{c|c} j_1 & j_2 \\ \hline j_3 & j_4 \\ \hline j_5 & j_6 \end{array}\right)^{\frac{\beta}{\alpha}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}$$

to be the cardinality of the following set

$$\Big\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1} \mid r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t,\eta))=j_1,\quad r(D^{\left[\substack{s-1\\ s+m-1}\right]\times k}(t,\eta))=j_2,$$
$$r(D^{\left[\substack{s-1\\ s+m}\right]\times(k-1)}(t,\eta))=j_3,\quad r(D^{\left[\substack{s-1\\ s+m}\right]\times k}(t,\eta))=j_4,$$
$$r(D^{\left[\substack{s\\ s+m}\right]\times(k-1)}(t,\eta))=j_5,\quad r(D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta))=j_6\Big\}.$$

- Consider the following partition of the matrix $D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta)$

$$\left(\begin{array}{ccccc|c}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1} \\
\hline
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{array}\right).$$

Let $(j_1,j_2,j_3,j_4)\in\mathbb{N}^4$, we define

$$\#\left(\begin{array}{c|c} j_1 & j_2 \\ \hline j_3 & j_4 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}$$

to be the cardinality of the following set

$$\Big\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1} \mid r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-1)}(t,\eta))=j_1,\quad r(D^{\left[\substack{s\\ s+m-1}\right]\times k}(t,\eta))=j_2,$$
$$r(D^{\left[\substack{s\\ s+m}\right]\times(k-1)}(t,\eta))=j_3,\quad r(D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta))=j_4\Big\}.$$

- *Set* $(t,\eta)=(\sum_{i\geq 1}\alpha_i T^{-i},\sum_{i\geq 1}\beta_i T^{-i})\in\mathbb{P}\times\mathbb{P}.$
  Let $\Gamma_i^{\left[\substack{s\\ s+m}\right]\times k}$ denote the number of double persymmetric $(2s+m)\times k$ rank i matrices of the form $\left[\frac{D_{s\times k}(t)}{D_{(s+m)\times k}(\eta)}\right]$,that is

$$\Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = Card\left\{(t,\eta)\in \mathbb{P}/\mathbb{P}_{k+s-1}\times \mathbb{P}/\mathbb{P}_{k+s+m-1} \mid r(D^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}(t,\eta)) = i\right\}.$$

- Set for $0 \leq i \leq \inf(2s+m,k)$

$$\Delta_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = \sigma_{i,i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right]\times k} - 3\cdot\sigma_{i-1,i-1,i-1}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right]\times k} + 2\cdot\sigma_{i-2,i-2,i-2}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right]\times k}.$$

- Set for $0 \leq j \leq s+m,\ k \geq s+j+1$

$$\Omega_{j+1}(1,m+s-1,k) = \Gamma_{j+1}^{\left[\begin{smallmatrix} 1 \\ 1+m+(s-1) \end{smallmatrix}\right]\times k} - 4\cdot\Gamma_j^{\left[\begin{smallmatrix} 1 \\ 1+m+(s-2) \end{smallmatrix}\right]\times k}.$$

- We define

$k-1$	k	
$\cdot$	$\cdot$	$D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix}\right]\times\cdot}$
j	j	α_s-
j	$j+1$	$\beta_{s+m}-$

  to be the following subset of $\mathbb{P}^2$

$$\Big\{(t,\eta) = (\textstyle\sum_{i\geq 1}\alpha_i T^{-i}, \sum_{i\geq 1}\beta_i T^{-i}) \in \mathbb{P}^2 \mid r(D^{\left[\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix}\right]\times(k-1)}(t,\eta)) = j,$$
$$r(D^{\left[\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix}\right]\times k}(t,\eta)) = j, \quad r(D^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k-1)}(t,\eta)) = j, \quad r(D^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}(t,\eta)) = j+1\Big\}$$

- We define

$k-1$	k	
j	j	$D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix}\right]\times\cdot}$
j	$j+1$	α_s-
j	$j+1$	$\beta_{s+m}-$

  to be the following subset of $\mathbb{P}^2$

$$\Big\{(t,\eta) = (\textstyle\sum_{i\geq 1}\alpha_i T^{-i}, \sum_{i\geq 1}\beta_i T^{-i}) \in \mathbb{P}^2 \mid r(D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix}\right]\times(k-1)}(t,\eta)) = j,$$
$$r(D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix}\right]\times k}(t,\eta)) = j, \quad r(D^{\left[\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix}\right]\times(k-1)}(t,\eta)) = j, \quad r(D^{\left[\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix}\right]\times k}(t,\eta)) = j+1,$$
$$r(D^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k-1)}(t,\eta)) = j, \quad r(D^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}(t,\eta)) = j+1\Big\}$$

- Similar expressions are defined in a similar way.
- We denote by $R_q(k,s,m)$ the number of solutions $(Y_1,Z_1,U_1,\ldots,Y_q,Z_q,U_q)$ of the polynomial equations

$$\begin{cases} Y_1Z_1 + Y_2Z_2 + \ldots + Y_qZ_q = 0 \\ Y_1U_1 + Y_2U_2 + \ldots + Y_qU_q = 0 \end{cases}$$

  satisfying the degree conditions

$$degY_i \leq k-1, \quad degZ_i \leq s-1, \quad degU_i \leq s+m-1 \quad for \quad 1 \leq i \leq q.$$

2. INTRODUCTION (REFER TO SECTION 1 AND SECTION 3)

The rational function field $\mathbb{F}_2(T)$ is completed with respect to an appropriate valuation to a field $\mathbb{K}$ (i.e. the field of Laurent Series). The unit interval of $\mathbb{K}$, that is, the open ball of radius 1 about 0, is a compact additive group. We shall use the Haar integral on this group. Let $\mathbb{K}^2$ be the 2-dimensional vectorspace over $\mathbb{K}$.

The main result of this paper is to obtain a formula for the number $\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times k}$ of double persymmetric $(2s+m)\times k$ rank i matrices.
In particular we shall show that if $0\leq i\leq \inf(s-1,k-1)$ the number of double persymmetric $(2s+m)\times k$ rank i matrices depends only on i and if $i\geq s+1,\ k\geq i$ the number of rank i matrices can be obtained from the formula for the number of rank s+1 matrices. The computation of the $\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times k}$ is based on a formula (given in [4]) of the number of rank i matrices of the form $\left[\frac{A}{b_-}\right]$ where A is persymmetric and b_- a one-row matrix with entries in $\mathbb{F}_2$. We observe that the $\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times k}$ where $0\leq i\leq \inf(2s+m,k)$ are solutions to the system

$$\begin{cases} \displaystyle\sum_{i=0}^{\inf(2s+m,k)} \Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times k} = 2^{2k+2s+m-2}, \\ \displaystyle\sum_{i=0}^{\inf(2s+m,k)} \Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times k}\cdot 2^{-i} = 2^{k+2s+m-2}+2^{2k-2}-2^{k-2}. \end{cases} \tag{2.1}$$

The first equation is obvious.
The second equation is a consequence of the identity

$$\begin{aligned}\int_{\mathbb{P}\times\mathbb{P}} g(t,\eta)dtd\eta &= Card\{(Y,Z,U), degY\leq k-1, degZ\leq s-1, degU\leq s+m-1 \mid Y\cdot Z = Y\cdot U = 0\} \\ &= 2^{2s+m}+2^{k}-1.\end{aligned}$$

2.1. SKETCH OF THE PROOF.

- Consider the following $\left[\substack{s \\ s+m}\right]\times k$ double persymmetric matrix

$$\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1} \\
\hline
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{array}\right).$$

- We establish the following recurrent formula for the number $\Gamma_i^{\left[\substack{s\\ s+m}\right]\times k}$ of $\left[\substack{s\\ s+m}\right]\times k$ double persymmetric matrices of rank i:

$$\Gamma_i^{\left[\substack{s\\ s+m}\right]\times k} = 2\cdot\Gamma_{i-1}^{\left[\substack{s-1\\ s-1+(m+1)}\right]\times k} + 4\cdot\Gamma_{i-1}^{\left[\substack{s\\ s+(m-1)}\right]\times k} - 8\cdot\Gamma_{i-2}^{\left[\substack{s-1\\ s-1+m}\right]\times k} + \Delta_i^{\left[\substack{s\\ s+m}\right]\times k}. \tag{1}$$

We observe that the $\Gamma_i^{\left[\substack{s\\ s+m}\right]\times k}$, $for\ 0 \leq i \leq \inf(2s+m,k)$ are solutions to the system

$$\begin{cases} \sum_{i=0}^{\inf(2s+m,k)} \Gamma_i^{\left[\substack{s\\ s+m}\right]\times k} = 2^{2k+2s+m-2}, \\ \sum_{i=0}^{\inf(2s+m,k)} \Gamma_i^{\left[\substack{s\\ s+m}\right]\times k}\cdot 2^{-i} = 2^{k+2s+m-2} + 2^{2k-2} - 2^{k-2}. \end{cases} \tag{2}$$

- We establish that the remainder in (1) satisfies the following equality

$$\Delta_i^{\left[\substack{s\\ s+m}\right]\times k} = \Delta_i^{\left[\substack{s\\ s+m}\right]\times (i+1)} \quad for\ all \quad 0 \leq i \leq 2s+m,\ k \geq i+1. \tag{3}$$

- We establish from (1) and (3) a recurrent formula for the difference

$$\Gamma_i^{\left[\substack{s\\ s+m}\right]\times (k+1)} - \Gamma_i^{\left[\substack{s\\ s+m}\right]\times k} \quad for\ 2 \leq i \leq 2s+m, \quad k \geq i+1.$$

In fact we prove that

$$(4)\quad \Gamma_{i}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times(k+1)} - \Gamma_{i}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times k} = 4\cdot\left[\Gamma_{i-1}^{\begin{bmatrix} s \\ s+(m-1) \end{bmatrix}\times(k+1)} - \Gamma_{i-1}^{\begin{bmatrix} s \\ s+(m-1) \end{bmatrix}\times k}\right] \quad \text{if} \quad 1\leq i\leq s-1, \quad k\geq i+1,$$

$$(5)\quad \Gamma_{s+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times(k+1)} - \Gamma_{s+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times k} = 4\cdot\left[\Gamma_{s+j-1}^{\begin{bmatrix} s \\ s+(m-1) \end{bmatrix}\times(k+1)} - \Gamma_{s+j-1}^{\begin{bmatrix} s \\ s+(m-1) \end{bmatrix}\times k}\right] + R(j,s,m,k) \quad \text{if } 0\leq j\leq \inf(s+m,k-s+1)$$

where R(j,s,m,k) is equal to

$$2^{s-1}\left[\Gamma_{j+1}^{\begin{bmatrix} 1 \\ 1+(s+m-1) \end{bmatrix}\times(k+1)} - \Gamma_{j+1}^{\begin{bmatrix} 1 \\ 1+(s+m-1) \end{bmatrix}\times k}\right] - 2^{s+1}\left[\Gamma_{j}^{\begin{bmatrix} 1 \\ 1+(s+m-2) \end{bmatrix}\times(k+1)} - \Gamma_{j}^{\begin{bmatrix} 1 \\ 1+(s+m-2) \end{bmatrix}\times k}\right].$$

- From a formula of $\Gamma_{i}^{\begin{bmatrix} 1 \\ 1+m \end{bmatrix}\times k}$ obtained in [4] we compute the remainder R(j,s,m,k) in (5).
  In fact we prove

$$(6)\quad R(j,s,m,k) = \begin{cases} 2^{k+s-1} & \text{if } j=0, \quad k\geq s+1, \\ -2^{k+s-1} & \text{if } j=1, \quad k\geq s+2, \\ 0 & \text{if } 2\leq j\leq s+m-1, \quad k\geq s+j+1, \\ -3\cdot 2^{2k+2s+m-2} & \text{if } j=s+m, \quad k\geq 2s+m+1. \end{cases}$$

- From (4) applying (2) we deduce by induction on i

$$(7)\quad \Gamma_{i}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times k} = \begin{cases} \Gamma_{i}^{\begin{bmatrix} i \\ i \end{bmatrix}\times(i+1)} = 21\cdot 2^{3i-4} - 3\cdot 2^{2i-3} & \text{if} \quad 1\leq i\leq s-1,\ k\geq i+1, \\ 2^{2s+2i+m-2} - 3\cdot 2^{3i-4} + 2^{2i-3} & \text{if} \quad 1\leq i=k\leq s+1. \end{cases}$$

  We have now computed $\Gamma_{i}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times k}$ $for\ 1\leq i\leq s-1,\ k\geq i.$

- From (5) applying (6) we obtain

$$(8)\quad \Gamma_{s+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times(k+1)} - \Gamma_{s+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times k} = 4\cdot\left[\Gamma_{s+j-1}^{\begin{bmatrix} s \\ s+(m-1) \end{bmatrix}\times(k+1)} - \Gamma_{s+j-1}^{\begin{bmatrix} s \\ s+(m-1) \end{bmatrix}\times k}\right]$$

$$+\begin{cases} 2^{k+s-1} & \text{if } j=0, \quad k\geq s+1, \\ -2^{k+s-1} & \text{if } j=1, \quad k\geq s+2, \\ 0 & \text{if } 2\leq j\leq s+m-1, \quad k\geq s+j+1, \\ -3\cdot 2^{2k+2s+m-2} & \text{if } j=s+m, \quad k\geq 2s+m+1. \end{cases}$$

- Using successively the recurrent formula in (8) we deduce
  In the case $m\in\{0,1\}$

$$\Gamma_{s+j}^{\begin{bmatrix} s \\ s \end{bmatrix}\times(k+1)} - \Gamma_{s+j}^{\begin{bmatrix} s \\ s \end{bmatrix}\times k} = \begin{cases} 3\cdot 2^{k+s-1} & \text{if } j=0,\ k>s, \\ 21\cdot 2^{k+s+3j-4} & \text{if } 1\leq j\leq s-1,\ k>s+j, \\ 3\cdot 2^{2k+2s-2} - 3\cdot 2^{k+4s-4} & \text{if } j=s,\ k>2s, \end{cases}$$

$$\Gamma_{s+j}^{\left[\substack{s\\ s+1}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\substack{s\\ s+1}\right]\times k} = \begin{cases} 2^{k+s-1} & \text{if } j=0,\ k>s,\\ 11\cdot 2^{k+s-1} & \text{if } j=1,\ k>s+1,\\ 21\cdot 2^{k+s+3j-5} & \text{if } 2\leq j\leq s,\ k>s+j,\\ 3\cdot 2^{2k+2s-1}-3\cdot 2^{k+4s-2} & \text{if } j=s+1,\ k>2s+1. \end{cases}$$

In the case $m\geq 2$

$$\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k} = \begin{cases} 2^{k+s-1} & \text{if } j=0,\ k>s,\\ 3\cdot 2^{k+s+2j-3} & \text{if } 1\leq j\leq m-1,\ k>s+j,\\ 11\cdot 2^{k+s+2m-3} & \text{if } j=m,\ k>s+m, \end{cases} \tag{9}$$

$$\Gamma_{s+m+j}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{s+m+j}^{\left[\substack{s\\ s+m}\right]\times k} = \begin{cases} 21\cdot 2^{k+s+2m+3j-4} & \text{if } 1\leq j\leq s-1,\ k>s+m+j,\\ 3\cdot 2^{2k+2s+m-2}-3\cdot 2^{k+4s+2m-4} & \text{if } j=s,\ k>2s+m. \end{cases} \tag{10}$$

- Consider the following array $\left(\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k}\right)_{0\leq j\leq s+m,\ k\geq s+j}$:

$$\begin{array}{ccccccc}
\Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times s} & \Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times(s+1)} & \Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times(s+2)} & \dots & \Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times k} & \dots & k\geq s\\
\Gamma_{s+1}^{\left[\substack{s\\ s+m}\right]\times(s+1)} & \Gamma_{s+1}^{\left[\substack{s\\ s+m}\right]\times(s+2)} & \Gamma_{s+1}^{\left[\substack{s\\ s+m}\right]\times(s+3)} & \dots & \Gamma_{s+1}^{\left[\substack{s\\ s+m}\right]\times k} & \dots & k\geq s+1\\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots\\
\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times(s+j)} & \Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times(s+j+1)} & \Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times(s+j+2)} & \dots & \Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k} & \dots & k\geq s+j\\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots\\
\Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times(s+m)} & \Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times(s+m+1)} & \Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times(s+m+2)} & \dots & \Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times k} & \dots & k\geq s+m\\
\Gamma_{s+m+1}^{\left[\substack{s\\ s+m}\right]\times(s+m+1)} & \Gamma_{s+m+1}^{\left[\substack{s\\ s+m}\right]\times(s+m+2)} & \Gamma_{s+m+1}^{\left[\substack{s\\ s+m}\right]\times(s+m+3)} & \dots & \Gamma_{s+m+1}^{\left[\substack{s\\ s+m}\right]\times k} & \dots & k\geq s+m+1\\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots\\
\Gamma_{s+m+1+j}^{\left[\substack{s\\ s+m}\right]\times(s+m+1+j)} & \Gamma_{s+m+1+j}^{\left[\substack{s\\ s+m}\right]\times(s+m+2+j)} & \Gamma_{s+m+1+j}^{\left[\substack{s\\ s+m}\right]\times(s+m+3+j)} & \dots & \Gamma_{s+m+1+j}^{\left[\substack{s\\ s+m}\right]\times k} & \dots & k\geq s+m+1+j\\
\Gamma_{2s+m-1}^{\left[\substack{s\\ s+m}\right]\times(2s+m-1)} & \Gamma_{2s+m-1}^{\left[\substack{s\\ s+m}\right]\times(2s+m)} & \Gamma_{2s+m-1}^{\left[\substack{s\\ s+m}\right]\times(2s+m+1)} & \dots & \Gamma_{2s+m-1}^{\left[\substack{s\\ s+m}\right]\times k} & \dots & k\geq 2s+m-1\\
\Gamma_{2s+m}^{\left[\substack{s\\ s+m}\right]\times(2s+m)} & \Gamma_{2s+m}^{\left[\substack{s\\ s+m}\right]\times(2s+m+1)} & \Gamma_{2s+m}^{\left[\substack{s\\ s+m}\right]\times(2s+m+2)} & \dots & \Gamma_{2s+m}^{\left[\substack{s\\ s+m}\right]\times k} & \dots & k\geq 2s+m.
\end{array}$$

- To compute the elements in the array $\left(\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k}\right)_{0\leq j\leq s+m,\ k\geq s+j}$, we proceed by induction on j as follows:
  Let l be a rational integer such that $0\leq l\leq s+m-1$.
  Assume that we have computed the elements in the following subarray $\left(\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k}\right)_{0\leq j\leq l,\ k\geq s+j}$.

Recall that from (7) $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}$ are known for $0\leq i\leq s-1,\ k\geq i.$

From the first equation in (2) with k = s+l+1, $\sum_{i=0}^{s+l+1}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times (s+l+1)} = 2^{4s+2l+m}$

we deduce $\Gamma_{s+l+1}^{\left[\substack{s\\ s+m}\right]\times (s+l+1)}$ since the terms $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times (s+l+1)}$ for $i\leq s+l$ are known.

From the equations in (2) with $k=s+l+2$ we have

$$\begin{cases}\sum_{i=0}^{s+l+2}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times (s+l+2)} = 2^{4s+2l+m+2},\\ \sum_{i=0}^{s+l+2}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times (s+l+2)}\cdot 2^{-i} = 2^{3s+m+l}+2^{2s+2l+2}-2^{s+l}.\end{cases}$$

We then deduce $\Gamma_{s+l+1}^{\left[\substack{s\\ s+m}\right]\times (s+1+2)}$ and $\Gamma_{s+l+2}^{\left[\substack{s\\ s+m}\right]\times (s+l+2)}$ since the terms $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times (s+l+2)}$ for $i\leq s+l$ are known.

Then from (9) or (10) we compute $\Gamma_{s+l+1}^{\left[\substack{s\\ s+m}\right]\times k}$ knowing $\Gamma_{s+l+1}^{\left[\substack{s\\ s+m}\right]\times (s+1+2)}$ for all $k\geq s+l+2$.

We have now computed any element in the $l+1$-th row and the first element in the $l+2$-th row in the array $\left(\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k}\right)_{0\leq j\leq s+m,\ k\geq s+j}$.

- Using the procedure above we prove successively by induction on j the following reduction formulas
$\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k} = 8^{j-1}\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+(m-(j-1))}\right]\times (k-(j-1))}\quad for\quad 1\leq j\leq m,\ k\geq s+j,$
$\Gamma_{s+m+1+j}^{\left[\substack{s\\ s+m}\right]\times k} = 8^{2j+m}\cdot\Gamma_{s-j+1}^{\left[\substack{s-j\\ s-j}\right]\times (k-m-2j)}\quad for\quad 0\leq j\leq s-1,\ k\geq s+m+1+j.$

2.2. **ORGANIZATION OF THE PAPER.** We proceed as follows:

In Section 1 are introduced main notations and definitions.

In Section 3, we state the main theorems in the two dimensional $\mathbb{K}$-vector space.

In Section 4, we establish results on exponential sums, (in $\mathbb{P}\times\mathbb{P}$) of the form $g(t,\eta), h(t,\eta)$ or similar sums and we show that they only depend on rank properties of some corresponding double persymmetric matrices with entries in $\mathbb{F}_2$.

Consider in particular the quadratic exponential sum in $\mathbb{P}\times\mathbb{P}$ defined by

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto g(t,\eta)=\sum_{degY\leq k-1}\sum_{degZ\leq s-1}E(tYZ)\sum_{degU\leq s+m-1}E(\eta YU)\in\mathbb{Z}.$$

We associate to the exponential sum $g(t,\eta)$ the following double persymmetric matrix $\left[\frac{D_{s\times k}(t)}{D_{(s+m)\times k}(\eta)}\right]$,

$$\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1} \\
\hline
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{array}\right).$$

We get

(2.2)

$$g(t,\eta)=\sum_{degY\leq k-1}\sum_{degZ\leq s-1}E(tYZ)\sum_{degU\leq s+m-1}E(\eta YU)=2^{2s+m+k-r(D^{\left[\substack{s\\s+m}\right]\times k}(t,\eta))}.$$

The proof of (2.2) is based on the following two similar identities

$$\sum_{degZ\leq s-1}E(tYZ)=2^{s}\quad if\ and\ only\ if\quad Y\in\ker D_{s\times k}(t),$$

$$\sum_{degU\leq s+m-1}E(\eta YU)=2^{s+m}\quad if\ and\ only\ if\quad Y\in\ker D_{(s+m)\times k}(\eta).$$

Furthermore we establish similar results concerning similar exponential sums.
For instance we have
Let $(t,\eta)\in\mathbb{P}\times\mathbb{P}$ and set

$$g_2(t,\eta)=\sum_{degY\leq k-1}\sum_{degZ\leq s-1}E(tYZ)\sum_{degU=s-1+m}E(\eta YU).$$

Then
(2.3)

$$g_2(t,\eta)=\begin{cases}2^{2s+m+k-1-r(D^{\left[\substack{s\\s+m}\right]\times k}(t,\eta))} & \text{if } r(D^{\left[\substack{s\\s+m}\right]\times k}(t,\eta))=r(D^{\left[\substack{s\\s+(m-1)}\right]\times k}(t,\eta)),\\ 0 & \text{otherwise .}\end{cases}$$

In the end of the section we obtain by (2.2), observing that $g(t,\eta)$ is constant on cosets of $\mathbb{P}_{k+s-1}\times\mathbb{P}_{k+s+m-1}$

(2.4)

$$
\begin{aligned}
R(q,k-1,s-1,s+m-1,0,0) &= \int_{\mathbb{P}\times\mathbb{P}} g^{q}(t,\eta)\,dt\,d\eta \\
&= 2^{(2s+m+k)(q-1)}\cdot 2^{-k+2}\cdot \sum_{i=0}^{\inf(2s+m,k)} \Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}\cdot 2^{-qi}.
\end{aligned}
$$

In Section 5, we establish a recurrent formula for the number $\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}$ of rank i matrices of the form $\left[\frac{A}{B}\right]$, where A is a $s\times k$ persymmetric matrix and B a $(s+m)\times k$ persymmetric matrix with entries in $\mathbb{F}_2$.

Consider the following two partitions of the matrix $D^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}(t,\eta)$

$$
\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1} \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{array}\right)
\text{ and }
\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1}
\end{array}\right).
$$

Obviously $\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}$ can be represented in the following two ways

$$
\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = \sigma_{i,i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right]\times k} + \sigma_{i-1,i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right]\times k} + \sigma_{i-1,i-1,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right]\times k} + \sigma_{i-2,i-1,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right]\times k}, \tag{2.5}
$$

$$
\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = \sigma_{i,i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \beta_{s+m-} \\ \alpha_{s-} \end{array}\right]\times k} + \sigma_{i-1,i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \beta_{s+m-} \\ \alpha_{s-} \end{array}\right]\times k} + \sigma_{i-1,i-1,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \beta_{s+m-} \\ \alpha_{s-} \end{array}\right]\times k} + \sigma_{i-2,i-1,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \beta_{s+m-} \\ \alpha_{s-} \end{array}\right]\times k}. \tag{2.6}
$$

We establish the recurrent formula by obtaining relations between the terms of the two expressions of $\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}$. By considering the following partition of the matrix $D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ \alpha_{s-}+\beta_{s+m-} \end{smallmatrix}\right]\times k}(t,\eta)$,

$$\left(\begin{array}{c|cccccc}
r_1 & \alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
r_2 & \alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
r_{s-1} & \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
r_s & \beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
r_{s+1} & \beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
r_{s+m} & \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
r_{2s+m-2} & \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
r_{2s+m-1} & \alpha_s+\beta_{s+m} & \alpha_{s+1}+\beta_{s+m+1} & \alpha_{s+2}+\beta_{s+m+2} & \ldots & \alpha_{s+k-2}+\beta_{s+k+m-2} & \alpha_{s+k-1}+\beta_{s+k+m-1}
\end{array}\right)$$

and performing the following row operations, adding the s+m+j-th row to the j-th row for $0 \leq j \leq s-2$, we obtain by elementary rank considerations the following identity

$$\sigma_{i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-}+\beta_{s+m-} \end{array}\right]\times k} = 2 \cdot \sigma_{i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \end{array}\right]\times k} \quad for\ 0 \leq i \leq \inf(2s+m-2,k). \tag{2.7}$$

Using the binomial formula, we can write

$$g_1^q = (h+f_1)^q = h^q + f_1^q + \sum_{i=1}^{q-1}\binom{q}{i}h^i \cdot f_1^{q-i}, \tag{2.8}$$

$$g_2^q = (h+f_2)^q = h^q + f_2^q + \sum_{i=1}^{q-1}\binom{q}{i}h^i \cdot f_2^{q-i}. \tag{2.9}$$

Integrating (2.8) on the unit interval of $\mathbb{K}^2$ we get by using results similar to (2.3)

$$\sum_{i=1}^{\inf(2s+m-1,k)}\left[\sigma_{i-1,i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \beta_{s+m-} \\ \alpha_{s-} \end{array}\right]\times k} - 2 \cdot \sigma_{i-1,i-1,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right]\times k}\right] \cdot 2^{-qi} = 0\ for\ all\ q \geq 2. \tag{2.10}$$

From (2.10) we obtain

$$(2.11)\qquad \sigma_{i-1,i,i}^{\begin{bmatrix} s-1 \\ \underline{s+m-1} \\ \beta_{s+m-} \\ \alpha_{s-} \\ \\ \\ \end{bmatrix}\times k} = 2\cdot\sigma_{i-1,i-1,i}^{\begin{bmatrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \\ \end{bmatrix}\times k} \qquad for\ 0\leq i\leq \inf(2s+m-1,k).$$

Integrating (2.9) on the unit interval of $\mathbb{K}^2$ we get in the same way

(2.12)

$$\sum_{i=1}^{\inf(2s+m-1,k)}\left[\sigma_{i-1,i,i}^{\begin{bmatrix} s-1 \\ \underline{s+m-1} \\ \beta_{s+m-} \\ \alpha_{s-} \\ \\ \\ \end{bmatrix}\times k} - \sigma_{i-1,i,i}^{\begin{bmatrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \\ \end{bmatrix}\times k} - \sigma_{i-1,i-1,i}^{\begin{bmatrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \\ \end{bmatrix}\times k} + \sigma_{i-1,i-1,i}^{\begin{bmatrix} s-1 \\ \underline{s+m-1} \\ \beta_{s+m-} \\ \alpha_{s-} \\ \\ \\ \end{bmatrix}\times k}\right]\cdot 2^{-qi} = 0 \quad for\ q\geq 2.$$

From (2.12) we get

(2.13)

$$\sigma_{i-1,i,i}^{\begin{bmatrix} s-1 \\ \underline{s+m-1} \\ \beta_{s+m-} \\ \alpha_{s-} \\ \\ \\ \end{bmatrix}\times k} + \sigma_{i-1,i-1,i}^{\begin{bmatrix} s-1 \\ \underline{s+m-1} \\ \beta_{s+m-} \\ \alpha_{s-} \\ \\ \\ \end{bmatrix}\times k} = \sigma_{i-1,i,i}^{\begin{bmatrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \\ \end{bmatrix}\times k} + \sigma_{i-1,i-1,i}^{\begin{bmatrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \\ \end{bmatrix}\times k} \qquad for\ 1\leq i\leq \inf(2s+m-1,k).$$

Combining (2.5), (2.6), (2.7), (2.11) and (2.13) we obtain the following recurrent formula for the number of $(2s+m)\times k$ double persymmetric matrices of rank i

(2.14)

$$\Gamma_{i}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times k} = 2\cdot\Gamma_{i-1}^{\begin{bmatrix} s-1 \\ s-1+(m+1) \end{bmatrix}\times k} + 4\cdot\Gamma_{i-1}^{\begin{bmatrix} s \\ s+(m-1) \end{bmatrix}\times k} - 8\cdot\Gamma_{i-2}^{\begin{bmatrix} s-1 \\ s-1+m \end{bmatrix}\times k} + \Delta_{i}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times k} \quad for\ 0\leq i\leq \inf(2s+m,k)$$

where $\Delta_{i}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times k}$ is equal to

(2.15)

$$\sigma_{i,i,i}^{\begin{bmatrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \\ \end{bmatrix}\times k} - 3\cdot\sigma_{i-1,i-1,i-1}^{\begin{bmatrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \\ \end{bmatrix}\times k} + 2\cdot\sigma_{i-2,i-2,i-2}^{\begin{bmatrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \\ \end{bmatrix}\times k}.$$

We observe that we have for instance

$$2\cdot\Gamma_{i-1}^{\left[\begin{array}{c} s-1\\ s-1+(m+1)\end{array}\right]\times k} = \sigma_{i-1,i-1,i-1}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \beta_{s+m-}\\ \alpha_{s-}\\ \\ \\ \end{array}\right]\times k} + \sigma_{i-1,i-1,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \beta_{s+m-}\\ \alpha_{s-}\\ \\ \\ \end{array}\right]\times k} + \sigma_{i-2,i-1,i-1}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \beta_{s+m-}\\ \alpha_{s-}\\ \\ \\ \end{array}\right]\times k} + \sigma_{i-2,i-1,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \beta_{s+m-}\\ \alpha_{s-}\\ \\ \\ \end{array}\right]\times k} .$$

In Section 6, we study rank properties of a partition of double persymmetric matrices by integrating some appropriate exponential sums on the unit interval of $\mathbb{K}^2$ with integral equal to zero.

Consider the following partition of the matrix $D^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \\ \end{array}\right]\times k}(t,\eta)$,

$$\left(\begin{array}{cccccc|c}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & & \alpha_k\\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & & \alpha_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots\\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & & \alpha_{s+k-2}\\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & & \beta_k\\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & & \beta_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots\\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & & \beta_{k+m}\\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots\\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & & \beta_{s+m+k-2}\\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & & \alpha_{s+k-1}\\
\hline
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & & \beta_{s+m+k-1}
\end{array}\right).$$

In fact we prove that for all $j\in[0,\inf(2s+m-2,k-1)]$ we have

(2.16)

$$\#\left(\begin{array}{c|c} j & j\\ \hline j & j\\ \hline j & j+1\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} = \#\left(\begin{array}{c|c} j & j\\ \hline j & j\\ \hline j & j\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}},$$

(2.17)

$$\#\left(\begin{array}{c|c} j & j\\ \hline j & j+1\\ \hline j & j+1\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} = \#\left(\begin{array}{c|c} j & j\\ \hline j & j\\ \hline j & j\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} + \#\left(\begin{array}{c|c} j & j\\ \hline j & j\\ \hline j & j+1\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}.$$

We deduce (2.16) from the fact that for all $q\geq 2$ the integral $\int_{\mathbb{P}\times\mathbb{P}}\phi(t,\eta)\cdot\theta_3^{q-1}(t,\eta)dtd\eta$ is equal to zero, that is

$$\sum_{j=0}^{\inf(2s+m-2,k-1)} a(s,m,k,q)\cdot 2^{-jq}\cdot\left[\#\left(\begin{array}{c|c} j & j\\ \hline j & j\\ \hline j & j\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} - \#\left(\begin{array}{c|c} j & j\\ \hline j & j\\ \hline j & j+1\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}\right]=0.$$

We deduce (2.17) in a similar way from the fact that for all $q \geq 2$ the integral $\int_{\mathbb{P}\times\mathbb{P}} \theta_1(t,\eta)\cdot\theta_2^{q-1}(t,\eta)dtd\eta$ is equal to zero, that is

$$\sum_{j=0}^{\inf(2s+m-2,k-1)} b(s,m,k,q)\cdot 2^{-jq}\cdot\Bigg[\#\left(\begin{array}{c|c} j & j\\ \hline j & j\\ \hline j & j\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} + \#\left(\begin{array}{c|c} j & j\\ \hline j & j\\ \hline j & j+1\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}$$
$$-\#\left(\begin{array}{c|c} j & j\\ \hline j & j+1\\ \hline j & j+1\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}\Bigg] = 0.$$

In Section 7, we study some rank properties of submatrices of double persymmetric matrices. Consider the following partition of the matrix $D^{\left[\begin{smallmatrix} s-1\\ \frac{s+m-1}{\alpha_{s-}}\\ \beta_{s+m-}\end{smallmatrix}\right]\times k}(t,\eta)$,

$$\left(\begin{array}{ccccc|c}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k\\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2}\\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k\\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2}\\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1}\\
\hline
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{array}\right).$$

In fact we prove that for all $j \in [0, \inf(2s+m-3, k-2)]$ we have

$$\#\left(\begin{array}{c|c} j & j+1\\ \hline j & j+1\\ \hline j & j+1\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} = 0. \tag{2.18}$$

We prove (2.18) by contradiction.
Assume on the contrary that there exist $j_0 \in [0, \inf(2s+m-3, k-2)]$ such that

$$\#\left(\begin{array}{c|c} j_0 & j_0+1\\ \hline j_0 & j_0+1\\ \hline j_0 & j_0+1\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} > 0.$$

We show that

$$\#\left(\begin{array}{c|c} j_0 & j_0+1 \\ \hline j_0 & j_0+1 \\ \hline j_0 & j_0+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} > 0 \Longrightarrow \#\left(\begin{array}{c|c} j_0-1 & j_0 \\ \hline j_0-1 & j_0 \\ \hline j_0-1 & j_0 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}_1/\mathbb{P}_{k+s-1}\times\mathbb{P}_1/\mathbb{P}_{k+s+m-1}} > 0$$

$$\Longrightarrow \ldots \Longrightarrow \#\left(\begin{array}{c|c} 0 & 1 \\ \hline 0 & 1 \\ \hline 0 & 1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}_{j_0}/\mathbb{P}_{k+s-1}\times\mathbb{P}_{j_0}/\mathbb{P}_{k+s+m-1}} > 0$$

which obviously contradicts

$$\#\left(\begin{array}{c|c} 0 & 1 \\ \hline 0 & 1 \\ \hline 0 & 1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}_{j_0}/\mathbb{P}_{k+s-1}\times\mathbb{P}_{j_0}/\mathbb{P}_{k+s+m-1}} = 0.$$

These inequalities are a consequence of rank properties of submatrices of double persymmetric matrices.

In Section 8, we study the remainder $\Delta_j^{\left[\substack{s\\ s+m}\right]\times k}$ in the recurrent formula and we prove in particular that

$$\Delta_j^{\left[\substack{s\\ s+m}\right]\times (k+1)} - \Delta_j^{\left[\substack{s\\ s+m}\right]\times k} = 0 \text{ if } k \geq j+1. \tag{2.19}$$

Consider still the following partition of the matrix $D^{\left[\substack{\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}}\right]\times k}(t,\eta)$

$$\left(\begin{array}{ccccc|c}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1} \\
\hline
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{array}\right).$$

Let $j \leq \inf(k-2, 2s+m-3)$. Then by elementary rank considerations and using (2.18) we obtain

(2.20)

$$\begin{aligned}\sigma_{j,j,j}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right] \times k} &= {}^{\#}\left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} + {}^{\#}\left(\begin{array}{c|c} j-1 & j \\ \hline j & j \\ \hline j & j \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} \\ &+ {}^{\#}\left(\begin{array}{c|c} j-1 & j \\ \hline j-1 & j \\ \hline j & j \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} + {}^{\#}\left(\begin{array}{c|c} j-1 & j \\ \hline j-1 & j \\ \hline j-1 & j \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} \\ &= {}^{\#}\left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}.\end{aligned}$$

Further by (2.16), (2.17) and (2.18) we have

(2.21)

$$\begin{aligned}4\cdot\sigma_{j,j,j}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right] \times (k-1)} &= {}^{\#}\left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} + {}^{\#}\left(\begin{array}{c|c} j & j \\ \hline j & j+1 \\ \hline j & j+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} \\ &+ {}^{\#}\left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} + {}^{\#}\left(\begin{array}{c|c} j & j+1 \\ \hline j & j+1 \\ \hline j & j+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} \\ &= 4\cdot{}^{\#}\left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}.\end{aligned}$$

From (2.20), (2.21) we deduce

$$\sigma_{j,j,j}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right] \times k} = \sigma_{j,j,j}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right] \times (j+1)} \quad \text{if } \ j \leq \inf(k-2, 2s+m-3). \tag{2.22}$$

Using (2.16), (2.17) we obtain easily

$$\sigma_{j,j,j}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right] \times (j+1)} = {}^{\#}\left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{j+s}\times\mathbb{P}/\mathbb{P}_{j+s+m}}, \tag{2.23}$$

$$
\begin{aligned}
(2.24)\qquad 4\cdot\sigma_{j,j,j}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times j} &= 4\cdot 4\cdot\Gamma_{j}^{\left[\substack{s-1\\ s-1+m}\right]\times j}\\
&= 4\cdot {}^{\#}\left(\begin{array}{c|c} j & j\\ \hline j & j\\ \hline j & j \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{j+s}\times\mathbb{P}/\mathbb{P}_{j+s+m}} + 4\cdot\Gamma_{j+1}^{\left[\substack{s-1\\ s-1+m}\right]\times(j+1)}.
\end{aligned}
$$

From (2.23), (2.24) we deduce Theorem 3.3.

Recalling that $\Delta_{i}^{\left[\substack{s\\ s+m}\right]\times k}$, is equal to

$$
\sigma_{i,i,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k} - 3\cdot\sigma_{i-1,i-1,i-1}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k} + 2\cdot\sigma_{i-2,i-2,i-2}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}
$$

we deduce easily Theorem 3.4.

We get (2.19) from Theorem 3.5.

In Section 9, we establish a recurrent formula for the difference

$$
\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}\qquad for\ 2\leq i\leq 2s+m,\quad k\geq i+1.
$$

In fact we prove that

(2.25)

$$
\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k} = 4\cdot\left[\Gamma_{i-1}^{\left[\substack{s\\ s+(m-1)}\right]\times(k+1)} - \Gamma_{i-1}^{\left[\substack{s\\ s+(m-1)}\right]\times k}\right]\quad \text{if}\quad 2\leq i\leq s-1,\quad k\geq i+1,
$$

(2.26)

$$
\begin{aligned}
&\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k} = 4\cdot\left[\Gamma_{s+j-1}^{\left[\substack{s\\ s+(m-1)}\right]\times(k+1)} - \Gamma_{s+j-1}^{\left[\substack{s\\ s+(m-1)}\right]\times k}\right] + 2^{s-1}\left[\Gamma_{j+1}^{\left[\substack{1\\ 1+(s+m-1)}\right]\times(k+1)} - \Gamma_{j+1}^{\left[\substack{1\\ 1+(s+m-1)}\right]\times k}\right]\\
&- 2^{s+1}\left[\Gamma_{j}^{\left[\substack{1\\ 1+(s+m-2)}\right]\times(k+1)} - \Gamma_{j}^{\left[\substack{1\\ 1+(s+m-2)}\right]\times k}\right]\text{if } 0\leq j\leq s+m,\quad k\geq s+j+1.
\end{aligned}
$$

To prove (2.25) we proceed as follows

Set for $1\leq i\leq \inf(2s+m,k)$

$$
\Omega_i(s,m,k) = \Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k} - 4\cdot\Gamma_{i-1}^{\left[\substack{s\\ s+(m-1)}\right]\times k}.
$$

Then, from the recurrent formula (2.14) we get

$$\Omega_i(s,m,k) = 2\cdot\Omega_{i-1}(s-1,m+1,k) + \Delta_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}. \tag{2.27}$$

Applying succesively (2.27) we obtain

(2.28)

$$\begin{aligned}\sum_{j=0}^{i-2} 2^j\cdot\Omega_{i-j}(s-j,m+j,k) &= \sum_{j=0}^{i-2} 2^{j+1}\cdot\Omega_{i-(j+1)}(s-(j+1),m+(j+1),k) + \sum_{j=0}^{i-2} 2^j\cdot\Delta_{i-j}^{\left[\begin{smallmatrix} s-j \\ s-j+(m+j) \end{smallmatrix}\right]\times k} \\ &= \sum_{j=1}^{i-1} 2^j\cdot\Omega_{i-j}(s-j,m+j,k) + \sum_{j=0}^{i-2} 2^j\cdot\Delta_{i-j}^{\left[\begin{smallmatrix} s-j \\ s-j+(m+j) \end{smallmatrix}\right]\times k}.\end{aligned}$$

Observing that

$$\Omega_1(s-(i-1),m+(i-1),k) = \Gamma_1^{\left[\begin{smallmatrix} s-i+1 \\ s+m \end{smallmatrix}\right]\times k} - 4\cdot\Gamma_0^{\left[\begin{smallmatrix} s-i+1 \\ s+(m-1) \end{smallmatrix}\right]\times k} = 9-4=5,$$

we get by (2.28) after some simplifications

(2.29)

$$\Omega_i(s,m,k) = 2^{i-1}\cdot\Omega_1(s-(i-1),m+(i-1),k) + \sum_{j=0}^{i-2} 2^j\cdot\Delta_{i-j}^{\left[\begin{smallmatrix} s-j \\ s-j+(m+j) \end{smallmatrix}\right]\times k} = 5\cdot 2^{i-1} + \sum_{j=0}^{i-2} 2^j\Delta_{i-j}^{\left[\begin{smallmatrix} s-j \\ s-j+(m+j) \end{smallmatrix}\right]\times k}.$$

Applying (2.19) and (2.29) we obtain (2.25).

We prove (2.26) in a similar way.

In Section 10, we compute explicitly

$$2^{s-1}\left[\Gamma_{j+1}^{\left[\begin{smallmatrix} 1 \\ 1+(s+m-1) \end{smallmatrix}\right]\times(k+1)} - \Gamma_{j+1}^{\left[\begin{smallmatrix} 1 \\ 1+(s+m-1) \end{smallmatrix}\right]\times k}\right] - 2^{s+1}\left[\Gamma_{j}^{\left[\begin{smallmatrix} 1 \\ 1+(s+m-2) \end{smallmatrix}\right]\times(k+1)} - \Gamma_{j}^{\left[\begin{smallmatrix} 1 \\ 1+(s+m-2) \end{smallmatrix}\right]\times k}\right] \overset{\text{def}}{=}$$
$R(j,s,m,k)$
if $0 \leq j \leq s+m, \quad k \geq s+j+1.$
In fact we prove

$$R(j,s,m,k) = \begin{cases} 2^{k+s-1} & \text{if } j=0, \quad k\geq s+1, \\ -2^{k+s-1} & \text{if } j=1, \quad k\geq s+2, \\ 0 & \text{if } 2\leq j\leq s+m-1, \quad k\geq s+j+1, \\ -3\cdot 2^{2k+2s+m-2} & \text{if } j=s+m, \quad k\geq 2s+m+1. \end{cases} \tag{2.30}$$

From (2.25), (2.26) and (2.30) we get for $s\geq 2, m\geq 0$

(2.31)

$$\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(k+1)}-\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}=4\cdot\left[\Gamma_{i-1}^{\left[\substack{s\\ s+(m-1)}\right]\times(k+1)}-\Gamma_{i-1}^{\left[\substack{s\\ s+(m-1)}\right]\times k}\right]\text{if } 1\leq i\leq s-1,\quad k\geq i+1,$$

(2.32)

$$\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times(k+1)}-\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k}=4\cdot\left[\Gamma_{s+j-1}^{\left[\substack{s\\ s+(m-1)}\right]\times(k+1)}-\Gamma_{s+j-1}^{\left[\substack{s\\ s+(m-1)}\right]\times k}\right]$$
$$+\begin{cases}2^{k+s-1} & \text{if } j=0,\quad k\geq s+1,\\ -2^{k+s-1} & \text{if } j=1,\quad k\geq s+2,\\ 0 & \text{if } 2\leq j\leq s+m-1,\quad k\geq s+j+1,\\ -3\cdot 2^{2k+2s+m-2} & \text{if } j=s+m,\quad k\geq 2s+m+1.\end{cases}$$

We establish (2.30) by applying the formula for $\Gamma_{i}^{\left[\substack{1\\ 1+m}\right]\times k}$ (obtained in [4], see Theorem 3.9, with $m\rightarrow s+m-1, m\rightarrow s+m-2$).

In Section 11, we establish a formula for $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}$ in the case $1\leq i\leq \inf(s-1,k-1),\quad m\geq 0.$
We show precisely

$$\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}=\begin{cases}1 & \text{if } i=0,\\ \Gamma_{i}^{\left[\substack{i\\ i}\right]\times(i+1)}\overset{\text{def}}{=}\gamma_i & \text{if } 1\leq i\leq \inf(s-1,k-1), m\geq 0\end{cases}\tag{2.33}$$

and

$$\Gamma_{i}^{\left[\substack{i\\ i}\right]\times(i+1)}\overset{\text{def}}{=}\gamma_i=21\cdot 2^{3i-4}-3\cdot 2^{2i-3}\text{ if } 1\leq i\leq \inf(s-1,k-1)\ m\geq 0.\tag{2.34}$$

From (2.33), (2.34) we deduce the following results

$$\Gamma_{k}^{\left[\substack{s\\ s+m}\right]\times k}=2^{2k+2s+m-2}-3\cdot 2^{3k-4}+2^{2k-3}\quad \textit{for } 1\leq k\leq s,\tag{2.35}$$

$$\sum_{j=0}^{s-1}\Gamma_{j}^{\left[\substack{s\\ s+m}\right]\times k}=3\cdot 2^{3s-4}-2^{2s-3}\quad \textit{for } k\geq s,\tag{2.36}$$

$$\sum_{j=0}^{s-1}\Gamma_{j}^{\left[\substack{s\\ s+m}\right]\times k}\cdot 2^{-j}=7\cdot 2^{2s-4}-3\cdot 2^{s-3}\quad \textit{for } k\geq s.\tag{2.37}$$

We prove (2.33) by induction on j.

Set

$$(2.38)\qquad (H_j)\quad \Gamma_{j}^{\left[\substack{s\\ s+m}\right]\times k} = \Gamma_{j}^{\left[\substack{j\\ j}\right]\times (j+1)} = \gamma_j \quad \text{if} \quad s \geq j+1,\ k \geq j+1,\ m \geq 0.$$

Applying (2.1) we get, using successively (2.31) for i = 1, 2, 3

$$(2.39)\qquad \Gamma_{1}^{\left[\substack{s\\ s+m}\right]\times k} = \Gamma_{1}^{\left[\substack{1\\ 1}\right]\times 2} = \gamma_1 \quad \text{if} \quad s \geq 2,\ k \geq 2,\ m \geq 0\ .$$

$$(2.40)\qquad \text{Using (2.39) we get} \quad \Gamma_{2}^{\left[\substack{s\\ s+m}\right]\times k} = \Gamma_{2}^{\left[\substack{2\\ 2}\right]\times 3} = \gamma_2 \quad \text{if} \quad s \geq 3,\ k \geq 3,\ m \geq 0\ .$$

(2.41)

$$\text{Using (2.39),(2.40) we get} \quad \Gamma_{3}^{\left[\substack{s\\ s+m}\right]\times k} = \Gamma_{3}^{\left[\substack{3\\ 3}\right]\times 4} = \gamma_3 \quad \text{if} \quad s \geq 4,\ k \geq 4,\ m \geq 0\ .$$

Thus (H_j) [see (2.38)] holds for $j \in \{1, 2, 3\}$.
Assume now that (H_j) holds for $j \in \{1, 2, 3, \ldots, i-1\}$, that is

(2.42)

$$\bigwedge_{j=1}^{j=i-1} (H_j) \qquad \Gamma_{j}^{\left[\substack{s\\ s+m}\right]\times k} = \Gamma_{j}^{\left[\substack{j\\ j}\right]\times (j+1)} = \gamma_j \quad \text{for all} \quad 1 \leq j \leq i-1,\ k \geq j+1,\ s \geq j+1\ m \geq 0.$$

From $\bigwedge_{j=1}^{j=i-1}(H_j)$ and (2.31) it follows
$\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times (k+1)} - \Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k} = 4 \cdot \left[\Gamma_{i-1}^{\left[\substack{s\\ s+(m-1)}\right]\times (k+1)} - \Gamma_{i-1}^{\left[\substack{s\\ s+(m-1)}\right]\times k}\right] = 0$ for all $s \geq i+1,\ k \geq i+1$.
Hence $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k} = \Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times (i+1)}$ for all $s \geq i+1,\ k \geq i+1$.
To get (2.33) we need only to show that $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times (i+1)}$ is equal to $\Gamma_{i}^{\left[\substack{i\\ i}\right]\times (i+1)}$ for all $s \geq i+1,\ m \geq 0$.
To do so we consider the following matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_i & \alpha_{i+1} \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{i+1} & \alpha_{i+2} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+i-2} & \alpha_{s+i-1} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+i-1} & \alpha_{s+i} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_i & \beta_{i+1} \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{i+1} & \beta_{i+2} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{i+m} & \beta_{i+m+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+i-2} & \beta_{s+m+i-1} \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+i-1} & \beta_{s+m+i}
\end{pmatrix}.$$

Using the equations (2.1) with k = i+1 we obtain

$$(2.43) \qquad \sum_{j=0}^{i+1} \Gamma_j^{\left[\substack{s \\ s+m}\right]\times(i+1)} = 2^{2i+2s+m+2}$$

and

$$(2.44) \qquad \sum_{j=0}^{i+1} \Gamma_j^{\left[\substack{s \\ s+m}\right]\times(i+1)} \cdot 2^{-j} = 2^{i+2s+m-1} + 2^{2i} - 2^{i-1}.$$

Setting s = i, m = 0 in (2.43), (2.44) we have

$$(2.45) \qquad \sum_{j=0}^{i+1} \Gamma_j^{\left[\substack{i \\ i}\right]\times(i+1)} = 2^{4i}$$

and

$$(2.46) \qquad \sum_{j=0}^{i+1} \Gamma_j^{\left[\substack{i \\ i}\right]\times(i+1)} \cdot 2^{-j} = 2^{3i-1} + 2^{2i} - 2^{i-1}.$$

From (2.43), (2.44), (2.45), (2.46) and $\bigwedge_{j=1}^{j=i-1}(H_j)$ we deduce (2.34).

In Section 12, using successively the recurrent formula (2.32) for the difference $\Gamma_{s+j}^{\left[\substack{s \\ s+j}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\substack{s \\ s+j}\right]\times k}$,
with $0 \leq j \leq s+m,\ k \geq s+j+1$ we deduce
In the case $m \in \{0,1\}$

$$(2.47) \quad \Gamma_{s+j}^{\left[\substack{s \\ s}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\substack{s \\ s}\right]\times k} = \begin{cases} 3 \cdot 2^{k+s-1} & \text{if } j = 0,\ k > s, \\ 21 \cdot 2^{k+s+3j-4} & \text{if } 1 \leq j \leq s-1,\ k > s+j, \\ 3 \cdot 2^{2k+2s-2} - 3 \cdot 2^{k+4s-4} & \text{if } j = s,\ k > 2s, \end{cases}$$

(2.48)

$$\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix}\right]\times k} = \begin{cases} 2^{k+s-1} & \text{if } j=0,\ k>s, \\ 11\cdot 2^{k+s-1} & \text{if } j=1,\ k>s+1, \\ 21\cdot 2^{k+s+3j-5} & \text{if } 2\leq j\leq s,\ k>s+j, \\ 3\cdot 2^{2k+2s-1}-3\cdot 2^{k+4s-2} & \text{if } j=s+1,\ k>2s+1. \end{cases}$$

In the case $m \geq 2$

$$\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = \begin{cases} 2^{k+s-1} & \text{if } j=0,\ k>s, \\ 3\cdot 2^{k+s+2j-3} & \text{if } 1\leq j\leq m-1,\ k>s+j, \\ 11\cdot 2^{k+s+2m-3} & \text{if } j=m,\ k>s+m, \end{cases} \tag{2.49}$$

(2.50)

$$\Gamma_{s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = \begin{cases} 21\cdot 2^{k+s+2m+3j-4} & \text{if } 1\leq j\leq s-1,\ k>s+m+j, \\ 3\cdot 2^{2k+2s+m-2}-3\cdot 2^{k+4s+2m-4} & \text{if } j=s,\ k>2s+m. \end{cases}$$

For instance to compute $\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times k}$ for $0 \leq j \leq s-1,\ k \geq s+j+1$ we proceed as follows :

From (2.32) with m = 0, j = 0, $k \geq s+1$ we get, using (2.33)

(2.51)

$$\Gamma_{s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times k} = 4\cdot\left[\Gamma_{s-1}^{\left[\begin{smallmatrix} s-1 \\ s-1+1 \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s-1}^{\left[\begin{smallmatrix} s-1 \\ s-1+1 \end{smallmatrix}\right]\times k}\right] + 2^{k+s-1},$$

(2.52)

$$\Gamma_{s-1}^{\left[\begin{smallmatrix} s-1 \\ s-1+1 \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s-1}^{\left[\begin{smallmatrix} s-1 \\ s-1+1 \end{smallmatrix}\right]\times k} = 4\cdot\left[\Gamma_{s-2}^{\left[\begin{smallmatrix} s-1 \\ s-1 \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s-2}^{\left[\begin{smallmatrix} s-1 \\ s-1 \end{smallmatrix}\right]\times k}\right] + 2^{k+s-2} = 2^{k+s-2}.$$

From (2.51), (2.52) we obtain

$$\Gamma_{s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times k} = 4\cdot 2^{k+s-2} + 2^{k+s-1} = 3\cdot 2^{k+s-1}. \tag{2.53}$$

We get in a similar way, using (2.53) with $s \to s-1$

$$\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times k} = 21\cdot 2^{k+s-1}. \tag{2.54}$$

Let $2 \leq j \leq s-1,\ k > s+j$.

By (2.32) we obtain

$$\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times k} = 4\cdot\left[\Gamma_{s+j-1}^{\left[\begin{smallmatrix} s-1 \\ (s-1)+1 \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s-1+j}^{\left[\begin{smallmatrix} s-1 \\ (s-1)+1 \end{smallmatrix}\right]\times k}\right] \quad \text{if } 2\leq j\leq s-1,$$

$$\Gamma_{s-1+j}^{\left[\begin{smallmatrix} s-1\\ (s-1)+1\end{smallmatrix}\right]\times(k+1)} - \Gamma_{s-1+j}^{\left[\begin{smallmatrix} s-1\\ (s-1)+1\end{smallmatrix}\right]\times k} = 4\cdot\left[\Gamma_{s-1+(j-1)}^{\left[\begin{smallmatrix} s-1\\ s-1\end{smallmatrix}\right]\times(k+1)} - \Gamma_{s-1+(j-1)}^{\left[\begin{smallmatrix} s-1\\ s-1\end{smallmatrix}\right]\times k}\right] \qquad \text{if } 2\leq j\leq s-1+1-1=s-1.$$

From the above equations we get

$$\Gamma_{s+j}^{\left[\begin{smallmatrix} s\\ s\end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\begin{smallmatrix} s\\ s\end{smallmatrix}\right]\times k} = 4^2\cdot\left[\Gamma_{s-1+(j-1)}^{\left[\begin{smallmatrix} s-1\\ s-1\end{smallmatrix}\right]\times(k+1)} - \Gamma_{s-1+(j-1)}^{\left[\begin{smallmatrix} s-1\\ s-1\end{smallmatrix}\right]\times k}\right] \quad if\ 2\leq j\leq s-1. \tag{2.55}$$

Using successively (2.55) we get from (2.54) with $s\rightarrow s-j+1$

(2.56)

$$\begin{aligned}\Gamma_{s+j}^{\left[\begin{smallmatrix} s\\ s\end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\begin{smallmatrix} s\\ s\end{smallmatrix}\right]\times k} &= 4^{2(j-1)}\cdot\left[\Gamma_{s-j+2}^{\left[\begin{smallmatrix} s-j+1)\\ s-j+1)\end{smallmatrix}\right]\times(k+1)} - \Gamma_{s-j+2}^{\left[\begin{smallmatrix} s-j+1\\ s-j+1\end{smallmatrix}\right]\times k}\right] \quad if\ 2\leq j\leq s-1\\ &= 4^{2(j-1)}\cdot 21\cdot 2^{k+s-j+1-1} = 21\cdot 2^{k+s+3j-4}.\end{aligned}$$

In Sections 13, 14, 15, 16 and 17 we compute $\Gamma_{s+j}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k}$ for $0\leq j\leq s+m,\ k\geq s+j,\ m\geq 0$.

To do so, consider the following array $\left(\Gamma_{s+j}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k}\right)_{0\leq j\leq s+m,\ k\geq s+j}$:

(2.57)

$$\begin{array}{cccccccc}
\Gamma_{s}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times s} & \Gamma_{s}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+1)} & \Gamma_{s}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+2)} & \dots & \Gamma_{s}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k} & \dots & k\geq s\\
\Gamma_{s+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+1)} & \Gamma_{s+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+2)} & \Gamma_{s+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+3)} & \dots & \Gamma_{s+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k} & \dots & k\geq s+1\\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots\\
\Gamma_{s+j}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+j)} & \Gamma_{s+j}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+j+1)} & \Gamma_{s+j}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+j+2)} & \dots & \Gamma_{s+j}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k} & \dots & k\geq s+j\\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots\\
\Gamma_{s+m}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+m)} & \Gamma_{s+m}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+m+1)} & \Gamma_{s+m}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+m+2)} & \dots & \Gamma_{s+m}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k} & \dots & k\geq s+m\\
\Gamma_{s+m+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+m+1)} & \Gamma_{s+m+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+m+2)} & \Gamma_{s+m+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+m+3)} & \dots & \Gamma_{s+m+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k} & \dots & k\geq s+m+1\\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots\\
\Gamma_{s+m+1+j}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+m+1+j)} & \Gamma_{s+m+1+j}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+m+2+j)} & \Gamma_{s+m+1+j}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+m+3+j)} & \dots & \Gamma_{s+m+1+j}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k} & \dots & k\geq s+m+1+j\\
\Gamma_{2s+m-1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(2s+m-1)} & \Gamma_{2s+m-1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(2s+m)} & \Gamma_{2s+m-1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(2s+m+1)} & \dots & \Gamma_{2s+m-1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k} & \dots & k\geq 2s+m-1\\
\Gamma_{2s+m}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(2s+m)} & \Gamma_{2s+m}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(2s+m+1)} & \Gamma_{2s+m}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(2s+m+2)} & \dots & \Gamma_{2s+m}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k} & \dots & k\geq 2s+m.
\end{array}$$

- To compute the elements in the array $\left(\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times k}\right)_{0\leq j\leq s+m,\ k\geq s+j}$, we proceed by induction on j as follows:
  Let l be a rational integer such that $0\leq l\leq s+m-1$.
  Assume that we have computed the elements in the following subarray $\left(\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times k}\right)_{0\leq j\leq l,\ k\geq s+j}$.

  Recall that from (2.33), (2.34) $\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times k}$ are known for $0\leq i\leq s-1,\ k\geq i$.

  From the first equation in (2.1) with k = s+l+1, $\sum_{i=0}^{s+l+1}\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times(s+l+1)} = 2^{4s+2l+m}$
  we deduce $\Gamma_{s+l+1}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times(s+l+1)}$ since the terms $\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times(s+l+1)}$ for $i\leq s+l$ are known.
  From the equations in (2.1) with $k=s+l+2$ we get

$$\begin{cases}\sum_{i=0}^{s+l+2}\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times(s+l+2)} = 2^{4s+2l+m+2},\\ \sum_{i=0}^{s+l+2}\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times(s+l+2)}\cdot 2^{-i} = 2^{3s+m+l}+2^{2s+2l+2}-2^{s+l}.\end{cases}$$

  We deduce $\Gamma_{s+l+1}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times(s+1+2)}$ and $\Gamma_{s+l+2}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times(s+l+2)}$ since the terms $\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times(s+l+2)}$ for $i\leq s+l$ are known.

  Then, from (2.49) or (2.50), in the case $m\geq 2$, we compute $\Gamma_{s+l+1}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times k}$ knowing $\Gamma_{s+l+1}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times(s+1+2)}$ for all $k\geq s+l+2$.
  We have now computed any element in the $l+1$-th row and the first element in the $l+2$-th row in the array $\left(\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times k}\right)_{0\leq j\leq s+m,\ k\geq s+j}$.

- From the first equation in (2.1) with k = s, $\sum_{i=0}^{s}\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times s} = 2^{4s+m-2}$
  we deduce $\Gamma_{s}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times s}$ since the terms $\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times s}$ for $i\leq s-1$ are known by (2.33), (2.34).
  From the equations in (2.1) with k = s+1 we obtain

$$\begin{cases}\sum_{i=0}^{s+1}\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times(s+1)} = 2^{4s+m},\\ \sum_{i=0}^{s+1}\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times(s+1)}\cdot 2^{-i} = 2^{3s+m-1}+2^{2s}-2^{s-1}.\end{cases}$$

  We deduce $\Gamma_{s}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times(s+1)}$ and $\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times(s+1)}$ since the terms $\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times(s+1)}$ for $i\leq s-1$ are known.

  Then, from (2.49) in the case $m\geq 2$ with j = 0 we compute $\Gamma_{s}^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times k}$ for all $k\geq s+1$.

We have now computed any element in the subarray $\left(\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k}\right)_{0\leq j\leq l,\ k\geq s+j}$ with $l=0$.

- From the equations in (2.1) with k = s+ 2, we have

$$\begin{cases}\sum_{i=0}^{s+2}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(s+2)} = 2^{4s+m+2},\\ \sum_{i=0}^{s+2}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(s+2)}\cdot 2^{-i} = 2^{3s+m}+2^{2s+2}-2^{s}.\end{cases}$$

  We deduce $\Gamma_{s+1}^{\left[\substack{s\\ s+m}\right]\times(s+2)}$ and $\Gamma_{s+2}^{\left[\substack{s\\ s+m}\right]\times(s+2)}$ since the terms $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(s+2)}$ for $i\leq s$ are known.

  Then from (2.49) in the case $m\geq 2$ with j = 1 we compute $\Gamma_{s+1}^{\left[\substack{s\\ s+m}\right]\times k}$ knowing $\Gamma_{s+1}^{\left[\substack{s\\ s+m}\right]\times(s+2)}$ for all $k\geq s+2$.

  We have now computed any element in the subarray $\left(\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k}\right)_{0\leq j\leq l,\ k\geq s+j}$ with $l=1$.

- From the equations in (2.1) with k = s+ 3, we get

$$\begin{cases}\sum_{i=0}^{s+3}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(s+3)} = 2^{4s+m+4},\\ \sum_{i=0}^{s+3}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(s+3)}\cdot 2^{-i} = 2^{3s+m+1}+2^{2s+4}-2^{s+1}.\end{cases}$$

  We deduce $\Gamma_{s+2}^{\left[\substack{s\\ s+m}\right]\times(s+3)}$ and $\Gamma_{s+3}^{\left[\substack{s\\ s+m}\right]\times(s+3)}$ since the terms $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(s+3)}$ for $i\leq s+1$ are known.

  Then from (2.46) with j = 2 we compute $\Gamma_{s+2}^{\left[\substack{s\\ s+m}\right]\times k}$ knowing $\Gamma_{s+2}^{\left[\substack{s\\ s+m}\right]\times(s+3)}$ for all $k\geq s+3$.

  We have now computed any element in the subarray $\left(\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k}\right)_{0\leq j\leq 2,\ k\geq s+j}$ with $l=2$.

  Further we deduce that

$$\Gamma_{s+2}^{\left[\substack{s\\ s+m}\right]\times k} = 8\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+m-1}\right]\times(k-1)}\quad\text{for } k\geq s+2,\ m\geq 2. \tag{2.58}$$

- To compute the elements in the subarray $\left(\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k}\right)_{2\leq j\leq m,\ k\geq s+j}$ we proceed by induction on j.

  To do so, let l be a rational integer such that $2\leq l\leq m$.

  Assume (H_{l-1}) $\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k} = 8^{j-1}\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+(m-(j-1))}\right]\times(k-(j-1))}$ for $1\leq j\leq l-1,\ k\geq s+j$.

  (H_{l-1}) holds for $l=2$ and for $l=3$ by (2.58).

To establish (H_l) we need only to show that

$$\Gamma_{s+l}^{\left[\substack{s\\s+m}\right]\times k} = 8^{l-1}\cdot\Gamma_{s+1}^{\left[\substack{s\\s+(m-(l-1))}\right]\times(k-(l-1))} \quad \text{for} \quad k\geq s+l. \tag{2.59}$$

In the case $k>s+l$ we proceed as follows:

From the equations in (2.1) with $k=s+l+1$, we have

$$\begin{cases}\sum_{i=0}^{s+l+1}\Gamma_{i}^{\left[\substack{s\\s+m}\right]\times(s+l+1)} = 2^{4s+m+2l},\\ \sum_{i=0}^{s+l+1}\Gamma_{i}^{\left[\substack{s\\s+m}\right]\times(s+l+1)}\cdot 2^{-i} = 2^{3s+m+l-1}+2^{2s+2l}-2^{s+l-1}.\end{cases}$$

We deduce $\Gamma_{s+l}^{\left[\substack{s\\s+m}\right]\times(s+l+1)}$ and $\Gamma_{s+l+1}^{\left[\substack{s\\s+m}\right]\times(s+l+1)}$ since by (H_{l-1}) the terms $\Gamma_{i}^{\left[\substack{s\\s+m}\right]\times(s+l+1)}$ for $i\leq s+l-1$ are known.

Then from (2.49) with $j=l$ we compute $\Gamma_{s+l}^{\left[\substack{s\\s+m}\right]\times k}$ for all $k\geq s+l+1$ and (2.59) follows.

The case $k=s+l$ is obtained in a similar way.

- To compute the elements in the remaining subarray $\left(\Gamma_{s+m+1+j}^{\left[\substack{s\\s+m}\right]\times k}\right)_{0\leq j\leq s-1,\ k\geq s+m+1+j}$ we proceed still by induction on j.

Let j = 0, a similar proof of (2.58) gives the following identity:

$$\Gamma_{s+m+1}^{\left[\substack{s\\s+m}\right]\times k} = 8^{m}\cdot\Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(k-m)} \quad \text{for} \quad k\geq s+m+1. \tag{2.60}$$

Let l be a rational integer such that $1\leq l\leq s-1$.

Assume (H_{l-1}) $\Gamma_{s+m+1+j}^{\left[\substack{s\\s+m}\right]\times k} = 8^{2j+m}\cdot\Gamma_{s-j+1}^{\left[\substack{s-j\\s-j}\right]\times(k-m-2j)}$ for $0\leq j\leq l-1,\ k\geq s+m+1+j$.

From (2.60) (H_{l-1}) holds for $l=1$.

To establish (H_l) we need only to show that

$$\Gamma_{s+m+1+l}^{\left[\substack{s\\s+m}\right]\times k} = 8^{2l+m}\cdot\Gamma_{s-l+1}^{\left[\substack{s-l\\s-l}\right]\times(k-(l-1))} \quad \text{for} \quad k\geq s+m+1+l. \tag{2.61}$$

In the case $k>s+m+1+l$ we proceed as follows :

From the equations in (2.1) with $k=s+m+l+2$, we get

$$\begin{cases}\sum_{i=0}^{s+m+l+2}\Gamma_{i}^{\left[\substack{s\\s+m}\right]\times(s+m+l+2)} = 2^{4s+3m+2l+2},\\ \sum_{i=0}^{s+m+l+2}\Gamma_{i}^{\left[\substack{s\\s+m}\right]\times(s+m+l+2)}\cdot 2^{-i} = 2^{3s+2m+l}+2^{2s+2m+2l+2}-2^{s+m+l}.\end{cases}$$

We deduce $\Gamma_{s+m+l+1}^{\left[\substack{s\\s+m}\right]\times(s+m+l+2)}$ and $\Gamma_{s+m+l+2}^{\left[\substack{s\\s+m}\right]\times(s+m+l+2)}$ since by (H_{l-1}) the terms $\Gamma_{i}^{\left[\substack{s\\s+m}\right]\times(s+m+l+2)}$ for $i\leq s+m+l$ are known.

Then from (2.50) with $j = l$ knowing $\Gamma_{s+m+l+1}^{\left[\substack{s\\ s+m}\right]\times(s+m+l+2)}$ we compute $\Gamma_{s+m+l+1}^{\left[\substack{s\\ s+m}\right]\times k}$ for all $k \geq s+m+l+1$ and (2.61) follows.

The case $k = s+m+l+1$ is obtained in a similar way.

3. STATEMENT OF RESULTS

Theorem 3.1. *Let q be a rational integer ≥ 1, then*

(3.1)

$$g_{k,s,m}(t,\eta) = g(t,\eta) = \sum_{degY\leq k-1}\ \sum_{degZ\leq s-1} E(tYZ) \sum_{degU\leq s+m-1} E(\eta YU) = 2^{2s+m+k-r(D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta))},$$

(3.2)

$$\int_{\mathbb{P}\times\mathbb{P}} g^{q}_{k,s,m}(t,\eta)dtd\eta = 2^{(2s+m+k)(q-1)}\cdot 2^{-k+2}\cdot \sum_{i=0}^{\inf(2s+m,k)} \Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}\cdot 2^{-qi}.$$

Theorem 3.2. *Let $s \geq 2,\ m \geq 0,\ k \geq 1$ and $0 \leq i \leq \inf(2s+m,k)$. Then we have the following recurrent formula for the number $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}$ of rank i matrices of the form $\left[\frac{A}{B}\right]$, such that A is a $s\times k$ persymmetric matrix and B a $(s+m)\times k$ persymmetric matrix with entries in $\mathbb{F}_2$:*

(3.3)

$$\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k} = 2\cdot\Gamma_{i-1}^{\left[\substack{s-1\\ s-1+(m+1)}\right]\times k} + 4\cdot\Gamma_{i-1}^{\left[\substack{s\\ s+(m-1)}\right]\times k} - 8\cdot\Gamma_{i-2}^{\left[\substack{s-1\\ s-1+m}\right]\times k} + \Delta_{i}^{\left[\substack{s\\ s+m}\right]\times k}$$

where the remainder $\Delta_{i}^{\left[\substack{s\\ s+m}\right]\times k}$ is equal to

$$\sigma_{i,i,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k} - 3\cdot\sigma_{i-1,i-1,i-1}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k} + 2\cdot\sigma_{i-2,i-2,i-2}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}\cdot \tag{3.4}$$

Recall that

$$\sigma_{i,i,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}$$

is equal to the cardinality of the following set

$$\left\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1} \mid r(D^{\left[\substack{s-1\\ s-1+m}\right]\times k}(t,\eta)) = r(D^{\left[\substack{s\\ s+m-1}\right]\times k}(t,\eta)) = r(D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta)) = i\right\}.$$

Theorem 3.3. *Let $s \geq 2$ and $m \geq 0$, we have in the following two cases :*
The case $1 \leq k \leq 2s+m-2$

$$\sigma_{i,i,i}^{\left[\begin{array}{c} s \\ s+m \\ \hline \alpha_{s-} \\ \beta_{s+m-} \\ \\ \end{array}\right]\times k} = \begin{cases} 1 & \text{if } i=0, \quad k \geq 1, \\ 4\cdot\Gamma_{i}^{\left[\substack{s-1 \\ s-1+m}\right]\times i} - \Gamma_{i+1}^{\left[\substack{s-1 \\ s-1+m}\right]\times (i+1)} & \text{if } 1 \leq i \leq k-1, \\ 4\cdot\Gamma_{k}^{\left[\substack{s-1 \\ s-1+m}\right]\times k} & \text{if } i=k. \end{cases} \tag{3.5}$$

The case $k \geq 2s+m-2$

$$\sigma_{i,i,i}^{\left[\begin{array}{c} s \\ s+m \\ \hline \alpha_{s-} \\ \beta_{s+m-} \\ \\ \end{array}\right]\times k} = \begin{cases} 1 & \text{if } i=0, \\ 4\cdot\Gamma_{i}^{\left[\substack{s-1 \\ s-1+m}\right]\times i} - \Gamma_{i+1}^{\left[\substack{s-1 \\ s-1+m}\right]\times (i+1)} & \text{if } 1 \leq i \leq 2s+m-3, \\ 4\cdot\Gamma_{2s+m-2}^{\left[\substack{s-1 \\ s-1+m}\right]\times (2s+m-2)} & \text{if } i=2s+m-2. \end{cases} \tag{3.6}$$

Theorem 3.4. *The remainder $\Delta_{i}^{\left[\substack{s \\ s+m}\right]\times k}$ in the recurrent formula is equal to*

(3.7)
$$\begin{cases} 1 & \text{if } i=0,\ k \geq 1, \\ 4\cdot\Gamma_{1}^{\left[\substack{s-1 \\ s-1+m}\right]\times 1} - \Gamma_{2}^{\left[\substack{s-1 \\ s-1+m}\right]\times 2} & \text{if } i=1,\ k \geq 2, \\ 4\cdot\Gamma_{1}^{\left[\substack{s-1 \\ s-1+m}\right]\times 1} - 3 & \text{if } i=1,\ k=1, \\ 7\cdot\Gamma_{2}^{\left[\substack{s-1 \\ s-1+m}\right]\times 2} - 12\cdot\Gamma_{1}^{\left[\substack{s-1 \\ s-1+m}\right]\times 1} - \Gamma_{3}^{\left[\substack{s-1 \\ s-1+m}\right]\times 3} + 2 & \text{if } i=2,\ k \geq 3, \\ 7\cdot\Gamma_{2}^{\left[\substack{s-1 \\ s-1+m}\right]\times 2} - 12\cdot\Gamma_{1}^{\left[\substack{s-1 \\ s-1+m}\right]\times 1} + 2 & \text{if } i=2,\ k=2, \\ 7\cdot\Gamma_{i}^{\left[\substack{s-1 \\ s-1+m}\right]\times i} - 14\cdot\Gamma_{i-1}^{\left[\substack{s-1 \\ s-1+m}\right]\times (i-1)} + 8\cdot\Gamma_{i-2}^{\left[\substack{s-1 \\ s-1+m}\right]\times (i-2)} - \Gamma_{i+1}^{\left[\substack{s-1 \\ s-1+m}\right]\times (i+1)} & \text{if } 3 \leq i \leq 2s+m-3, k \geq i+1, \\ 7\cdot\Gamma_{i}^{\left[\substack{s-1 \\ s-1+m}\right]\times i} - 14\cdot\Gamma_{i-1}^{\left[\substack{s-1 \\ s-1+m}\right]\times (i-1)} + 8\cdot\Gamma_{i-2}^{\left[\substack{s-1 \\ s-1+m}\right]\times (i-2)} & \text{if } 3 \leq i \leq 2s+m-3,\ k=i, \\ 7\cdot\Gamma_{2s+m-2}^{\left[\substack{s-1 \\ s-1+m}\right]\times (2s+m-2)} - 14\cdot\Gamma_{2s+m-3}^{\left[\substack{s-1 \\ s-1+m}\right]\times (2s+m-3)} + 8\cdot\Gamma_{2s+m-4}^{\left[\substack{s-1 \\ s-1+m}\right]\times (2s+m-4)} & \text{if } i=2s+m-2,\ k \geq i, \\ -14\cdot\Gamma_{2s+m-2}^{\left[\substack{s-1 \\ s-1+m}\right]\times (2s+m-2)} + 8\cdot\Gamma_{2s+m-3}^{\left[\substack{s-1 \\ s-1+m}\right]\times (2s+m-3)} & \text{if } i=2s+m-1,\ k \geq i, \\ 8\cdot\Gamma_{2s+m-2}^{\left[\substack{s-1 \\ s-1+m}\right]\times (2s+m-2)} & \text{if } i=2s+m,\ k \geq i. \end{cases}$$

Theorem 3.5. *We have*

$$\Delta_{i}^{\left[\substack{s \\ s+m}\right]\times k} = \Delta_{i}^{\left[\substack{s \\ s+m}\right]\times (i+1)} \qquad \text{for } i \in [0, 2s+m-3],\ k \geq i+1, \tag{3.8}$$

$$\Delta_{i}^{\left[\substack{s \\ s+m}\right]\times k} = \Delta_{i}^{\left[\substack{s \\ s+m}\right]\times i} \qquad \text{for } i \in \{2s+m-2, 2s+m-1, 2s+m\},\ k \geq i. \tag{3.9}$$

Theorem 3.6. *We have for all $m \geq 0$*

$$\Gamma_{j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k+1)} - \Gamma_{j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = 0 \quad \textit{if} \quad 0 \leq j \leq s-1,\ k > j. \tag{3.10}$$

We have in the cases $m \in \{0,1\}$

$$\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times k} = \begin{cases} 3\cdot 2^{k+s-1} & \textit{if } j=0,\ k>s, \\ 21\cdot 2^{k+s+3j-4} & \textit{if } 1\leq j\leq s-1,\ k>s+j, \\ 3\cdot 2^{2k+2s-2} - 3\cdot 2^{k+4s-4} & \textit{if } j=s,\ k>2s, \end{cases} \tag{3.11}$$

(3.12)

$$\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix}\right]\times k} = \begin{cases} 2^{k+s-1} & \textit{if } j=0,\ k>s, \\ 11\cdot 2^{k+s-1} & \textit{if } j=1,\ k>s+1, \\ 21\cdot 2^{k+s+3j-5} & \textit{if } 2\leq j\leq s,\ k>s+j, \\ 3\cdot 2^{2k+2s-1} - 3\cdot 2^{k+4s-2} & \textit{if } j=s+1,\ k>2s+1. \end{cases}$$

In the case $m \geq 2$

$$\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = \begin{cases} 2^{k+s-1} & \textit{if } j=0,\ k>s, \\ 3\cdot 2^{k+s+2j-3} & \textit{if } 1\leq j\leq m-1,\ k>s+j, \\ 11\cdot 2^{k+s+2m-3} & \textit{if } j=m,\ k>s+m, \end{cases} \tag{3.13}$$

(3.14)

$$\Gamma_{s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+m+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = \begin{cases} 21\cdot 2^{k+s+2m+3j-4} & \textit{if } 1\leq j\leq s-1,\ k>s+m+j, \\ 3\cdot 2^{2k+2s+m-2} - 3\cdot 2^{k+4s+2m-4} & \textit{if } j=s,\ k>2s+m. \end{cases}$$

Theorem 3.7. *We have for* $m \geq 1$

$$\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = 8^{j-1}\cdot \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+(m-(j-1)) \end{smallmatrix}\right]\times(k-(j-1))} \qquad \textit{if } 1\leq j\leq m,\ k\geq s+j, \tag{3.15}$$

(3.16)

$$\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+(m-(j-1)) \end{smallmatrix}\right]\times(s+1)} = 2^{4s+(m-(j-1))} - 3\cdot 2^{3s-1} + 2^{2s-1} \qquad \textit{if } \ 1\leq j\leq m,\ k=s+j,$$

(3.17)

$$\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+(m-(j-1)) \end{smallmatrix}\right]\times(k-(j-1))} = 3\cdot 2^{k-j+s} + 21\cdot[2^{3s-1} - 2^{2s-1}] \qquad \textit{if } 1\leq j\leq m-1,\ k>s+j,$$

(3.18)

$$\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix}\right]\times(k-(m-1))} = 11\cdot 2^{k-m+s} + 21\cdot 2^{3s-1} - 11\cdot 2^{2s-1} \qquad \textit{if } j=m,\ k>s+m.$$

Theorem 3.8. *We have for* $m \geq 0$

(3.19)

$$\Gamma_{s+m+1+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = 8^{2j+m}\cdot \Gamma_{s-j+1}^{\left[\begin{smallmatrix} s-j \\ s-j \end{smallmatrix}\right]\times(k-m-2j)} \qquad \textit{if} \quad 0\leq j\leq s-1,\ k\geq s+m+1+j,$$

(3.20)
$$\Gamma_{s-j+1}^{\left[\substack{s-j\\ s-j}\right]\times(s-j+1)} = 2^{4s-4j} - 3\cdot 2^{3s-3j-1} + 2^{2s-2j-1} \qquad \text{if} \quad 0\leq j\leq s-1,\ k=s+m+1+j,$$

(3.21)
$$\Gamma_{s-j+1}^{\left[\substack{s-j\\ s-j}\right]\times(k-m-2j)} = 21\cdot[2^{k-m-3j+s-1} + 2^{3s-3j-1} - 5\cdot 2^{2s-2j-1}] \quad \text{if}\ \ 0\leq j\leq s-2,\ k>s+m+1+j,$$

(3.22)
$$\Gamma_{2}^{\left[\substack{1\\ 1}\right]\times(k-m-2s+2)} = 2^{2(k-m)-4s+4} - 3\cdot 2^{k-m-2s+2} + 2 \qquad \text{if} \quad j=s-1,\ k>2s+m.$$

Theorem 3.9. *We have*

(3.23)
$$\Gamma_{i}^{\left[\substack{s\\ s}\right]\times k} = \begin{cases} 1 & \text{if } i=0,\ k\geq 1,\\ 21\cdot 2^{3i-4} - 3\cdot 2^{2i-3} & \text{if } 1\leq i\leq s-1,\ k>i,\\ 3\cdot 2^{k+s-1} + 21\cdot 2^{3s-4} - 27\cdot 2^{2s-3} & \text{if } i=s,\ k>s,\\ 21\cdot[2^{k-2s+3i-4} + 2^{3i-4} - 5\cdot 2^{4i-2s-5}] & \text{if } s+1\leq i\leq 2s-1,\ k>i,\\ 2^{2k+2s-2} - 3\cdot 2^{k+4s-4} + 2^{6s-5} & \text{if } i=2s,\ k>2s. \end{cases}$$

Theorem 3.10. *We have*

$$\Gamma_{i}^{\left[\substack{s\\ s}\right]\times i} = \begin{cases} 2^{2s+2i-2} - 3\cdot 2^{3i-4} + 2^{2i-3} & \text{if } 1\leq i\leq s,\\ 2^{2s+2i-2} - 3\cdot 2^{3i-4} + 2^{4i-2s-5} & \text{if } s+1\leq i\leq 2s. \end{cases} \tag{3.24}$$

Theorem 3.11. *We have*

(3.25)
$$\Gamma_{i}^{\left[\substack{s\\ s+1}\right]\times k} = \begin{cases} 1 & \text{if } i=0,\ k\geq 1,\\ 21\cdot 2^{3i-4} - 3\cdot 2^{2i-3} & \text{if } 1\leq i\leq s-1,\ k>i,\\ 2^{k+s-1} + 21\cdot 2^{3s-4} - 11\cdot 2^{2s-3} & \text{if } i=s,\ k>s,\\ 11\cdot 2^{k+s-1} + 21\cdot 2^{3s-1} - 53\cdot 2^{2s-1} & \text{if } i=s+1,\ k>s+1,\\ 21\cdot[2^{k-2s+3i-5} + 2^{3i-4} - 5\cdot 2^{4i-2s-6}] & \text{if } s+2\leq i\leq 2s,\ k>i,\\ 2^{2k+2s-1} - 3\cdot 2^{k+4s-2} + 2^{6s-2} & \text{if } i=2s+1,\ k>2s+1. \end{cases}$$

Theorem 3.12. *We have*

$$\Gamma_{i}^{\left[\substack{s\\ s+1}\right]\times i} = \begin{cases} 2^{2s+2i-1} - 3\cdot 2^{3i-4} + 2^{2i-3} & \text{if } 1\leq i\leq s+1,\\ 2^{2s+2i-2} - 3\cdot 2^{3i-4} + 2^{4i-2s-6} & \text{if } s+2\leq i\leq 2s+1. \end{cases} \tag{3.26}$$

Theorem 3.13. *We have for $m\geq 2$*

(3.27)

$$\Gamma_i^{\left[\substack{s\\ s+m}\right]\times k} = \begin{cases} 1 & \text{if } i=0,\ k\geq 1,\\ 21\cdot 2^{3i-4}-3\cdot 2^{2i-3} & \text{if } 1\leq i\leq s-1,\ k>i,\\ 2^{k+s-1}+21\cdot 2^{3s-4}-11\cdot 2^{2s-3} & \text{if } i=s,\ k>s,\\ 3\cdot 2^{k-s+2i-3}+21\cdot\left[2^{3i-4}-2^{3i-s-4}\right] & \text{if } s+1\leq i\leq s+m-1,\ k>i,\\ 11\cdot 2^{k+s+2m-3}+21\cdot 2^{3s+3m-4}-53\cdot 2^{2s+3m-4} & \text{if } i=s+m,\ k>s+m,\\ 21\cdot\left[2^{k-2s+3i-m-4}+2^{3i-4}-5\cdot 2^{4i-2s-m-5}\right] & \text{if } s+m+1\leq i\leq 2s+m-1,\ k>i,\\ 2^{2k+2s+m-2}-3\cdot 2^{k+4s+2m-4}+2^{6s+3m-5} & \text{if } i=2s+m,\ k>2s+m. \end{cases}$$

Theorem 3.14. *We have for $m\geq 2$*

(3.28)

$$\Gamma_i^{\left[\substack{s\\ s+m}\right]\times i} = \begin{cases} 2^{2s+2i+m-2}-3\cdot 2^{3i-4}+2^{2i-3} & \text{if } 1\leq i\leq s+1,\\ 2^{2s+2i+m-2}-3\cdot 2^{3i-4}+2^{3i-s-4} & \text{if } s+2\leq i\leq s+m+1,\\ 2^{2s+2i+m-2}-3\cdot 2^{3i-4}+2^{4i-2s-m-5} & \text{if } s+m+2\leq i\leq 2s+m. \end{cases}$$

Theorem 3.15. *We denote by $R_q(k,s,m)$ the number of solutions $(Y_1,Z_1,U_1,\ldots,Y_q,Z_q,U_q)$ of the polynomial equations*

$$\begin{cases} Y_1Z_1+Y_2Z_2+\ldots+Y_qZ_q=0,\\ Y_1U_1+Y_2U_2+\ldots+Y_qU_q=0, \end{cases}$$

satisfying the degree conditions

$$degY_i\leq k-1,\quad degZ_i\leq s-1,\quad degU_i\leq s+m-1\quad for\quad 1\leq i\leq q.$$

Then

$$R_q(q,k,s,m)=\int_{\mathbb{P}\times\mathbb{P}} g_{k,s,m}^{q}(t,\eta)dtd\eta \tag{3.29}$$

$$= 2^{(2s+m+k)(q-1)}\cdot 2^{-k+2}\cdot\sum_{i=0}^{\inf(2s+m,k)}\Gamma_i^{\left[\substack{s\\ s+m}\right]\times k}\cdot 2^{-qi}.$$

Example. Let q = 3, k = 4, s = 3, m = 2. Then

$$\Gamma_i^{\left[\substack{3\\ 3+2}\right]\times 4} = \begin{cases} 1 & \text{if } i=0,\\ 9 & \text{if } i=1,\\ 78 & \text{if } i=2,\\ 648 & \text{if } i=3,\\ 15648 & \text{if } i=4. \end{cases}$$

Hence the number $R_3(4,3,2)$ of solutions $(Y_1,Z_1,U_1,Y_2,Z_2,U_2,Y_3,Z_3,U_3)$ of the polynomial equations

$$\begin{cases} Y_1Z_1+Y_2Z_2+Y_3Z_3=0,\\ Y_1U_1+Y_2U_2+Y_3U_3=0, \end{cases}$$

satisfying the degree conditions

$$degY_i\leq 3,\quad degZ_i\leq 2,\quad degU_i\leq 4\quad for\quad 1\leq i\leq 3$$

is equal to

$$\begin{aligned}\int_{\mathbb{P}\times\mathbb{P}} g^3_{4,3,2}(t,\eta)dtd\eta &= 2^{22}\cdot\sum_{i=0}^{4}\Gamma_i^{\left[\substack{3\\3+2}\right]\times 4}\cdot 2^{-3i}\\ &= 2^{22}\cdot[1+9\cdot 2^{-3}+78\cdot 2^{-6}+648\cdot 2^{-9}+15648\cdot 2^{-12}]\\ &= 35356672.\end{aligned}$$

Example. Let q = 4, k = 6, s = 5, m = 0. Then

$$\Gamma_i^{\left[\substack{5\\5}\right]\times 6} = \begin{cases} 1 & \text{if } i=0,\\ 9 & \text{if } i=1,\\ 78 & \text{if } i=2,\\ 648 & \text{if } i=3,\\ 5280 & \text{if } i=4,\\ 42624 & \text{if } i=5,\\ 999936 & \text{if } i=6.\end{cases}$$

Hence the number $R_4(6,5,0)$ of solutions $(Y_1,Z_1,U_1,Y_2,Z_2,U_2,Y_3,Z_3,U_3,Y_4,Z_4,U_4)$ of the polynomial equations

$$\begin{cases} Y_1Z_1+Y_2Z_2+Y_3Z_3+Y_4Z_4=0,\\ Y_1U_1+Y_2U_2+Y_3U_3+Y_4U_4=0,\end{cases}$$

satisfying the degree conditions

$$degY_i\leq 5,\quad degZ_i\leq 4,\quad degU_i\leq 4\quad for\quad 1\leq i\leq 4$$

is equal to

$$\begin{aligned}&\int_{\mathbb{P}\times\mathbb{P}} g^4_{4,5,0}(t,\eta)dtd\eta\\ &= 2^{44}\cdot\sum_{i=0}^{6}\Gamma_i^{\left[\substack{5\\5}\right]\times 6}\cdot 2^{-4i}\\ &= 2^{44}\cdot[1+9\cdot 2^{-4}+78\cdot 2^{-8}+648\cdot 2^{-12}+5280\cdot 2^{-16}+42624\cdot 2^{-20}+999936\cdot 2^{-24}]\\ &= 37014016\cdot 2^{20}.\end{aligned}$$

Example. The fraction of square double persymmetric $\left[\substack{s\\s+m}\right]\times(2s+m)$ matrices which are invertible is equal to $\frac{\Gamma_{2s+m}^{\left[\substack{s\\s+m}\right]\times(2s+m)}}{\sum_{i=0}^{2s+m}\Gamma_i^{\left[\substack{s\\s+m}\right]\times(2s+m)}}=\frac{3}{8}.$

4. EXPONENTIAL SUMS FORMULAS ON $\mathbb{K}^2$

In this section we compute exponential quadratic sums in $\mathbb{P}^2$ and show that they only depend on rank properties of some associated double persymmetric matrices. The following propositions are proved in [2].

Proposition 4.1. *The following holds :*

- *For every rational integer j, the measure of* $\mathbb{P}_j$ *is* 2^{-j}.
- *For every* $A\in\mathbb{F}_2[T]$, $E(A)=1$.
- *For* $u\in\mathbb{K}$ $\nu(u)\geq 2\Rightarrow$ $E(u)=1$.

Proposition 4.2. *Let j be a rational integer and $u \in \mathbb{K}$. Then*

$$\int_{\mathbb{P}_j} E(ut)dt = \begin{cases} 2^{-j} & if \quad \nu(u) > -j, \\ 0 & otherwise. \end{cases}$$

Proposition 4.3. *Let j be a rational integer and let $u \in \mathbb{K}$. Then*

$$\sum_{degB \leq j} E(Bu) = \begin{cases} 2^{j+1} & if \quad \nu(\{u\}) > j+1, \\ 0 & \qquad otherwise. \end{cases}$$

Lemma 4.4. *Let $Y \in \mathbb{F}_2[T]$ and a a rational integer ≥ 1. Then*

$$\sum_{degZ \leq a-1} E(tYZ) = \begin{cases} 2^{a} & if \quad \nu(\{tY\}) > a, \\ 0 & \qquad otherwise. \end{cases}$$

Proof. The Lemma follows from the Proposition 4.3 with u = tY and j = a -1. ☐

Lemma 4.5. *Let $t \in \mathbb{P}$ and $Y \in \mathbb{F}_2[T]$, $degY \leq b-1$. Then*

$$\nu(\{tY\}) > a \Longleftrightarrow Y \in \ker D_{a \times b}(t).$$

.

Proof. Let $Y = \sum_{j=0}^{b-1} \gamma_j T^j$, $\gamma_j \in \mathbb{F}_2$. Then

$$tY = (\sum_{i \geq 1} \alpha_i T^{-i})(\sum_{j=0}^{b-1} \gamma_j T^j) = \sum_{i \geq 1} \sum_{j=0}^{b-1} \alpha_i \gamma_j T^{-(i-j)}$$

and

$$\{tY\} = \left(\sum_{i=1}^{b} \alpha_i \gamma_{i-1} \right) T^{-1} + \left(\sum_{i=2}^{b+1} \alpha_i \gamma_{i-2} \right) T^{-2} + \ldots + \left(\sum_{i=s}^{b+a-1} \alpha_i \gamma_{i-a} \right) T^{-a} + \ldots$$

Therefore $\nu(\{tY\}) \geq a$ if and only if

$$\begin{bmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \ldots & \alpha_b \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 & \ldots & \alpha_{b+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ldots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ldots & \vdots \\ \alpha_a & \alpha_{a+1} & \alpha_{a+2} & \alpha_{a+3} & \alpha_{a+4} & \ldots & \alpha_{b+a-1} \end{bmatrix} \begin{bmatrix} \gamma_0 \\ \gamma_1 \\ \gamma_2 \\ \gamma_3 \\ \gamma_4 \\ \vdots \\ \gamma_{b-1} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

$$\Longleftrightarrow D_{a \times b}(t) \begin{pmatrix} \gamma_0 \\ \gamma_1 \\ \vdots \\ \gamma_{b-1} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \Longleftrightarrow Y = \begin{pmatrix} \gamma_0 \\ \gamma_1 \\ \vdots \\ \gamma_{b-1} \end{pmatrix} \in Ker\, D_{a \times b}(t).$$

☐

Lemma 4.6. *Let $(t, \eta) \in \mathbb{P} \times \mathbb{P}$. Then*

(4.1)
$$g(t, \eta) = \sum_{degY \leq k-1} \sum_{degZ \leq s-1} E(tYZ) \sum_{degU \leq s+m-1} E(\eta YU) = 2^{2s+m+k-r(D^{\left[\overset{s}{s+m}\right] \times k}(t,\eta))}.$$

Proof. By lemma 4.4 and lemma 4.5 we obtain with $a \rightarrow s \; and \; b \rightarrow k$

$$\sum_{degZ \leq s-1} E(tYZ) = 2^{s} \quad if \; and \; only \; if \quad Y \in \ker D_{s \times k}(t).$$

Similarly with $t \rightarrow \eta, \; a \rightarrow s+m \; and \; b \rightarrow k$ we get

$$\sum_{degU \leq s+m-1} E(\eta YU) = 2^{s+m} \quad if \; and \; only \; if \quad Y \in \ker D_{(s+m)\times k}(\eta).$$

We then deduce

$$\sum_{degZ \leq s-1} E(tYZ) \sum_{degU \leq s+m-1} E(\eta YU) = 2^{2s+m} \quad if \; and \; only \; if \quad Y \in \ker D^{\left[\substack{s \\ s+m}\right]\times k}(t,\eta)$$

and we obtain

$$\sum_{degY \leq k-1} \sum_{degZ \leq s-1} E(tYZ) \sum_{degU \leq s+m-1} E(\eta YU) = 2^{2s+m} \sum_{\substack{degY \leq k-1 \\ Y \in \ker D^{\left[\substack{s \\ s+m}\right]\times k}(t,\eta)}} 1 = 2^{2s+m} \cdot 2^{k - r(D^{\left[\substack{s \\ s+m}\right]\times k}(t,\eta))}.$$

□

Lemma 4.7. *Let* $(t,\eta) \in \mathbb{P} \times \mathbb{P}$ *and set*

$$g_1(t,\eta) = \sum_{degY \leq k-1} \sum_{degZ = s-1} E(tYZ) \sum_{degU \leq s+m-1} E(\eta YU).$$

Then

(4.2)

$$g_1(t,\eta) = \begin{cases} 2^{2s+m+k-1-r(D^{\left[\substack{s \\ s+m}\right]\times k}(t,\eta))} & if \; r(D^{\left[\substack{s-1 \\ s+m}\right]\times k}(t,\eta)) = r(D^{\left[\substack{s \\ s+m}\right]\times k}(t,\eta)), \\ 0 & otherwise \; . \end{cases}$$

Proof. We have

$$g_1(t,\eta) = \sum_{degY \leq k-1} \sum_{degZ \leq s-1} E(tYZ) \sum_{degU \leq s+m-1} E(\eta YU) - \sum_{degY \leq k-1} \sum_{degZ \leq s-2} E(tYZ) \sum_{degU \leq s+m-1} E(\eta YU).$$

By (4.1) we obtain

$$g_1(t,\eta) = 2^{2s+m+k-r(D^{\left[\substack{s \\ s+m}\right]\times k}(t,\eta))} - 2^{2s+m+k-1-r(D^{\left[\substack{s-1 \\ s+m}\right]\times k}(t,\eta))}.$$

Now either

$$r(D^{\left[\substack{s \\ s+m}\right]\times k}(t,\eta)) = r(D^{\left[\substack{s-1 \\ s+m}\right]\times k}(t,\eta))$$

or

$$r(D^{\left[\substack{s \\ s+m}\right]\times k}(t,\eta)) = r(D^{\left[\substack{s-1 \\ s+m}\right]\times k}(t,\eta)) + 1.$$

Lemma 4.7 follows. □

Lemma 4.8. *Let* $(t,\eta) \in \mathbb{P} \times \mathbb{P}$ *and set*

$$g_2(t,\eta) = \sum_{degY \leq k-1} \sum_{degZ \leq s-1} E(tYZ) \sum_{degU = s-1+m} E(\eta YU),$$

$$f_1(t,\eta) = \sum_{degY \leq k-1} \sum_{degZ = s-1} E(tYZ) \sum_{degU \leq s-2+m} E(\eta YU),$$

$$f_2(t,\eta) = \sum_{degY\leq k-1}\sum_{degZ\leq s-2} E(tYZ) \sum_{degU=s-2+(m+1)} E(\eta YU).$$

Then

(4.3)

$$g_2(t,\eta) = \begin{cases} 2^{2s+m+k-1-r(D^{\left[\overset{s}{s+m}\right]\times k}(t,\eta))} & \textit{if } r(D^{\left[\overset{s}{s+m}\right]\times k}(t,\eta)) = r(D^{\left[\overset{s}{s+(m-1)}\right]\times k}(t,\eta)),\\ 0 & \textit{otherwise,} \end{cases}$$

(4.4)

$$f_1(t,\eta) = \begin{cases} 2^{2s+m+k-2-r(D^{\left[\overset{s-1}{s-1+m}\right]\times k}(t,\eta))} & \textit{if } r(D^{\left[\overset{s-1}{s-1+m}\right]\times k}(t,\eta)) = r(D^{\left[\overset{s}{s+(m-1)}\right]\times k}(t,\eta)),\\ 0 & \textit{otherwise,} \end{cases}$$

(4.5)

$$f_2(t,\eta) = \begin{cases} 2^{2s+m+k-2-r(D^{\left[\overset{s-1}{s-1+m}\right]\times k}(t,\eta))} & \textit{if } r(D^{\left[\overset{s-1}{s-1+m}\right]\times k}(t,\eta)) = r(D^{\left[\overset{s-1}{s-1+(m+1)}\right]\times k}(t,\eta)),\\ 0 & \textit{otherwise.} \end{cases}$$

Proof. Similarly to the proof of Lemma 4.7 □

Lemma 4.9. *Set* $(t,\eta) = (\sum_{j\geq 1}\alpha_j T^{-j}, \sum_{j\geq 1}\beta_j T^{-j}) \in \mathbb{P}\times\mathbb{P}$ *and* $Y = \sum_{j=1}^{k-1}\delta_j T^j \in \mathbb{F}_2[T]$, $degY \leq k-1$.
Then

$$E(Y(tT^{s-1}+\eta T^{s+m-1})) = \begin{cases} 1 & \textit{if } \sum_{j=1}^{k-1}(\alpha_{s+j}+\beta_{s+m+j})\delta_j = 0,\\ -1 & \textit{if } \sum_{j=1}^{k-1}(\alpha_{s+j}+\beta_{s+m+j})\delta_j = 1. \end{cases} \tag{4.6}$$

Proof. We have

$$\begin{aligned} Y(tT^{s-1}+\eta T^{s+m-1}) &= \sum_{j=0}^{k-1}\delta_j T^j \cdot \left[\sum_{i\geq 1}\alpha_i T^{-i}\cdot T^{s-1} + \sum_{l\geq 1}\beta_l T^{-l}\cdot T^{s+m-1}\right]\\ &= \sum_{j=0}^{k-1}\left[\sum_{i\geq 1}\alpha_i\delta_j T^{s-1-i+j} + \sum_{l\geq 1}\beta_l\delta_j T^{s+m-1-l+j}\right]\\ &= \ldots + \left[\sum_{j=0}^{k-1}(\alpha_{s+j}+\beta_{s+m+j})\delta_j\right]\cdot T^{-1} + \ldots \end{aligned}$$

Now

$$\nu\left(\left\{Y(tT^{s-1}+\eta T^{s+m-1})\right\}\right) > 1 \Longleftrightarrow \sum_{j=0}^{k-1}(\alpha_{s+j}+\beta_{s+m+j})\delta_j = 0,$$

$$\nu\left(\left\{Y(tT^{s-1}+\eta T^{s+m-1})\right\}\right) = 1 \Longleftrightarrow \sum_{j=0}^{k-1}(\alpha_{s+j}+\beta_{s+m+j})\delta_j = 1,$$

which proves Lemma 4.9. □

Lemma 4.10. *Let* $(t,\eta)\in\mathbb{P}\times\mathbb{P}$ *and set*

$$h(t,\eta) = \sum_{degY\leq k-1}\sum_{degZ=s-1} E(tYZ) \sum_{degU=s+m-1} E(\eta YU).$$

Then
(4.7)

$$h(t,\eta)=\begin{cases} 2^{2s+m+k-2-r(D^{\left[\begin{smallmatrix} s-1\\ s-1+m\end{smallmatrix}\right]\times k}(t,\eta))} & \textit{if } r(D^{\left[\begin{smallmatrix} s-1\\ s-1+m\end{smallmatrix}\right]\times k}(t,\eta))=r(D^{\left[\begin{smallmatrix} s-1\\ s+m-1\\ \alpha_{s-}+\beta_{s+m-}\end{smallmatrix}\right]\times k}(t,\eta)),\\ 0 & \textit{otherwise .}\end{cases}$$

Proof. We have by the proof of lemma 4.6 and equation (4.6)

$$\begin{aligned}
h(t,\eta) &= \sum_{degY\leq k-1}\sum_{degZ=s-1}E(tYZ)\sum_{degU=s+m-1}E(\eta YU)\\
&= \sum_{degY\leq k-1}E(Y(tT^{s-1}+\eta T^{s+m-1}))\sum_{degZ\leq s-2}E(tYZ)\sum_{degU\leq s+m-2}E(\eta YU)\\
&= 2^{2s+m-2}\sum_{\substack{degY\leq k-1\\ Y\in\ker D^{\left[\begin{smallmatrix} s-1\\ s-1+m\end{smallmatrix}\right]\times k}}}E(Y(tT^{s-1}+\eta T^{s+m-1}))\\
&= 2^{2s+m-2}\left[\sum_{\substack{degY\leq k-1\\ Y\in\ker D^{\left[\begin{smallmatrix} s-1\\ s-1+m\end{smallmatrix}\right]\times k}(t,\eta)\\ \sum_{j=1}^{k-1}(\alpha_{s+j}+\beta_{s+m+j})\delta_j=0}}1-\sum_{\substack{degY\leq k-1\\ Y\in\ker D^{\left[\begin{smallmatrix} s-1\\ s-1+m\end{smallmatrix}\right]\times k}(t,\eta)\\ \sum_{j=1}^{k-1}(\alpha_{s+j}+\beta_{s+m+j})\delta_j=1}}1\right]\\
&= 2^{2s+m-2}\left[\sum_{\substack{degY\leq k-1\\ Y\in\ker D^{\left[\begin{smallmatrix} s-1\\ s+m-1\\ \alpha_{s-}+\beta_{s+m-}\end{smallmatrix}\right]\times k}(t,\eta)}}1-\left(\sum_{\substack{degY\leq k-1\\ Y\in\ker D^{\left[\begin{smallmatrix} s-1\\ s+m-1\end{smallmatrix}\right]\times k}(t,\eta)}}1-\sum_{\substack{degY\leq k-1\\ Y\in\ker D^{\left[\begin{smallmatrix} s-1\\ s+m-1\\ \alpha_{s-}+\beta_{s+m-}\end{smallmatrix}\right]\times k}(t,\eta)}}1\right)\right]\\
&= 2^{2s+m-2}\left[2\cdot\sum_{\substack{degY\leq k-1\\ Y\in\ker D^{\left[\begin{smallmatrix} s-1\\ s+m-1\\ \alpha_{s-}+\beta_{s+m-}\end{smallmatrix}\right]\times k}(t,\eta)}}1-\sum_{\substack{degY\leq k-1\\ Y\in\ker D^{\left[\begin{smallmatrix} s-1\\ s+m-1\end{smallmatrix}\right]\times k}(t,\eta)}}1\right]\\
&= 2^{2s+m-2}\left[2\cdot 2^{k-r(D^{\left[\begin{smallmatrix} s-1\\ s+m-1\\ \alpha_{s-}+\beta_{s+m-}\end{smallmatrix}\right]\times k}(t,\eta))}-2^{k-r(D^{\left[\begin{smallmatrix} s-1\\ s-1+m\end{smallmatrix}\right]\times k}(t,\eta))}\right]
\end{aligned}$$

which proves Lemma 4.10. □

Lemma 4.11. *Let* $(t,\eta)\in\mathbb{P}\times\mathbb{P}$ *and set*

$$\begin{aligned}
h(t,\eta) &= \sum_{degY\leq k-1}\sum_{degZ=s-1}E(tYZ)\sum_{degU=s+m-1}E(\eta YU),\\
v(t,\eta) &= \sum_{degY\leq k-1}\sum_{degZ\leq s-2}E(tYZ)\sum_{degU\leq s+m-2}E(\eta YU),\\
f_1(t,\eta) &= \sum_{degY\leq k-1}\sum_{degZ=s-1}E(tYZ)\sum_{degU\leq s+m-2}E(\eta YU),
\end{aligned}$$

$$f_2(t,\eta) = \sum_{degY \leq k-1} \sum_{degZ \leq s-2} E(tYZ) \sum_{degU = s+m-1} E(\eta YU).$$

Then we have

$$h^q(t,\eta) = h(t,\eta) \cdot v^{q-1}(t,\eta) \text{ for every rational integer } q \geq 2, \tag{4.8}$$

$$f_1^q(t,\eta) = f_1(t,\eta) \cdot v^{q-1}(t,\eta) \text{ for every rational integer } q \geq 2, \tag{4.9}$$

$$f_2^q(t,\eta) = f_2(t,\eta) \cdot v^{q-1}(t,\eta) \text{ for every rational integer } q \geq 2. \tag{4.10}$$

Proof. We get obviously

$$h^2(t,\eta) = \Big[2^{2s+m-2} \sum_{\substack{degY_1 \leq k-1 \\ Y_1 \in \ker D^{\left[\substack{s-1 \\ s-1+m}\right] \times k}}} E(Y_1(tT^{s-1}+\eta T^{s+m-1}))\Big] \cdot \Big[2^{2s+m-2} \sum_{\substack{degY_2 \leq k-1 \\ Y_2 \in \ker D^{\left[\substack{s-1 \\ s-1+m}\right] \times k}}} E(Y_2(tT^{s-1}+\eta T^{s+m-1}))\Big].$$

We set

$$\begin{cases} Y_1 + Y_2 = Y_3 & degY_3 \leq k-1, \\ \quad Y_2 = Y_4 & degY_4 \leq k-1. \end{cases}$$

Then we obtain

$$\begin{aligned} h^2(t,\eta) &= 2^{(2s+m-2)2} \sum_{\substack{degY_3 \leq k-1 \\ Y_3 \in \ker D^{\left[\substack{s-1 \\ s-1+m}\right] \times k}}} \sum_{\substack{degY_4 \leq k-1 \\ Y_4 \in \ker D^{\left[\substack{s-1 \\ s-1+m}\right] \times k}}} E((Y_4 + Y_3 + Y_4)(tT^{s-1} + \eta T^{s+m-1})) \\ &= \Big[2^{2s+m-2} \sum_{\substack{degY_3 \leq k-1 \\ Y_3 \in \ker D^{\left[\substack{s-1 \\ s-1+m}\right] \times k}}} E(Y_3(tT^{s-1} + \eta T^{s+m-1}))\Big] \cdot \Big[2^{2s+m-2} \sum_{\substack{degY_4 \leq k-1 \\ Y_4 \in \ker D^{\left[\substack{s-1 \\ s-1+m}\right] \times k}}} 1\Big] \\ &= h(t,\eta) \cdot v(t,\eta). \end{aligned}$$

By recurrence on q we get $h^q(t,\eta) = h(t,\eta) \cdot v^{q-1}(t,\eta)$.

The proof of (4.9) and (4.10) is similar to the proof of (4.10), that is

$$f_1^2(t,\eta) = \Big[2^{2s+m-2} \sum_{\substack{degY_1 \leq k-1 \\ Y_1 \in \ker D^{\left[\substack{s-1 \\ s-1+m}\right] \times k}}} E(Y_1 tT^{s-1})\Big] \cdot \Big[2^{2s+m-2} \sum_{\substack{degY_2 \leq k-1 \\ Y_2 \in \ker D^{\left[\substack{s-1 \\ s-1+m}\right] \times k}}} E(Y_2 tT^{s-1})\Big].$$

We set

$$\begin{cases} Y_1 + Y_2 = Y_3 & degY_3 \leq k-1, \\ \quad Y_2 = Y_4 & degY_4 \leq k-1. \end{cases}$$

Then we obtain

$$f_1^2(t,\eta) = 2^{(2s+m-2)2} \sum_{\substack{degY_3 \leq k-1 \\ Y_3 \in \ker D^{\left[\substack{s-1 \\ s-1+m}\right] \times k}}} \sum_{\substack{degY_4 \leq k-1 \\ Y_4 \in \ker D^{\left[\substack{s-1 \\ s-1+m}\right] \times k}}} E(Y_3 tT^{s-1})$$

$$= \Big[2^{2s+m-2}\sum_{\substack{degY_3\leq k-1\\ Y_3\in\ker D^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times k}}} E(Y_3tT^{s-1})\Big]\cdot\Big[2^{2s+m-2}\sum_{\substack{degY_4\leq k-1\\ Y_4\in\ker D^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times k}}} 1\Big]$$
$$= f_1(t,\eta)\cdot v(t,\eta).$$

By recurrence on q we get respectively

$$f_1^q(t,\eta) = f_1(t,\eta)\cdot v^{q-1}(t,\eta),$$
$$f_2^q(t,\eta) = f_2(t,\eta)\cdot v^{q-1}(t,\eta).$$

□

Lemma 4.12. *Let $(t,\eta)\in\mathbb{P}\times\mathbb{P}$ and $q\geq 2$, then we have for $1\leq i\leq q-1$*

$$h^i f_1^{q-i} = h^i f_2^{q-i} = \begin{cases} v^q & \textit{if } f_1f_2\neq 0,\\ 0 & \textit{otherwise,}\end{cases} \tag{4.11}$$

that is

$$h^i(t,\eta)f_1^{q-i}(t,\eta) = h^i(t,\eta)f_2^{q-i}(t,\eta) \tag{4.12}$$

is equal to

(4.13)
$$\begin{cases} 2^{q\cdot(2s+m+k-2-r(D^{\left[\begin{smallmatrix}s-1\\ s+m-1\end{smallmatrix}\right]\times k)}(t,\eta)))} & \textit{if } r(D^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times k}(t,\eta)) = r(D^{\left[\begin{smallmatrix}s\\ s+(m-1)\end{smallmatrix}\right]\times k}(t,\eta)) = r(D^{\left[\begin{smallmatrix}s-1\\ s+m\end{smallmatrix}\right]\times k}(t,\eta)),\\ 0 & \textit{otherwise.}\end{cases}$$

Proof. We have

$$f_1(t,\eta)h(t,\eta) = \Big[2^{2s+m-2}\sum_{\substack{degY_1\leq k-1\\ Y_1\in\ker D^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times k}}} E(Y_1(tT^{s-1}))\Big]\cdot\Big[2^{2s+m-2}\sum_{\substack{degY_2\leq k-1\\ Y_2\in\ker D^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times k}}} E(Y_2(tT^{s-1}+\eta T^{s+m-1}))\Big]$$
$$= 2^{(2s+m-2)2}\sum_{\substack{degY_1\leq k-1\\ Y_1\in\ker D^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times k}}}\sum_{\substack{degY_2\leq k-1\\ Y_2\in\ker D^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times k}}} E(t(Y_1+Y_2)T^{s-1})E(\eta Y_2T^{s+m-1}).$$

We set
$$\begin{cases} Y_1+Y_2 = Y_3 & degY_3\leq k-1,\\ Y_2 = Y_4 & degY_4\leq k-1.\end{cases}$$

Then we obtain

$$f_1(t,\eta)h(t,\eta) = 2^{(2s+m-2)2}\sum_{\substack{degY_3\leq k-1\\ Y_3\in\ker D^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times k}}}\sum_{\substack{degY_4\leq k-1\\ Y_4\in\ker D^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times k}}} E(tY_3T^{s-1})E(\eta Y_4T^{s+m-1})$$
$$= \Big[2^{2s+m-2}\sum_{\substack{degY_3\leq k-1\\ Y_3\in\ker D^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times k}}} E(tY_3T^{s-1})\Big]\cdot\Big[2^{2s+m-2}\sum_{\substack{degY_4\leq k-1\\ Y_4\in\ker D^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times k}}} E(\eta Y_4T^{s+m-1})\Big]$$
$$= f_1(t,\eta)\cdot f_2(t,\eta).$$

In the same way we get

$$f_2(t,\eta)\cdot h(t,\eta) = f_1(t,\eta)\cdot f_2(t,\eta).$$

Then

$$f_1(t,\eta)\cdot h(t,\eta) = f_2(t,\eta)\cdot h(t,\eta) = f_1(t,\eta)\cdot f_2(t,\eta). \tag{4.14}$$

By respectively (4.8), (4.9) and (4.10) we deduce

$$h = \begin{cases} v & \text{if } h \neq 0, \\ 0 & \text{otherwise,} \end{cases} \tag{4.15}$$

$$f_1 = \begin{cases} v & \text{if } f_1 \neq 0, \\ 0 & \text{otherwise,} \end{cases} \tag{4.16}$$

$$f_2 = \begin{cases} v & \text{if } f_2 \neq 0, \\ 0 & \text{otherwise.} \end{cases} \tag{4.17}$$

Now by (4.14), (4.15), (4.16) and (4.17) we have for $1 \leq i \leq q-1$

$$h^i f_1^{q-i} = h^i f_2^{q-i} = \begin{cases} v^i \cdot v^{q-i} = v^q & \text{if } h\cdot f_1 = h\cdot f_2 = f_1\cdot f_2 \neq 0, \\ 0 & \text{otherwise.} \end{cases} \tag{4.18}$$

Finally from (4.4), (4.5) and (4.18) we deduce (4.13). ☐

Lemma 4.13. *We denote by $R_q(k,s,m)$ the number of solutions $(Y_1, Z_1, U_1, \ldots, Y_q, Z_q, U_q)$ of the polynomial equations*

$$\begin{cases} Y_1Z_1 + Y_2Z_2 + \ldots + Y_qZ_q = 0, \\ Y_1U_1 + Y_2U_2 + \ldots + Y_qU_q = 0, \end{cases}$$

satisfying the degree conditions

$$deg Y_i \leq k-1, \quad deg Z_i \leq s-1, \quad deg U_i \leq s+m-1 \quad for \quad 1 \leq i \leq q.$$

Then

$$\begin{aligned} R_q(k,s,m) &= \int_{\mathbb{P}\times\mathbb{P}} g_{k,s,m}^q(t,\eta)\,dt\,d\eta \\ &= 2^{(2s+m+k)(q-1)}\cdot 2^{-k+2}\cdot \sum_{i=0}^{\inf(2s+m,k)} \Gamma_i^{\left[\substack{s \\ s+m}\right]\times k}\cdot 2^{-qi}. \end{aligned} \tag{4.19}$$

Proof. We have by (4.1) observing that $g(t,\eta)$ is constant on cosets of $\mathbb{P}_{k+s-1}\times\mathbb{P}_{k+s+m-1}$

$$\begin{aligned} \int_{\mathbb{P}^2} g^q(t,\eta)\,dt\,d\eta &= \sum_{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} 2^{q(2s+m+k-r(D^{\left[\substack{s \\ s+m}\right]\times k}(t,\eta))} \int_{\mathbb{P}_{k+s-1}} dt \int_{\mathbb{P}_{k+s+m-1}} dt\,d\eta \\ &= \sum_{i=0}^{\inf(2s+m,k)} \sum_{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} 2^{q(2s+m+k-i)} \int_{\mathbb{P}_{k+s-1}} dt \int_{\mathbb{P}_{k+s+m-1}} dt\,d\eta \\ &= \sum_{i=0}^{\inf(2s+m,k)} \Gamma_i^{\left[\substack{s \\ s+m}\right]\times k}\cdot 2^{-qi}\cdot 2^{q(2s+m+k)}\cdot 2^{-(k+s-1)}\cdot 2^{-(k+s+m-1)}. \end{aligned}$$

☐

5. A RECURRENT FORMULA FOR THE NUMBER $\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}$ OF RANK i MATRICES OF THE FORM $\left[\frac{A}{B}\right]$, WHERE A IS A $s\times k$ PERSYMMETRIC MATRIX AND B a $(s+m)\times k$ PERSYMMETRIC MATRIX WITH ENTRIES IN $\mathbb{F}_2$

In this section we establish a recurrent formula for the number of rank i matrices of the form $\left[\frac{A}{B}\right]$, where A and B are persymmetric.

Lemma 5.1. *We have*

$$\begin{aligned}(5.1)\quad \int_{\mathbb{P}^2} g_1^q(t,\eta)dtd\eta &= \int_{\{(t,\eta)\in\mathbb{P}^2 \mid h(t,\eta)\neq 0\}} v^q(t,\eta)dtd\eta + \int_{\{(t,\eta)\in\mathbb{P}^2 \mid f_1(t,\eta)\neq 0\}} v^q(t,\eta)dtd\eta \\ &\quad + (2^q-2)\cdot\int_{\{(t,\eta)\in\mathbb{P}^2 \mid f_1(t,\eta)\cdot f_2(t,\eta)\neq 0\}} v^q(t,\eta)dtd\eta.\end{aligned}$$

Proof. By the binomial theorem we obtain

$$(5.2)\qquad g_1^q = (h+f_1)^q = h^q + f_1^q + \sum_{i=1}^{q-1}\binom{q}{i} h^i\cdot f_1^{q-i}.$$

By integrating (5.2) on the the unit interval of $\mathbb{K}^2$ and using (4.15),(4.16) and (4.9) we get

(5.3)

$$\begin{aligned}\int_{\mathbb{P}^2} g_1^q(t,\eta)dtd\eta &= \int_{\mathbb{P}^2}(h+f_1)^q(t,\eta)dtd\eta \\ &= \int_{\mathbb{P}^2} h^q(t,\eta)dtd\eta + \int_{\mathbb{P}^2} f_1^q(t,\eta)dtd\eta + \int_{\mathbb{P}^2}\sum_{i=1}^{q-1}\binom{q}{i}h^i(t,\eta)\cdot f_1^{q-i}(t,\eta)dtd\eta \\ &= \int_{\{(t,\eta)\in\mathbb{P}^2 \mid h(t,\eta)\neq 0\}} v^q(t,\eta)dtd\eta + \int_{\{(t,\eta)\in\mathbb{P}^2 \mid f_1(t,\eta)\neq 0\}} v^q(t,\eta)dtd\eta \\ &\quad + (2^q-2)\cdot\int_{\{(t,\eta)\in\mathbb{P}^2 \mid f_1(t,\eta)\cdot f_2(t,\eta)\neq 0\}} v^q(t,\eta)dtd\eta.\end{aligned}$$

□

Lemma 5.2. *We have*

$$(5.4)\qquad \int_{\mathbb{P}^2} g_1^q(t,\eta)dtd\eta = 2^{q(2s+m+k-1)}\cdot 2^{-(2k+2s+m-2)}\cdot\sum_{i=0}^{\inf(2s+m-1,k)}\sigma_{i,i}^{\left[\begin{array}{c} s-1 \\ s+m \\ \hline \alpha_{s-}\end{array}\right]\times k}\cdot 2^{-qi}.$$

Proof. We have by (4.2) observing that $g_1(t,\eta)$ is constant on cosets of $\mathbb{P}_{k+s-1}\times\mathbb{P}_{k+s+m-1}$

$$\begin{aligned}\int_{\mathbb{P}^2} g_1^q(t,\eta)dtd\eta &= \sum_{\substack{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1} \\ r(D^{\left[\begin{smallmatrix} s-1 \\ s+m \end{smallmatrix}\right]\times k}(t,\eta)) = r(D^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}(t,\eta))}} 2^{q(2s+m+k-1-r(D^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}(t,\eta)))}\int_{\mathbb{P}_{k+s-1}}dt\int_{\mathbb{P}_{k+s+m-1}}d\eta \\ &= \sum_{i=0}^{\inf(2s+m-1,k)}\sum_{\substack{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1} \\ r(D^{\left[\begin{smallmatrix} s-1 \\ s+m \end{smallmatrix}\right]\times k}(t,\eta)) = r(D^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}(t,\eta)) = i}} 2^{q(2s+m+k-1-i)}\int_{\mathbb{P}_{k+s-1}}dt\int_{\mathbb{P}_{k+s+m-1}}d\eta\end{aligned}$$

$$= \sum_{i=0}^{\inf(2s+m-1,k)} \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m}{\alpha_{s-}} \end{smallmatrix}\right]\times k} \cdot 2^{-qi} \cdot 2^{q(2s+m+k-1)} \cdot 2^{-(k+s-1)} \cdot 2^{-(k+s+m-1)}.$$

□

Lemma 5.3. *We have*

$$\int_{\mathbb{P}^2} f_1^q(t,\eta)dtd\eta = 2^{q(2s+m+k-2)} \cdot 2^{-(2k+2s+m-3)} \cdot \sum_{i=0}^{\inf(2s+m-2,k)} \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \end{smallmatrix}\right]\times k} \cdot 2^{-qi}. \tag{5.5}$$

Proof. Similar to the proof of Lemma 5.2 using (4.4) and observing that $f_1(t,\eta)$ is constant on cosets of $\mathbb{P}_{k+s-1} \times \mathbb{P}_{k+s+m-2}$.

□

Lemma 5.4. *We have*

$$\int_{\{(t,\eta)\in\mathbb{P}^2 \mid f_1(t,\eta)\cdot f_2(t,\eta)\neq 0\}} v^q(t,\eta)dtd\eta = 2^{q(2s+m+k-2)} \cdot 2^{-(2k+2s+m-2)} \cdot \sum_{i=0}^{\inf(2s+m-2,k)} \sigma_{i,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k} \cdot 2^{-qi}. \tag{5.6}$$

Proof. We have by (4.12) and (4.13)

$$\int_{\{(t,\eta)\in\mathbb{P}^2 \mid f_1(t,\eta)\cdot f_2(t,\eta)\neq 0\}} v^q(t,\eta)dtd\eta$$

$$= \sum_{\substack{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1} \\ r(D^{\left[\begin{smallmatrix} s-1 \\ s-1+m \end{smallmatrix}\right]\times k}(t,\eta)) = r(D^{\left[\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix}\right]\times k}(t,\eta)) = r(D^{\left[\begin{smallmatrix} s-1 \\ s-1+m+1 \end{smallmatrix}\right]\times k}(t,\eta))}} 2^{q(2s+m+k-2-r(D^{\left[\begin{smallmatrix} s-1 \\ s-1+m \end{smallmatrix}\right]\times k}(t,\eta)))} \int_{\mathbb{P}_{k+s-1}} dt \int_{\mathbb{P}_{k+s+m-1}} d\eta$$

$$= \sum_{i=0}^{\inf(2s+m-2,k)} \sum_{\substack{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1} \\ r(D^{\left[\begin{smallmatrix} s-1 \\ s-1+m \end{smallmatrix}\right]\times k}(t,\eta)) = r(D^{\left[\begin{smallmatrix} s \\ s+m-1 \end{smallmatrix}\right]\times k}(t,\eta)) = r(D^{\left[\begin{smallmatrix} s-1 \\ s-1+m+1 \end{smallmatrix}\right]\times k}(t,\eta)) = i}} 2^{q(2s+m+k-2-i)} \int_{\mathbb{P}_{k+s-1}} dt \int_{\mathbb{P}_{k+s+m-1}} d\eta$$

$$= \sum_{i=0}^{\inf(2s+m-2,k)} \sigma_{i,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k} \cdot 2^{-qi} \cdot 2^{q(2s+m+k-2)} \cdot 2^{-(k+s-1)} \cdot 2^{-(k+s+m-1)}.$$

□

Lemma 5.5. *We have*

$$\sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}+\beta_{s+m-}} \end{smallmatrix}\right]\times k} = 2 \cdot \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \end{smallmatrix}\right]\times k}.$$

Proof. We consider the following $(2s+m-1)\times k$ matrix denoted by $D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \\ \alpha_{s-}+\beta_{s+m-} \end{smallmatrix}\right]\times k}(t,\eta)$ (see Section 1)

$$\left(\begin{array}{l|cccccc}
r_1 & \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\
r_2 & \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
r_{s-1} & \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
r_s & \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\
r_{s+1} & \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
r_{s+m} & \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
r_{2s+m-2} & \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
r_{2s+m-1} & \alpha_s+\beta_{s+m} & \alpha_{s+1}+\beta_{s+m+1} & \alpha_{s+2}+\beta_{s+m+2} & \dots & \alpha_{s+k-2}+\beta_{s+k+m-2} & \alpha_{s+k-1}+\beta_{s+k+m-1}
\end{array}\right).$$

↑

We recall that the rank of a matrix does not change under elementary row operations.
On the above matrix, we add the s+m+j-th row to the j-th row for $0 \leq j \leq s-2$ obtaining

$$\left(\begin{array}{l|cccccc}
r_1+r_{s+m} & \alpha_1+\beta_{m+1} & \alpha_2+\beta_{m+2} & \alpha_3+\beta_{m+3} & \dots & \alpha_{k-1}+\beta_{m+k-1} & \alpha_k+\beta_{m+k} \\
r_2+r_{s+m+1} & \alpha_2+\beta_{m+2} & \alpha_3+\beta_{m+3} & \alpha_4+\beta_{m+4} & \dots & \alpha_k+\beta_{m+k} & \alpha_{k+1}+\beta_{m+k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
r_{s-1}+r_{2s+m-2} & \alpha_{s-1}+\beta_{s-1+m} & \alpha_s+\beta_{s+m} & \alpha_{s+1}+\beta_{s+m+1} & \dots & \alpha_{s+k-3}+\beta_{s+k+m-3} & \alpha_{s+k-2}+\beta_{s+k+m-2} \\
r_s & \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\
r_{s+1} & \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \\
r_{s+m} & \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
r_{2s+m-2} & \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
r_{2s+m-1} & \alpha_s+\beta_{s+m} & \alpha_{s+1}+\beta_{s+m+1} & \alpha_{s+2}+\beta_{s+m+2} & \dots & \alpha_{s+k-2}+\beta_{s+k+m-2} & \alpha_{s+k-1}+\beta_{s+k+m-1}
\end{array}\right).$$

↑

In the above matrix we set $\begin{cases} \alpha_i+\beta_{i+m} & =\mu_i \quad \text{for} \quad 1 \leq i \leq k+s-2, \\ \beta_j & =\tau_j \quad \text{for} \quad 1 \leq j \leq k+s+m-2. \end{cases}$

(Remark that the map $\kappa : \mathbb{F}_2^{2k+2s+m-4} \longmapsto \mathbb{F}_2^{2k+2s+m-4}$ defined by
$(\alpha_1, \alpha_2, \dots, \alpha_{k+s-2}, \beta_1, \beta_2, \dots, \beta_{k+s+m-2}) \longmapsto (\mu_1, \mu_2, \dots, \mu_{k+s-2}, \tau_1, \tau_2, \dots, \tau_{k+s+m-2})$

is an isomorphisme). We then obtain the below matrix

↓

$$\begin{array}{c@{\quad}c}
\begin{array}{l}
r_1\\ r_2\\ \vdots\\ r_{s-1}\\ r_s\\ r_{s+1}\\ \vdots\\ r_{s+m}\\ \vdots\\ r_{2s+m-2}\\ \hline r_{2s+m-1}
\end{array}
&
\left(\begin{array}{cccccc}
\mu_1 & \mu_2 & \mu_3 & \cdots & \mu_{k-1} & \mu_k\\
\mu_2 & \mu_3 & \mu_4 & \cdots & \mu_k & \mu_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\mu_{s-1} & \mu_s & \mu_{s+1} & \cdots & \mu_{s+k-3} & \mu_{s+k-2}\\
\tau_1 & \tau_2 & \tau_3 & \cdots & \tau_{k-1} & \tau_k\\
\tau_2 & \tau_3 & \tau_4 & \cdots & \tau_k & \tau_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\tau_{m+1} & \tau_{m+2} & \tau_{m+3} & \cdots & \tau_{k+m-1} & \tau_{k+m}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\tau_{s+m-1} & \tau_{s+m} & \tau_{s+m+1} & \cdots & \tau_{s+m+k-3} & \tau_{s+m+k-2}\\
\hline
\mu_s & \mu_{s+1} & \mu_{s+2} & \cdots & \mu_{s+k-2} & \alpha_{s+k-1}+\beta_{s+k+m-1}
\end{array}\right)
\end{array}.$$

Comparing the above matrix with the below matrix $\updownarrow$

$$\begin{array}{c@{\quad}c}
\begin{array}{l}
r_1\\ r_2\\ \vdots\\ r_{s-1}\\ r_s\\ r_{s+1}\\ \vdots\\ r_{s+m}\\ \vdots\\ r_{2s+m-2}\\ \hline r_{2s+m-1}
\end{array}
&
\left(\begin{array}{cccccc}
\mu_1 & \mu_2 & \mu_3 & \cdots & \mu_{k-1} & \mu_k\\
\mu_2 & \mu_3 & \mu_4 & \cdots & \mu_k & \mu_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\mu_{s-1} & \mu_s & \mu_{s+1} & \cdots & \mu_{s+k-3} & \mu_{s+k-2}\\
\tau_1 & \tau_2 & \tau_3 & \cdots & \tau_{k-1} & \tau_k\\
\tau_2 & \tau_3 & \tau_4 & \cdots & \tau_k & \tau_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\tau_{m+1} & \tau_{m+2} & \tau_{m+3} & \cdots & \tau_{k+m-1} & \tau_{k+m}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\tau_{s+m-1} & \tau_{s+m} & \tau_{s+m+1} & \cdots & \tau_{s+m+k-3} & \tau_{s+m+k-2}\\
\hline
\mu_s & \mu_{s+1} & \mu_{s+2} & \cdots & \mu_{s+k-2} & \mu_{s+k-1}
\end{array}\right)
\end{array}$$

and observing that for all $\mu_{s+k-1} \in \mathbb{F}_2$

$$Card\left\{(\alpha_{s+k-1},\beta_{s+k+m-1}) \in \mathbb{F}_2^2 | \quad \alpha_{s+k-1}+\beta_{s+k+m-1} = \mu_{s+k-1}\right\} = 2$$

we obtain the following equality

$$Card\left\{(t,\eta) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \mid r(D^{\left[\overset{s-1}{s-1+m}\right]\times k}(t,\eta)) = r(D^{\left[\begin{smallmatrix} s-1\\ s+m-1\\ \alpha_{s-}+\beta_{s+m-}\end{smallmatrix}\right]\times k}(t,\eta)) = i\right\}$$
$$= 2 \cdot Card\left\{(t,\eta) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-2} \mid r(D^{\left[\overset{s-1}{s-1+m}\right]\times k}(t,\eta)) = r(D^{\left[\overset{s}{s-1+m}\right]\times k}(t,\eta)) = i\right\}.$$

Alternatively consider the following equivalences

$$r\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\alpha_s+\beta_{s+m} & \alpha_{s+1}+\beta_{s+m+1} & \alpha_{s+2}+\beta_{s+m+2} & \ldots & \alpha_{s+k-2}+\beta_{s+m+k-2} & \alpha_{s+k-1}+\beta_{s+m+k-1}
\end{array}\right)$$

$$= r\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2}
\end{pmatrix}$$

$$\Longleftrightarrow$$

$$r\left(\begin{array}{cccccc}
\alpha_1+\beta_{m+1} & \alpha_2+\beta_{m+2} & \alpha_3+\beta_{m+3} & \ldots & \alpha_{k-1}+\beta_{m+k-1} & \alpha_k+\beta_{m+k} \\
\alpha_2+\beta_{m+2} & \alpha_3+\beta_{m+3} & \alpha_4+\beta_{m+4} & \ldots & \alpha_k+\beta_{m+k} & \alpha_{k+1}+\beta_{m+k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1}+\beta_{s-1+m} & \alpha_s+\beta_{s+m} & \alpha_{s+1}+\beta_{s+m+1} & \ldots & \alpha_{s+k-3}+\beta_{s+k+m-3} & \alpha_{s+k-2}+\beta_{s+k+m-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\alpha_s+\beta_{s+m} & \alpha_{s+1}+\beta_{s+m+1} & \alpha_{s+2}+\beta_{s+m+2} & \ldots & \alpha_{s+k-2}+\beta_{s+m+k-2} & \alpha_{s+k-1}+\beta_{s+m+k-1}
\end{array}\right)$$

$$= r\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2}
\end{pmatrix}$$

$$\Longleftrightarrow$$

$$r\left(\begin{array}{cccccc} \mu_1 & \mu_2 & \mu_3 & \cdots & \mu_{k-1} & \mu_k \\ \mu_2 & \mu_3 & \mu_4 & \cdots & \mu_k & \mu_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \mu_{s-1} & \mu_s & \mu_{s+1} & \cdots & \mu_{s+k-3} & \mu_{s+k-2} \\ \tau_1 & \tau_2 & \tau_3 & \cdots & \tau_{k-1} & \tau_k \\ \tau_2 & \tau_3 & \tau_4 & \cdots & \tau_k & \tau_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \tau_{m+1} & \tau_{m+2} & \tau_{m+3} & \cdots & \tau_{k+m-1} & \tau_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \tau_{s+m-1} & \tau_{s+m} & \tau_{s+m+1} & \cdots & \tau_{s+m+k-3} & \tau_{s+m+k-2} \\ \hline \mu_s & \mu_{s+1} & \mu_{s+2} & \cdots & \mu_{s+k-2} & \alpha_{s+k-1}+\beta_{s+k+m-1} \end{array}\right) = r\left(\begin{array}{cccccc} \mu_1 & \mu_2 & \mu_3 & \cdots & \mu_{k-1} & \mu_k \\ \mu_2 & \mu_3 & \mu_4 & \cdots & \mu_k & \mu_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \mu_{s-1} & \mu_s & \mu_{s+1} & \cdots & \mu_{s+k-3} & \mu_{s+k-2} \\ \tau_1 & \tau_2 & \tau_3 & \cdots & \tau_{k-1} & \tau_k \\ \tau_2 & \tau_3 & \tau_4 & \cdots & \tau_k & \tau_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \tau_{m+1} & \tau_{m+2} & \tau_{m+3} & \cdots & \tau_{k+m-1} & \tau_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \tau_{s+m-1} & \tau_{s+m} & \tau_{s+m+1} & \cdots & \tau_{s+m+k-3} & \tau_{s+m+k-2} \end{array}\right).$$

We then get

$$\sigma_{i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-}+\beta_{s+m-} \end{array}\right]\times k} = 2\cdot\sigma_{i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \end{array}\right]\times k}.$$

☐

Lemma 5.6. *We have*

(5.7)

$$\int_{\mathbb{P}^2} h^q(t,\eta)dtd\eta = 2^{q(2s+m+k-2)}\cdot 2^{-(2k+2s+m-2)}\cdot\sum_{i=0}^{\inf(2s+m-2,k)}\sigma_{i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-}+\beta_{s+m-} \end{array}\right]\times k}\cdot 2^{-qi}.$$

Proof. We have by (4.7) observing that $h(t,\eta)$ is constant on cosets of $\mathbb{P}_{k+s-1}\times\mathbb{P}_{k+s+m-1}$

$$\begin{aligned} \int_{\mathbb{P}^2} h^q(t,\eta)dtd\eta &= \sum_{\substack{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1} \\ r(D^{\left[\begin{smallmatrix} s-1 \\ s-1+m \end{smallmatrix}\right]\times k}(t,\eta))=r(D^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \alpha_{s-}+\beta_{s+m-} \end{array}\right]\times k}(t,\eta))}} 2^{q(2s+m+k-2-r(D^{\left[\begin{smallmatrix} s-1 \\ s-1+m \end{smallmatrix}\right]\times k}(t,\eta)))}\int_{\mathbb{P}_{k+s-1}} dt\int_{\mathbb{P}_{k+s+m-1}} d\eta \\ &= \sum_{i=0}^{\inf(2s+m-2,k)}\sum_{\substack{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1} \\ r(D^{\left[\begin{smallmatrix} s-1 \\ s-1+m \end{smallmatrix}\right]\times k}(t,\eta))=r(D^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \alpha_{s-}+\beta_{s+m-} \end{array}\right]\times k}(t,\eta))=i}} 2^{q(2s+m+k-2-i)}\int_{\mathbb{P}_{k+s-1}} dt\int_{\mathbb{P}_{k+s+m-1}} d\eta \\ &= \sum_{i=0}^{\inf(2s+m-2,k)}\sigma_{i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-}+\beta_{s+m-} \end{array}\right]\times k}\cdot 2^{-qi}\cdot 2^{q(2s+m+k-2)}\cdot 2^{-(k+s-1)}\cdot 2^{-(k+s+m-1)}. \end{aligned}$$

☐

Lemma 5.7. *We have*

$$\int_{\mathbb{P}^2} h^q(t,\eta)dtd\eta = \int_{\mathbb{P}^2} f_1^q(t,\eta)dtd\eta. \tag{5.8}$$

Proof. Immediately obtained by comparing (5.5) and (5.7) using Lemma 5.5. ☐

Lemma 5.8. *We have for* $1 \leq i \leq \inf(2s+m-1,k)$

$$\sigma_{i-1,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\beta_{s+m-}} \\ \alpha_{s-} \end{smallmatrix}\right]\times k} = 2\cdot\sigma_{i-1,i-1,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k} . \tag{5.9}$$

Proof. From (5.1), (5.4),(5.5),(5.7) and (5.8) we have

$$\begin{aligned}
\int_{\mathbb{P}^2} g_1^q &= \int_{\mathbb{P}^2} h^q + \int_{\mathbb{P}^2} f_1^q + (2^q-2)\cdot\int_{f_1 f_2\neq 0} v^q \\
&= 2\cdot\int_{\mathbb{P}^2} f_1^q + (2^q-2)\cdot\int_{f_1 f_2\neq 0} v^q \\
&\Leftrightarrow 2^{q(2s+m+k-1)}\cdot 2^{-(2k+2s+m-2)}\cdot \sum_{i=0}^{\inf(2s+m-1,k)} \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m}{\alpha_{s-}} \end{smallmatrix}\right]\times k}\cdot 2^{-qi} \\
&= 2\cdot 2^{q(2s+m+k-2)}\cdot 2^{-(2k+2s+m-3)}\cdot \sum_{i=0}^{\inf(2s+m-2,k)} \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \end{smallmatrix}\right]\times k}\cdot 2^{-qi} \\
&+ (2^q-2)\cdot 2^{q(2s+m+k-2)}\cdot 2^{-(2k+2s+m-2)}\cdot \sum_{i=0}^{\inf(2s+m-2,k)} \sigma_{i,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k}\cdot 2^{-qi} \\
&\Leftrightarrow \sum_{i=0}^{\inf(2s+m-1,k)} \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m}{\alpha_{s-}} \end{smallmatrix}\right]\times k}\cdot 2^{-qi} \\
&= 2^{2-q}\cdot \sum_{i=0}^{\inf(2s+m-2,k)} \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \end{smallmatrix}\right]\times k}\cdot 2^{-qi} + (2^q-2)\cdot 2^{-q} \sum_{i=0}^{\inf(2s+m-2,k)} \sigma_{i,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k}\cdot 2^{-qi} \\
&\Leftrightarrow \sum_{i=0}^{\inf(2s+m-1,k)} \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m}{\alpha_{s-}} \end{smallmatrix}\right]\times k}\cdot 2^{-qi} \\
&= \sum_{i=0}^{\inf(2s+m-2,k)} 4\cdot\sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \end{smallmatrix}\right]\times k}\cdot 2^{-q(i+1)} \\
&+ \sum_{i=0}^{\inf(2s+m-2,k)} \sigma_{i,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k}\cdot 2^{-qi} - \sum_{i=0}^{\inf(2s+m-2,k)} 2\cdot\sigma_{i,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k}\cdot 2^{-q(i+1)}
\end{aligned}$$

(5.10)

$$\Leftrightarrow \sum_{i=0}^{\inf(2s+m-1,k)} \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m}{\alpha_{s-}} \end{smallmatrix}\right]\times k} \cdot 2^{-qi}$$

$$= \sum_{i=1}^{\inf(2s+m-2,k)+1} 4\cdot \sigma_{i-1,i-1}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \end{smallmatrix}\right]\times k} \cdot 2^{-qi}$$

$$+ \sum_{i=0}^{\inf(2s+m-2,k)} \sigma_{i,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k} \cdot 2^{-qi} - \sum_{i=1}^{\inf(2s+m-2,k)+1} 2\cdot \sigma_{i-1,i-1,i-1}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k} \cdot 2^{-qi}.$$

The case $k \leq 2s+m-2$

Observing that $4\cdot\sigma_{k,k}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \end{smallmatrix}\right]\times k} = 2\cdot\sigma_{k,k,k}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k}$ we deduce from (5.10)

(5.11)

$$\sum_{i=1}^{k}\left[\sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m}{\alpha_{s-}} \end{smallmatrix}\right]\times k} - 4\cdot\sigma_{i-1,i-1}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \end{smallmatrix}\right]\times k} - \sigma_{i,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k} + 2\cdot\sigma_{i-1,i-1,i-1}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k}\right]\cdot 2^{-qi} = 0 \; \textit{for all } q \geq 2.$$

Now we have obviously

$$\sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m}{\alpha_{s-}} \end{smallmatrix}\right]\times k} = \sigma_{i-1,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\beta_{s+m-}} \\ \alpha_{s-} \end{smallmatrix}\right]\times k} + \sigma_{i,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\beta_{s+m-}} \\ \alpha_{s-} \end{smallmatrix}\right]\times k}, \tag{5.12}$$

$$2\cdot\sigma_{i-1,i-1}^{\left[\begin{smallmatrix} s-1 \\ \frac{s-1+m}{\alpha_{s-}} \end{smallmatrix}\right]\times k} = \sigma_{i-1,i-1,i-1}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k} + \sigma_{i-1,i-1,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k}. \tag{5.13}$$

By (5.11), (5.12) and (5.13) we obtain for all $q \geq 2$

(5.14)

$$\sum_{i=1}^{k}\left[\sigma_{i-1,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\beta_{s+m-}} \\ \alpha_{s-} \end{smallmatrix}\right]\times k} + \sigma_{i,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\beta_{s+m-}} \\ \alpha_{s-} \end{smallmatrix}\right]\times k} - 2\cdot\sigma_{i-1,i-1,i-1}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k} - 2\cdot\sigma_{i-1,i-1,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k} - \sigma_{i,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k} + 2\cdot\sigma_{i-1,i-1,i-1}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k}\right]\cdot 2^{-qi} = 0.$$

From (5.14) we deduce

$$\sum_{i=1}^{k}\left[\sigma_{i-1,i,i}^{\begin{bmatrix} s-1 \\ \frac{s+m-1}{\beta_{s+m-}} \\ \alpha_{s-} \end{bmatrix}\times k} - 2\cdot\sigma_{i-1,i-1,i}^{\begin{bmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{bmatrix}\times k}\right]\cdot 2^{-qi} = 0 \; for \; all \; q \geq 2. \tag{5.15}$$

The case $k \geq 2s+m-1$

(5.16)

$$\begin{aligned}\sum_{i=0}^{2s+m-1}\sigma_{i,i}^{\begin{bmatrix} s-1 \\ \frac{s+m}{\alpha_{s-}} \end{bmatrix}\times k}\cdot 2^{-qi} &= \sum_{i=1}^{2s+m-1} 4\cdot\sigma_{i-1,i-1}^{\begin{bmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \end{bmatrix}\times k}\cdot 2^{-qi} \\ &+ \sum_{i=0}^{2s+m-2}\sigma_{i,i,i}^{\begin{bmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{bmatrix}\times k}\cdot 2^{-qi} - \sum_{i=1}^{2s+m-1} 2\cdot\sigma_{i-1,i-1,i-1}^{\begin{bmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{bmatrix}\times k}\cdot 2^{-qi}.\end{aligned}$$

Observing that $\sigma_{2s+m-1,2s+m-1,2s+m-1}^{\begin{bmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{bmatrix}\times k} = 0$ we deduce from (5.16), (5.12) and (5.13) that for all $q \geq 2$

(5.17)

$$\sum_{i=1}^{2s+m-1}\left[\sigma_{i-1,i,i}^{\begin{bmatrix} s-1 \\ \frac{s+m-1}{\beta_{s+m-}} \\ \alpha_{s-} \end{bmatrix}\times k} + \sigma_{i,i}^{\begin{bmatrix} s-1 \\ \frac{s+m-1}{\beta_{s+m-}} \\ \alpha_{s-} \end{bmatrix}\times k} - 2\cdot\sigma_{i-1,i-1,i-1}^{\begin{bmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{bmatrix}\times k} - 2\cdot\sigma_{i-1,i-1,i}^{\begin{bmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{bmatrix}\times k} - \sigma_{i,i,i}^{\begin{bmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{bmatrix}\times k} + 2\cdot\sigma_{i-1,i-1,i-1}^{\begin{bmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{bmatrix}\times k}\right]\cdot 2^{-qi} = 0.$$

From (5.17) we obtain

$$\sum_{i=1}^{2s+m-1}\left[\sigma_{i-1,i,i}^{\begin{bmatrix} s-1 \\ \frac{s+m-1}{\beta_{s+m-}} \\ \alpha_{s-} \end{bmatrix}\times k} - 2\cdot\sigma_{i-1,i-1,i}^{\begin{bmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{bmatrix}\times k}\right]\cdot 2^{-qi} = 0 \; for \; all \; q \geq 2 \tag{5.18}$$

and Lemma 5.8 is proved. □

Lemma 5.9.

(5.19)

$$\int_{\mathbb{P}^2} g_2^q(t,\eta)dtd\eta = \int_{\{(t,\eta)\in\mathbb{P}^2 | h(t,\eta)\neq 0\}} v^q(t,\eta)dtd\eta + \int_{\{(t,\eta)\in\mathbb{P}^2 | f_2(t,\eta)\neq 0\}} v^q(t,\eta)dtd\eta$$
$$+ (2^q-2)\cdot \int_{\{(t,\eta)\in\mathbb{P}^2 | f_1(t,\eta)\cdot f_2(t,\eta)\neq 0\}} v^q(t,\eta)dtd\eta.$$

Proof. Similar to the proof of Lemma 5.1. □

Lemma 5.10. *We have for* $1 \leq i \leq \inf(2s+m-1,k)$

$$\sigma_{i-1,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\beta_{s+m-}} \\ \alpha_{s-} \end{smallmatrix}\right]\times k} + \sigma_{i-1,i-1,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\beta_{s+m-}} \\ \alpha_{s-} \end{smallmatrix}\right]\times k} = \sigma_{i-1,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k} + \sigma_{i-1,i-1,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k}. \tag{5.20}$$

Proof. From (5.1) and (5.19) we get for all $q \geq 2$

(5.21)

$$\int_{\mathbb{P}^2} g_1^q(t,\eta)dtd\eta - \int_{\mathbb{P}^2} g_2^q(t,\eta)dtd\eta = \int_{\{(t,\eta)\in\mathbb{P}^2 | f_1(t,\eta)\neq 0\}} v^q(t,\eta)dtd\eta - \int_{\{(t,\eta)\in\mathbb{P}^2 | f_2(t,\eta)\neq 0\}} v^q(t,\eta)dtd\eta.$$

By (5.21) we get for all $q \geq 2$ the following equivalences

$$2^{q(2s+m+k-1)}\cdot 2^{-(2k+2s+m-2)}\cdot \sum_{i=0}^{\inf(2s+m-1,k)} \left[\sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m}{\alpha_{s-}} \end{smallmatrix}\right]\times k} - \sigma_{i,i}^{\left[\begin{smallmatrix} s \\ \frac{s+m-1}{\beta_{s+m-}} \end{smallmatrix}\right]\times k}\right]\cdot 2^{-qi}$$

$$= 2^{q(2s+m+k-2)}\cdot 2^{-(2k+2s+m-3)}\cdot \sum_{i=0}^{\inf(2s+m-2,k)} \left[\sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \end{smallmatrix}\right]\times k} - \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\beta_{s+m-}} \end{smallmatrix}\right]\times k}\right]\cdot 2^{-qi}$$

$$\Leftrightarrow \sum_{i=0}^{\inf(2s+m-1,k)} \left[\sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m}{\alpha_{s-}} \end{smallmatrix}\right]\times k} - \sigma_{i,i}^{\left[\begin{smallmatrix} s \\ \frac{s+m-1}{\beta_{s+m-}} \end{smallmatrix}\right]\times k}\right]\cdot 2^{-qi} = \sum_{i=0}^{\inf(2s+m-2,k)} 2\cdot \left[\sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \end{smallmatrix}\right]\times k} - \sigma_{i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\beta_{s+m-}} \end{smallmatrix}\right]\times k}\right]\cdot 2^{-q(i+1)}$$

$$\Leftrightarrow \sum_{i=0}^{\inf(2s+m-1,k)} \left[\sigma_{i-1,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\beta_{s+m-}} \\ \alpha_{s-} \end{smallmatrix}\right]\times k} - \sigma_{i-1,i,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k}\right]\cdot 2^{-qi} = \sum_{i=1}^{\inf(2s+m-2,k)+1} \left[\sigma_{i-1,i-1,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\alpha_{s-}} \\ \beta_{s+m-} \end{smallmatrix}\right]\times k} - \sigma_{i-1,i-1,i}^{\left[\begin{smallmatrix} s-1 \\ \frac{s+m-1}{\beta_{s+m-}} \\ \alpha_{s-} \end{smallmatrix}\right]\times k}\right]\cdot 2^{-qi}$$

(5.22)

$$\Leftrightarrow \sum_{i=1}^{\inf(2s+m-1,k)} \left[\sigma_{i-1,i,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\beta_{s+m-}\\ \alpha_{s-}}}\end{bmatrix}\times k} - \sigma_{i-1,i,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\alpha_{s-}\\ \beta_{s+m-}}}\end{bmatrix}\times k} - \sigma_{i-1,i-1,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\alpha_{s-}\\ \beta_{s+m-}}}\end{bmatrix}\times k} + \sigma_{i-1,i-1,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\beta_{s+m-}\\ \alpha_{s-}}}\end{bmatrix}\times k} \right] \cdot 2^{-qi} = 0.$$

Now (5.22) holds for all $q \geq 2$ which deduce Lemma 5.10.

□

Lemma 5.11. *We have*

$$\begin{aligned}(5.23)\qquad \Gamma_i^{\left[\substack{s\\ s+m}\right]\times k} &= \sigma_{i,i,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\alpha_{s-}\\ \beta_{s+m-}}}\end{bmatrix}\times k} + \sigma_{i-1,i,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\alpha_{s-}\\ \beta_{s+m-}}}\end{bmatrix}\times k} + \sigma_{i-1,i-1,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\alpha_{s-}\\ \beta_{s+m-}}}\end{bmatrix}\times k} + \sigma_{i-2,i-1,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\alpha_{s-}\\ \beta_{s+m-}}}\end{bmatrix}\times k} \\ &= \sigma_{i,i,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\beta_{s+m-}\\ \alpha_{s-}}}\end{bmatrix}\times k} + \sigma_{i-1,i,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\beta_{s+m-}\\ \alpha_{s-}}}\end{bmatrix}\times k} + \sigma_{i-1,i-1,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\beta_{s+m-}\\ \alpha_{s-}}}\end{bmatrix}\times k} + \sigma_{i-2,i-1,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\beta_{s+m-}\\ \alpha_{s-}}}\end{bmatrix}\times k}.\end{aligned}$$

Proof. The proof is obvious.

□

Lemma 5.12. *We have*

$$(5.24)\qquad \sigma_{i-2,i-1,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\alpha_{s-}\\ \beta_{s+m-}}}\end{bmatrix}\times k} = \sigma_{i-2,i-1,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\beta_{s+m-}\\ \alpha_{s-}}}\end{bmatrix}\times k}.$$

Proof. (5.24) follows from (5.23) and (5.20).

□

Lemma 5.13. *The following holds :*

- $$\sigma_{i-1,i,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\beta_{s+m-}\\ \alpha_{s-}}}\end{bmatrix}\times k} = 2 \cdot \sigma_{i-1,i-1,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\alpha_{s-}\\ \beta_{s+m-}}}\end{bmatrix}\times k}$$
- $$\sigma_{i-1,i,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\beta_{s+m-}\\ \alpha_{s-}}}\end{bmatrix}\times k} + \sigma_{i-1,i-1,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\beta_{s+m-}\\ \alpha_{s-}}}\end{bmatrix}\times k} = \sigma_{i-1,i,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\alpha_{s-}\\ \beta_{s+m-}}}\end{bmatrix}\times k} + \sigma_{i-1,i-1,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\alpha_{s-}\\ \beta_{s+m-}}}\end{bmatrix}\times k}$$
- $$\sigma_{i-2,i-1,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\alpha_{s-}\\ \beta_{s+m-}}}\end{bmatrix}\times k} = \sigma_{i-2,i-1,i}^{\begin{bmatrix}\frac{\substack{s-1\\ s+m-1}}{\substack{\beta_{s+m-}\\ \alpha_{s-}}}\end{bmatrix}\times k}$$

- $$\sigma_{i,i,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k} = \sigma_{i,i,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \beta_{s+m-}\\ \alpha_{s-}\\ \\ \end{array}\right]\times k}$$

- $$\Gamma_i^{\left[\begin{smallmatrix} s\\ s+m \end{smallmatrix}\right]\times k} = \sigma_{i,i,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k} + \sigma_{i-1,i,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k} + \sigma_{i-1,i-1,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k} + \sigma_{i-2,i-1,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}$$

- $$\Gamma_i^{\left[\begin{smallmatrix} s\\ s+m \end{smallmatrix}\right]\times k} = \sigma_{i,i,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \beta_{s+m-}\\ \alpha_{s-}\\ \\ \end{array}\right]\times k} + \sigma_{i-1,i,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \beta_{s+m-}\\ \alpha_{s-}\\ \\ \end{array}\right]\times k} + \sigma_{i-1,i-1,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \beta_{s+m-}\\ \alpha_{s-}\\ \\ \end{array}\right]\times k} + \sigma_{i-2,i-1,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \beta_{s+m-}\\ \alpha_{s-}\\ \\ \end{array}\right]\times k} .$$

Proof. The Theorem follows from Lemmas 5.8, 5.10, 5.11 and 5.12.

□

Lemma 5.14. *Let $s \geq 2,\ m \geq 0,\ k \geq 1$ and $0 \leq i \leq \inf(2s+m,k)$. Then we have the following recurrent formula for the number of rank i matrices of the form $\left[\frac{A}{B}\right]$, such that A is a $s \times k$ persymmetric matrix and B a $(s+m) \times k$ persymmetric matrix with entries in $\mathbb{F}_2$*

(5.25)

$$\Gamma_i^{\left[\begin{smallmatrix} s\\ s+m \end{smallmatrix}\right]\times k} = 2\cdot\Gamma_{i-1}^{\left[\begin{smallmatrix} s-1\\ s-1+(m+1) \end{smallmatrix}\right]\times k} + 4\cdot\Gamma_{i-1}^{\left[\begin{smallmatrix} s\\ s+(m-1) \end{smallmatrix}\right]\times k} - 8\cdot\Gamma_{i-2}^{\left[\begin{smallmatrix} s-1\\ s-1+m \end{smallmatrix}\right]\times k} + \Delta_i^{\left[\begin{smallmatrix} s\\ s+m \end{smallmatrix}\right]\times k}$$

where the remainder $\Delta_i^{\left[\begin{smallmatrix} s\\ s+m \end{smallmatrix}\right]\times k}$ *is equal to*

$$\sigma_{i,i,i}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k} - 3\cdot\sigma_{i-1,i-1,i-1}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k} + 2\cdot\sigma_{i-2,i-2,i-2}^{\left[\begin{array}{c} s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k} . \tag{5.26}$$

Proof. Consider the following two matrices

$$\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \cdots & \alpha_{k-1} & \alpha_k\\
\alpha_2 & \alpha_3 & \alpha_4 & \cdots & \alpha_k & \alpha_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \cdots & \alpha_{s+k-3} & \alpha_{s+k-2}\\
\beta_1 & \beta_2 & \beta_3 & \cdots & \beta_{k-1} & \beta_k\\
\beta_2 & \beta_3 & \beta_4 & \cdots & \beta_k & \beta_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \cdots & \beta_{k+m-1} & \beta_{k+m}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \cdots & \beta_{s+m+k-3} & \beta_{s+m+k-2}\\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \cdots & \alpha_{s+k-2} & \alpha_{s+k-1}\\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \cdots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{array}\right) \text{ and } \left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \cdots & \alpha_{k-1} & \alpha_k\\
\alpha_2 & \alpha_3 & \alpha_4 & \cdots & \alpha_k & \alpha_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \cdots & \alpha_{s+k-3} & \alpha_{s+k-2}\\
\beta_1 & \beta_2 & \beta_3 & \cdots & \beta_{k-1} & \beta_k\\
\beta_2 & \beta_3 & \beta_4 & \cdots & \beta_k & \beta_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \cdots & \beta_{k+m-1} & \beta_{k+m}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \cdots & \beta_{s+m+k-3} & \beta_{s+m+k-2}\\
\hline
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \cdots & \beta_{s+m+k-2} & \beta_{s+m+k-1}\\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \cdots & \alpha_{s+k-2} & \alpha_{s+k-1}
\end{array}\right).$$

We have by Lemma 5.13

$$\begin{aligned}
&2\cdot\Gamma_{i-1}^{\left[\substack{s-1\\ s-1+(m+1)}\right]\times k}+4\cdot\Gamma_{i-1}^{\left[\substack{s\\ s+(m-1)}\right]\times k}-8\cdot\Gamma_{i-2}^{\left[\substack{s-1\\ s-1+m}\right]\times k}+\Delta_{i}^{\left[\substack{s\\ s+m}\right]\times k}\\
&=\sigma_{i-1,i-1,i-1}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \overset{\beta_{s+m-}}{\alpha_{s-}}\\ \\ \end{array}\right]\times k}+\sigma_{i-1,i-1,i}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \overset{\beta_{s+m-}}{\alpha_{s-}}\\ \\ \end{array}\right]\times k}+\sigma_{i-2,i-1,i-1}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \overset{\beta_{s+m-}}{\alpha_{s-}}\\ \\ \end{array}\right]\times k}+\sigma_{i-2,i-1,i}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \overset{\beta_{s+m-}}{\alpha_{s-}}\\ \\ \end{array}\right]\times k}\\
&+2\cdot\sigma_{i-1,i-1,i-1}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}+2\cdot\sigma_{i-2,i-1,i-1}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}+2\cdot\sigma_{i-1,i-1,i}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}+2\cdot\sigma_{i-2,i-1,i}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}\\
&-2\cdot\sigma_{i-2,i-2,i-2}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}-2\cdot\sigma_{i-2,i-2,i-1}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}-2\cdot\sigma_{i-2,i-1,i-1}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}-2\cdot\sigma_{i-2,i-1,i}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}\\
&+\sigma_{i,i,i}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}-3\cdot\sigma_{i-1,i-1,i-1}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}+2\cdot\sigma_{i-2,i-2,i-2}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}\\
&=\sigma_{i-1,i-1,i}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \overset{\beta_{s+m-}}{\alpha_{s-}}\\ \\ \end{array}\right]\times k}+\sigma_{i-2,i-1,i-1}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \overset{\beta_{s+m-}}{\alpha_{s-}}\\ \\ \end{array}\right]\times k}+\sigma_{i-2,i-1,i}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \overset{\beta_{s+m-}}{\alpha_{s-}}\\ \\ \end{array}\right]\times k}+2\cdot\sigma_{i-1,i-1,i}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}\\
&-2\cdot\sigma_{i-2,i-2,i-1}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}+\sigma_{i,i,i}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}\\
&=\sigma_{i,i,i}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \overset{\beta_{s+m-}}{\alpha_{s-}}\\ \\ \end{array}\right]\times k}+\sigma_{i-1,i,i}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \overset{\beta_{s+m-}}{\alpha_{s-}}\\ \\ \end{array}\right]\times k}+\sigma_{i-1,i-1,i}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \overset{\beta_{s+m-}}{\alpha_{s-}}\\ \\ \end{array}\right]\times k}+\sigma_{i-2,i-1,i}^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \overset{\beta_{s+m-}}{\alpha_{s-}}\\ \\ \end{array}\right]\times k}\\
&=\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}.
\end{aligned}$$

□

6. RANK PROPERTIES OF A PARTITION OF DOUBLE PERSYMMETRIC MATRICES

Consider the following partition of the matrix

$$D^{\left[\begin{array}{c}\frac{s-1}{s+m-1}\\ \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}$$

$$\left(\begin{array}{ccccc|c}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1} \\
\hline
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{array}\right).$$

By integrating some appropriate exponential sums on the unit interval of $\mathbb{K}^2$ with integral equal to zero,
we deduce the following rank formulas for all $j \in [0, \inf(2s+m-2, k-1)]$:

$$\#\left(\begin{array}{c|c} j & j \\ \hline j & j+1 \\ \hline j & j+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} = 2\cdot\#\left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} = 2\cdot\#\left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}.$$

Lemma 6.1. *Let $(t,\eta) \in \mathbb{P}\times\mathbb{P}$ and set*

$$\psi(t,\eta) = \sum_{degY=k-1}\sum_{degZ\leq s-1} E(tYZ) \sum_{degU\leq s-1+m} E(\eta YU).$$

Then
(6.1)
$$\psi(t,\eta) = \begin{cases} 2^{2s+m+k-1-r(D^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times k}(t,\eta))} & \text{if } r(D^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times k}(t,\eta)) = r(D^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times (k-1)}(t,\eta)), \\ 0 & \text{otherwise.} \end{cases}$$

Proof. We have

$$\begin{aligned}
\psi(t,\eta) &= \sum_{degY=k-1}\sum_{degZ\leq s-1} E(tYZ) \sum_{degU\leq s+m-1} E(\eta YU) \\
&= \sum_{degY\leq k-1}\sum_{degZ\leq s-1} E(tYZ) \sum_{degU\leq s+m-1} E(\eta YU) - \sum_{degY\leq k-2}\sum_{degZ\leq s-1} E(tYZ) \sum_{degU\leq s+m-1} E(\eta YU) \\
&= 2^{2s+m}\cdot \sum_{\substack{degY\leq k-1 \\ Y\in\ker D^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times k}(t,\eta)}} 1 - 2^{2s+m}\cdot \sum_{\substack{degY\leq k-1 \\ Y\in\ker D^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times (k-1)}(t,\eta)}} 1 \\
&= 2^{2s+m}\cdot 2^{k-r(D^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times k}(t,\eta))} - 2^{2s+m}\cdot 2^{k-1-r(D^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times (k-1)}(t,\eta))}.
\end{aligned}$$

□

Lemma 6.2. *We have*
(6.2) $\phi^2(t,\eta) = \phi_1(t,\eta)\cdot\phi_2(t,\eta)$, *where*

$$\begin{aligned}
\phi(t,\eta) &= \sum_{degY=k-1}\sum_{degZ\leq s-1} E(tYZ)\sum_{degU=s+m-1} E(\eta YU),\\
\phi_1(t,\eta) &= \sum_{degY=k-1}\sum_{degZ\leq s-1} E(tYZ)\sum_{degU\leq s+m-2} E(\eta YU) \quad \textit{and}\\
\phi_2(t,\eta) &= \sum_{degY\leq k-2}\sum_{degZ\leq s-1} E(tYZ)\sum_{degU=s+m-1} E(\eta YU).
\end{aligned}$$

Proof. We obtain

$$\begin{aligned}
&\phi^2(t,\eta)\\
&= \left[2^{2s+m-1}\cdot \sum_{\substack{degY_1=k-1\\ Y_1\in\ker D^{\left[\overset{s}{s+m-1}\right]\times k}(t,\eta)}} E(\eta Y_1T^{s+m-1})\right]\cdot\left[2^{2s+m-1}\cdot \sum_{\substack{degY_2=k-1\\ Y_2\in\ker D^{\left[\overset{s}{s+m-1}\right]\times k}(t,\eta)}} E(\eta Y_2T^{s+m-1})\right].
\end{aligned}$$

We set

$$\begin{cases} Y_1+Y_2=Y_3 & degY_3\leq k-2,\\ \quad Y_1=Y_4 & degY_4=k-1.\end{cases}$$

Then we get obviously

$$\begin{cases} Y_1\in\ker D^{\left[\overset{s}{s+m-1}\right]\times k}(t,\eta) & degY_1=k-1,\\ Y_2\in\ker D^{\left[\overset{s}{s+m-1}\right]\times k}(t,\eta) & degY_2=k-1,\end{cases}$$

$$\Leftrightarrow\begin{cases} Y_3\in\ker D^{\left[\overset{s}{s+m-1}\right]\times (k-1)}(t,\eta) & degY_3\leq k-2,\\ Y_4\in\ker D^{\left[\overset{s}{s+m-1}\right]\times k}(t,\eta) & degY_4=k-1.\end{cases}$$

And we obtain

$$\begin{aligned}
\phi^2(t,\eta) &= 2^{(2s+m-1)2}\sum_{\substack{degY_4=k-1\\ Y_4\in\ker D^{\left[\overset{s}{s+m-1}\right]\times k}(t,\eta)}}\ \sum_{\substack{degY_3\leq k-2\\ Y_3\in\ker D^{\left[\overset{s}{s+m-1}\right]\times (k-1)}(t,\eta)}} E(\eta Y_3T^{s+m-1})\\
&= \left[2^{2s+m-1}\sum_{\substack{degY_4=k-1\\ Y_4\in\ker D^{\left[\overset{s}{s+m-1}\right]\times k}(t,\eta)}} 1\right]\cdot\left[2^{2s+m-1}\sum_{\substack{degY_3\leq k-2\\ Y_3\in\ker D^{\left[\overset{s}{s+m-1}\right]\times (k-1)}(t,\eta)}} E(\eta Y_3T^{s+m-1})\right]\\
&= \phi_1(t,\eta)\cdot\phi_2(t,\eta).
\end{aligned}$$

□

Lemma 6.3. *We have the following equivalences*

$$\begin{aligned}
\phi(t,\eta)\neq 0 &\Leftrightarrow \phi^2(t,\eta)\neq 0\\
&\Leftrightarrow \phi_1(t,\eta)\cdot\phi_2(t,\eta)\neq 0\\
&\Leftrightarrow r(D^{\left[\overset{s}{s+m-1}\right]\times (k-1)}(t,\eta)) = r(D^{\left[\overset{s}{s+m-1}\right]\times k}(t,\eta)) = r(D^{\left[\overset{s}{s+m}\right]\times (k-1)}(t,\eta)).
\end{aligned}$$

Proof. By Lemma 6.1 with $m \rightarrow m-1$ we have
(6.3)

$$\phi_1(t,\eta) = \begin{cases} 2^{2s+m+k-2-r(D^{\left[\begin{smallmatrix} s \\ s+(m-1)\end{smallmatrix}\right]\times k}(t,\eta))} & \text{if } r(D^{\left[\begin{smallmatrix} s \\ s+(m-1)\end{smallmatrix}\right]\times k}(t,\eta)) = r(D^{\left[\begin{smallmatrix} s \\ s+(m-1)\end{smallmatrix}\right]\times (k-1)}(t,\eta)), \\ 0 & \text{otherwise} \end{cases}$$

and by (4.5) with $k \rightarrow k-1,\ s \rightarrow s+1$ and $m \rightarrow m-1$ we have
(6.4)

$$\phi_2(t,\eta) = \begin{cases} 2^{2s+m+k-2-r(D^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times (k-1)}(t,\eta))} & \text{if } r(D^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times (k-1)}(t,\eta)) = r(D^{\left[\begin{smallmatrix} s \\ s+(m-1)\end{smallmatrix}\right]\times (k-1)}(t,\eta)), \\ 0 & \text{otherwise.} \end{cases}$$

Lemma 6.3 follows then from (6.3) and (6.4).

□

Lemma 6.4. *Let* $(t,\eta) \in \mathbb{P}\times\mathbb{P}$, *then*

$$\phi(t,\eta) = \sum_{degY=k-1}\sum_{degZ\leq s-1} E(tYZ) \sum_{degU=s+m-1} E(\eta YU)$$

is given by

$$\begin{cases} 2^{2s+m+k-j-2} & \textit{if} \quad (t,\eta) \in \begin{array}{|c|c|c|} k-1 & k & \\ \hline \cdot & \cdot & D^{\left[\begin{smallmatrix} s-1 \\ s+m-1\end{smallmatrix}\right]\times\cdot} \\ \hline j & j & \alpha_s - \\ \hline j & j & \beta_{s+m} - \end{array} \\ -2^{2s+m+k-j-2} & \textit{if} \quad (t,\eta) \in \begin{array}{|c|c|c|} k-1 & k & \\ \hline \cdot & \cdot & D^{\left[\begin{smallmatrix} s-1 \\ s+m-1\end{smallmatrix}\right]\times\cdot} \\ \hline j & j & \alpha_s - \\ \hline j & j+1 & \beta_{s+m} - \end{array} \\ 0 & \textit{otherwise.} \end{cases}$$

Proof. We consider the following two cases in which by Lemma 6.3 $\phi(t,\eta)$ is different from zero.

First case :
$r(D^{\left[\begin{smallmatrix} s \\ s+m-1\end{smallmatrix}\right]\times (k-1)}(t,\eta)) = r(D^{\left[\begin{smallmatrix} s \\ s+m-1\end{smallmatrix}\right]\times k}(t,\eta)) = r(D^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times (k-1)}(t,\eta)) = r(D^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times (k-1)}(t,\eta)) =$
j.

Second case :
$r(D^{\left[\begin{smallmatrix} s \\ s+m-1\end{smallmatrix}\right]\times (k-1)}(t,\eta)) = r(D^{\left[\begin{smallmatrix} s \\ s+m-1\end{smallmatrix}\right]\times k}(t,\eta)) = r(D^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times (k-1)}(t,\eta)) = j$ and $r(D^{\left[\begin{smallmatrix} s \\ s+m\end{smallmatrix}\right]\times k}(t,\eta)) =$
$j+1$.
We obtain using (4.3) :

In the first case

$$\begin{aligned}
\phi(t,\eta) &= \sum_{degY=k-1}\sum_{degZ\leq s-1} E(tYZ)\sum_{degU=s+m-1} E(\eta YU)\\
&= \sum_{degY\leq k-1}\sum_{degZ\leq s-1} E(tYZ)\sum_{degU=s+m-1} E(\eta YU) - \sum_{degY\leq k-2}\sum_{degZ\leq s-1} E(tYZ)\sum_{degU=s+m-1} E(\eta YU)\\
&= 2^{2s+m+k-1-j} - 2^{2s+m+k-2-j} = 2^{2s+m+k-2-j}.
\end{aligned}$$

In the second case

$$\begin{aligned}
\phi(t,\eta) &= \sum_{degY=k-1}\sum_{degZ\leq s-1} E(tYZ)\sum_{degU=s+m-1} E(\eta YU)\\
&= \sum_{degY\leq k-1}\sum_{degZ\leq s-1} E(tYZ)\sum_{degU=s+m-1} E(\eta YU) - \sum_{degY\leq k-2}\sum_{degZ\leq s-1} E(tYZ)\sum_{degU=s+m-1} E(\eta YU)\\
&= 0 - 2^{2s+m+k-2-j} = -2^{2s+m+k-2-j}.
\end{aligned}$$

And otherwise $\phi(t,\eta)$ is equal to zero.

$\square$

Lemma 6.5. *We have*

$$\int_{\mathbb{P}\times\mathbb{P}} \phi^{2q+1}(t,\eta)dtd\eta = 0.$$

Proof. The integral $\int_{\mathbb{P}\times\mathbb{P}} \phi^{2q+1}(t,\eta)dtd\eta$ is equal to the number of solutions $(Y_1, Z_1, U_1, Y_2, Z_2, U_2, \ldots, Y_{2q+1}, Z_{2q+1}, U_{2q+1})$ of the polynomial equations

$$\begin{cases} Y_1Z_1 + Y_2Z_2 + \ldots + Y_{2q+1}Z_{2q+1} = 0,\\ Y_1U_1 + Y_2U_2 + \ldots + Y_{2q+1}U_{2q+1} = 0,\end{cases}$$

satisfying the degree conditions
$\deg Y_i = k-1, \quad \deg Z_i \leq s-1 \quad \deg U_i = s+m-1 \quad for \quad 1 \leq i \leq 2q+1.$
Now 2q+1 is odd so $\deg\sum_{i=1}^{2q+1} Y_iU_i$ is equal to k+s+m-2. The Lemma follows.

$\square$

Lemma 6.6. *Let $j \in \mathbb{N}$ such that $0 \leq j \leq \inf(2s+m-1, k-1)$, then*

$$\#\left(\begin{array}{c|c} j & j \\ \hline j & j \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} = \#\left(\begin{array}{c|c} j & j \\ \hline j & j+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}.$$

Proof. We define

$$\begin{aligned}
\mathbb{A} = \{(t,\eta) \in \mathbb{P}^2 \mid r(D^{\left[\overset{s}{s+m-1}\right]\times(k-1)}(t,\eta)) = r(D^{\left[\overset{s}{s+m-1}\right]\times k}(t,\eta))\\
= r(D^{\left[\overset{s}{s+m}\right]\times(k-1)}(t,\eta)) = r(D^{\left[\overset{s}{s+m}\right]\times k}(t,\eta))\}
\end{aligned}$$

and

$$\begin{aligned}
\mathbb{B} = \{(t,\eta) \in \mathbb{P}^2 \mid r(D^{\left[\overset{s}{s+m-1}\right]\times(k-1)}(t,\eta)) = r(D^{\left[\overset{s}{s+m-1}\right]\times k}(t,\eta)) = r(D^{\left[\overset{s}{s+m}\right]\times(k-1)}(t,\eta)),\\
r(D^{\left[\overset{s}{s+m}\right]\times k}(t,\eta)) = r(D^{\left[\overset{s}{s+m-1}\right]\times(k-1)}(t,\eta)) + 1\}.
\end{aligned}$$

By lemma 6.4 we have by observing that $\phi(t,\eta)$ is constant on cosets of $\mathbb{P}_{k+s-1}\times\mathbb{P}_{k+s+m-1}$

$$\int_{\mathbb{P}\times\mathbb{P}}\phi^{2q+1}(t,\eta)dt\eta=\int_{\mathbb{A}}\phi^{2q+1}(t,\eta)dt\eta+\int_{\mathbb{B}}\phi^{2q+1}(t,\eta)dt\eta$$

$$=\sum_{(t,\eta)\in\mathbb{A}\bigcap\left(\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\right)}2^{(2s+m+k-2-r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-1)}(t,\eta))(2q+1)}\int_{\mathbb{P}_{s+k-1}}dt\int_{\mathbb{P}_{s+k+m-1}}d\eta$$

$$+\sum_{(t,\eta)\in\mathbb{B}\bigcap\left(\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\right)}-2^{(2s+m+k-2-r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-1)}(t,\eta))(2q+1)}\int_{\mathbb{P}_{s+k-1}}dt\int_{\mathbb{P}_{s+k+m-1}}d\eta$$

$$=$$

$$\sum_{j=0}^{\inf(2s+m-1,k-1)}2^{(2s+m+k-2-j)(2q+1)}\,.\,\#\left(\begin{array}{c|c} j & j\\ \hline j & j\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}\int_{\mathbb{P}_{s+k-1}}dt\int_{\mathbb{P}_{s+k+m-1}}d\eta$$
$$-\sum_{j=0}^{\inf(2s+m-1,k-1)}2^{(2s+m+k-2-j)(2q+1)}\,.\,\#\left(\begin{array}{c|c} j & j\\ \hline j & j+1\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}\int_{\mathbb{P}_{s+k-1}}dt\int_{\mathbb{P}_{s+k+m-1}}d\eta.$$

Now by Lemma 6.5 we get for all $q\in\mathbb{N}$

$$\sum_{j=0}^{\inf(2s+m-1,k-1)}2^{(-j)(2q+1)}\,.\,\left(\#\left(\begin{array}{c|c} j & j\\ \hline j & j\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}-\#\left(\begin{array}{c|c} j & j\\ \hline j & j+1\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}\right)=0,$$

which proves Lemma 6.6.

□

Lemma 6.7. *Let* $(t,\eta)\in\mathbb{P}^2$, *q be a rational integer* ≥ 2 *and*

$$\theta_1(t,\eta)=\sum_{degY=k-1}\sum_{degZ=s-1}E(tYZ)\sum_{degU\leq s+m-2}E(\eta YU),$$
$$\theta_2(t,\eta)=\sum_{degY\leq k-2}\sum_{degZ\leq s-1}E(tYZ)\sum_{degU=s+m-1}E(\eta YU)$$

Then $\theta_1(t,\eta)$ *is given by*

$$
(6.5)\qquad
\begin{cases}
2^{2s+m+k-j-3} & \text{if} \quad (t,\eta) \in
\begin{array}{|c|c|c|}
k-1 & k & \\ \hline
j & j & D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix}\right]\times\cdot} \\ \hline
j & j & \alpha_s- \\ \hline
\cdot & \cdot & \beta_{s+m}-
\end{array} \\[2em]
-2^{2s+m+k-j-3} & \text{if} \quad (t,\eta) \in
\begin{array}{|c|c|c|}
k-1 & k & \\ \hline
j & j & D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix}\right]\times\cdot} \\ \hline
j & j+1 & \alpha_s- \\ \hline
\cdot & \cdot & \beta_{s+m}-
\end{array} \\
0 & \text{otherwise},
\end{cases}
$$

and $\theta_2(t,\eta)$ *is given by*

$$
(6.6)\qquad
\begin{cases}
2^{2s+m+k-2-j} & \text{if} \quad (t,\eta) \in
\begin{array}{|c|c|c|}
k-1 & k & \\ \hline
\cdot & \cdot & D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix}\right]\times\cdot} \\ \hline
j & \cdot & \alpha_s- \\ \hline
j & \cdot & \beta_{s+m}-
\end{array} \\[2em]
0 & \text{otherwise},
\end{cases}
$$

Proof. The proofs of (6.5), (6.6) are respectively similar to the proofs of Lemmas 6.4, 4.7. ☐

Lemma 6.8. *Let* $(t,\eta) \in \mathbb{P}^2$ *and* q *be a rational integer* ≥ 2, *then* $\theta_1(t,\eta)\cdot\theta_2^{q-1}(t,\eta)$ *is equal to*

$$
(6.7)\qquad
\begin{cases}
2^{q-1}\cdot 2^{(2s+m+k-3-j)q} & \text{if} \quad (t,\eta) \in
\begin{array}{|c|c|c|}
k-1 & k & \\ \hline
j & j & D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix}\right]\times\cdot} \\ \hline
j & j & \alpha_s- \\ \hline
j & \cdot & \beta_{s+m}-
\end{array} \\[2em]
-2^{q-1}\cdot 2^{(2s+m+k-3-j)q} & \text{if} \quad (t,\eta) \in
\begin{array}{|c|c|c|}
k-1 & k & \\ \hline
j & j & D^{\left[\begin{smallmatrix} s-1 \\ s+m-1 \end{smallmatrix}\right]\times\cdot} \\ \hline
j & j+1 & \alpha_s- \\ \hline
j & j+1 & \beta_{s+m}-
\end{array} \\
0 & \text{otherwise}.
\end{cases}
$$

Proof. We consider the following partition of the matrix

$$
D^{\left[\begin{array}{c} s-1 \\ \hline s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \\ \\ \\ \end{array}\right]\times k}
$$

$$\left(\begin{array}{ccccc|c}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1} \\
\hline
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{array}\right).$$

Obviously by (6.6) we have

(6.8)

$$\theta_2^{q-1}(t,\eta)=\begin{cases}2^{(2s+m+k-2-j)(q-1)} & \text{if } r(D^{\left[\substack{s\\ s-1+m}\right]\times(k-1)}(t,\eta))=r(D^{\left[\substack{s\\ s+m}\right]\times(k-1)}(t,\eta))=j,\\ 0 & \text{otherwise .}\end{cases}$$

Then by considering the above matrix, using (6.5), (6.8) and elementary rank properties we get

$\theta_1(t,\eta)\cdot\theta_2^{q-1}(t,\eta)$ is equal to $2^{q-1}\cdot 2^{(2s+m+k-3-j)q}$ if and only if

$$r(D^{\left[\substack{s-1\\ s-1+m}\right]\times(k-1)}(t,\eta))=r(D^{\left[\substack{s-1\\ s+m-1}\right]\times k}(t,\eta))=r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-1)}(t,\eta))$$
$$=r(D^{\left[\substack{s\\ s+m-1}\right]\times k}(t,\eta))=r(D^{\left[\substack{s\\ s+m}\right]\times(k-1)}(t,\eta))=j$$

and

$$r(D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta))=j \text{ or } j+1$$

$\theta_1(t,\eta)\cdot\theta_2^{q-1}(t,\eta)$ is equal to $-2^{q-1}\cdot 2^{(2s+m+k-3-j)q}$ if and only if

$$r(D^{\left[\substack{s-1\\ s-1+m}\right]\times(k-1)}(t,\eta))=r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-1)}(t,\eta))=r(D^{\left[\substack{s\\ s+m}\right]\times(k-1)}(t,\eta))=r(D^{\left[\substack{s-1\\ s+m-1}\right]\times k}(t,\eta))=j$$

and

$$r(D^{\left[\substack{s-1\\ s+m-1}\right]\times k}(t,\eta))=r(D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta))=j+1$$

And in all the other cases $\theta_1(t,\eta)\cdot\theta_2^{q-1}(t,\eta)$ is equal to zero, which proves Lemma 6.8. ☐

Lemma 6.9. *We have*

$$\int_{\mathbb{P}\times\mathbb{P}}\theta_1(t,\eta)\cdot\theta_2^{q-1}(t,\eta)dtd\eta=0.$$

Proof. The integral $\int_{\mathbb{P}\times\mathbb{P}}\theta_1(t,\eta)\cdot\theta_2^{q-1}(t,\eta)dtd\eta$ is equal to the number of solutions $(Y,Z,U,Y_1,Z_1,U_1,\ldots,Y_{q-1},Z_{q-1},U_{q-1})$ of the polynomial equations

$$\begin{cases} YZ+Y_1Z_1+\ldots+Y_{q-1}Z_{q-1}=0,\\ YU+Y_1U_1+\ldots+Y_{q-1}U_{q-1}=0,\end{cases}$$

satisfying the degree conditions
$\deg Y=k-1,\quad \deg Z=s-1\quad \deg U\leq s+m-2$
$\deg Y_i\leq k-2,\quad \deg Z_i\leq s-1\quad \deg U_i=s+m-1\quad for 1\leq i\leq q-1.$
By degree considerations we have that $\deg\left[YZ+Y_1Z_1+\ldots+Y_{q-1}Z_{q-1}\right]$ is equal to s + k - 2.
Lemma 6.9 follows. ☐

Lemma 6.10. *We have for all $j\in\mathbb{Z}$ such that $0\leq j\leq\inf(2s+m-2,k-1)$,*

$$\#\left(\begin{array}{c|c} j & j\\ \hline j & j\\ \hline j & j\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}+\#\left(\begin{array}{c|c} j & j\\ \hline j & j\\ \hline j & j+1\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}-\#\left(\begin{array}{c|c} j & j\\ \hline j & j+1\\ \hline j & j+1\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}=0.$$

Recall that

$$\#\left(\begin{array}{c|c} j_1 & j_2\\ \hline j_3 & j_4\\ \hline j_5 & j_6\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}$$

denotes the cardinality of the following set

$$\Big\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\mid r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t,\eta))=j_1,\quad r(D^{\left[\substack{s-1\\ s+m-1}\right]\times k}(t,\eta))=j_2,$$
$$r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-1)}(t,\eta))=j_3,\quad r(D^{\left[\substack{s\\ s+m-1}\right]\times k}(t,\eta))=j_4,$$
$$r(D^{\left[\substack{s\\ s+m}\right]\times(k-1)}(t,\eta))=j_5,\quad r(D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta))=j_6\Big\}$$

for $(j_1,j_2,j_3,j_4,j_5,j_6)\in\mathbb{N}^6$.

Proof. We define

$$\mathbb{A}=\{(t,\eta)\in\mathbb{P}^2\mid r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t,\eta))=r(D^{\left[\substack{s-1\\ s+m-1}\right]\times k}(t,\eta))$$
$$=r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-1)}(t,\eta))=r(D^{\left[\substack{s\\ s+m-1}\right]\times k}(t,\eta))=r(D^{\left[\substack{s\\ s+m}\right]\times(k-1)}(t,\eta))\}$$

and

$$\mathbb{B} = \{(t,\eta)\in\mathbb{P}^2 \mid r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t,\eta)) = r(D^{\left[\substack{s-1\\ s+m-1}\right]\times k}(t,\eta)) = r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-1)}(t,\eta))$$
$$= r(D^{\left[\substack{s\\ s+m}\right]\times(k-1)}(t,\eta)), \quad r(D^{\left[\substack{s\\ s+m-1}\right]\times k}(t,\eta)) = r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t,\eta))+1\}.$$

By (6.7) we have, observing that $\theta_1(t,\eta)\cdot\theta_2^{q-1}(t,\eta)$ is constant on cosets of $\mathbb{P}_{k+s-1}\times\mathbb{P}_{k+s+m-1}$

$$\int_{\mathbb{P}\times\mathbb{P}}\theta_1(t,\eta)\cdot\theta_2^{q-1}(t,\eta)dt\eta = \int_{\mathbb{A}}\theta_1(t,\eta)\cdot\theta_2^{q-1}(t,\eta)dt\eta + \int_{\mathbb{B}}\theta_1(t,\eta)\cdot\theta_2^{q-1}(t,\eta)dt\eta$$

$$= \sum_{(t,\eta)\in\mathbb{A}\bigcap\left(\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\right)} 2^{q-1}\cdot 2^{(2s+m+k-3-r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t,\eta))q}\int_{\mathbb{P}_{s+k-1}}dt\int_{\mathbb{P}_{s+k+m-1}}d\eta$$

$$+ \sum_{(t,\eta)\in\mathbb{B}\bigcap\left(\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\right)} -2^{q-1}\cdot 2^{(2s+m+k-3-r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t,\eta))q}\int_{\mathbb{P}_{s+k-1}}dt\int_{\mathbb{P}_{s+k+m-1}}d\eta$$

$$=$$

$$\sum_{j=0}^{\inf(2s+m-2,k-1)} 2^{q-1}\cdot 2^{(2s+m+k-3-j)q}\,.\,\#\left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}\cdot 2^{-(2k+2s+m-2)}$$

$$+\sum_{j=0}^{\inf(2s+m-2,k-1)} 2^{q-1}\cdot 2^{(2s+m+k-3-j)q}\,.\,\#\left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}\cdot 2^{-(2k+2s+m-2)}$$

$$-\sum_{j=0}^{\inf(2s+m-2,k-1)} 2^{q-1}\cdot 2^{(2s+m+k-3-j)q}\,.\,\#\left(\begin{array}{c|c} j & j \\ \hline j & j+1 \\ \hline j & j+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}\cdot 2^{-(2k+2s+m-2)}.$$

Now by Lemma 6.9 we get for all $q\geq 2$

$$\sum_{j=0}^{\inf(2s+m-2,k-1)} 2^{-jq}\,.\left[\#\left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} +\#\left(\begin{array}{c|c} j & j \\ \hline j & j \\ \hline j & j+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}\right.$$
$$\left. -\#\left(\begin{array}{c|c} j & j \\ \hline j & j+1 \\ \hline j & j+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}\right] = 0.$$

□

Lemma 6.11. *Let* $(t,\eta)\in\mathbb{P}\times\mathbb{P}$,*and* q *be a rational integer* ≥ 2.
Set

$$\phi_3(t,\eta) = \sum_{degY\leq k-2}\sum_{degZ=s-1}E(tYZ)\sum_{degU\leq s+m-2}E(\eta YU).$$

Then we have

(6.9)

$$\phi_3^{q-1}(t,\eta)=\begin{cases} 2^{(2s+m+k-3-j)\cdot(q-1)} & \textit{if} \quad (t,\eta)\in \begin{array}{|c|c|c|} k-1 & k & \\ \hline j & \cdot & D^{\left[\begin{smallmatrix} s-1 \\ s+m-1\end{smallmatrix}\right]\times\cdot} \\ \hline j & \cdot & \alpha_s- \\ \hline \cdot & \cdot & \beta_{s+m}- \end{array} \\ 0 & \textit{otherwise} . \end{cases}$$

Proof. Lemma 6.11 follows from (4.4) with $k \to k-1$. ☐

Lemma 6.12. *Let $(t,\eta)\in\mathbb{P}^2$, and q be a rational integer ≥ 2, then $\phi(t,\eta)\cdot\theta_3^{q-1}(t,\eta)$ is equal to*

$$\begin{cases} 2\cdot 2^{(2s+m+k-3-j)q} & \textit{if} \quad (t,\eta)\in \begin{array}{|c|c|c|} k-1 & k & \\ \hline j & j & D^{\left[\begin{smallmatrix} s-1 \\ s+m-1\end{smallmatrix}\right]\times\cdot} \\ \hline j & j & \alpha_s- \\ \hline j & j & \beta_{s+m}- \end{array} \\ -2\cdot 2^{(2s+m+k-3-j)q} & \textit{if} \quad (t,\eta)\in \begin{array}{|c|c|c|} k-1 & k & \\ \hline j & j & D^{\left[\begin{smallmatrix} s-1 \\ s+m-1\end{smallmatrix}\right]\times\cdot} \\ \hline j & j & \alpha_s- \\ \hline j & j+1 & \beta_{s+m}- \end{array} \\ 0 & \textit{otherwise}. \end{cases} \tag{6.10}$$

Proof. Recall that by Lemma 6.4 $\phi(t,\eta)$ is equal to

$$\begin{cases} 2^{2s+m+k-j-2} & \text{if} \quad (t,\eta)\in \begin{array}{|c|c|c|} k-1 & k & \\ \hline \cdot & \cdot & D^{\left[\begin{smallmatrix} s-1 \\ s+m-1\end{smallmatrix}\right]\times\cdot} \\ \hline j & j & \alpha_s- \\ \hline j & j & \beta_{s+m}- \end{array} \\ -2^{2s+m+k-j-2} & \text{if} \quad (t,\eta)\in \begin{array}{|c|c|c|} k-1 & k & \\ \hline \cdot & \cdot & D^{\left[\begin{smallmatrix} s-1 \\ s+m-1\end{smallmatrix}\right]\times\cdot} \\ \hline j & j & \alpha_s- \\ \hline j & j+1 & \beta_{s+m}- \end{array} \\ 0 & \text{otherwise}. \end{cases}$$

Then by (6.9) we obtain by elementary rank properties (6.10) and Lemma 6.12 is proved.

☐

Lemma 6.13. *We have*

$$\int_{\mathbb{P}\times\mathbb{P}}\phi(t,\eta)\cdot\theta_3^{q-1}(t,\eta)dtd\eta=0.$$

Proof. The integral $\int_{\mathbb{P}\times\mathbb{P}}(t,\eta)\phi(t,\eta)\cdot\theta_3^{q-1}(t,\eta)dtd\eta$ is equal to the number of solutions
$(Y,Z,U,Y_1,Z_1,U_1,\ldots,Y_{q-1},Z_{q-1},U_{q-1})$ of the polynomial equations

$$\left\{\begin{array}{l} YZ+Y_1Z_1+\ldots+Y_{q-1}Z_{q-1}=0,\\ YU+Y_1U_1+\ldots+Y_{q-1}U_{q-1}=0,\end{array}\right.$$

satisfying the degree conditions
$\deg Y=k-1,\quad \deg Z\leq s-1,\quad \deg U=s+m-1,$
$\deg Y_i\leq k-2,\quad \deg Z_i=s-1,\quad \deg U_i\leq s+m-2\quad for 1\leq i\leq q-1.$
By degree considerations we have that deg $[YU+Y_1U_1+\ldots+Y_{q-1}U_{q-1}]$ is equal to k+s+m-2.
Lemma 6.13 follows.

□

Lemma 6.14. *We have for all $j\in\mathbb{Z}$ such that $0\leq j\leq\inf(2s+m-2,k-1)$.*

$$\#\left(\begin{array}{c|c} j & j\\ \hline j & j\\ \hline j & j\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} -\#\left(\begin{array}{c|c} j & j\\ \hline j & j\\ \hline j & j+1\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} =0$$

Recall that

$$\#\left(\begin{array}{c|c} j_1 & j_2\\ \hline j_3 & j_4\\ \hline j_5 & j_6\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}$$

denotes the cardinality of the following set

$$\Big\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\mid r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t,\eta))=j_1,\quad r(D^{\left[\substack{s-1\\ s+m-1}\right]\times k}(t,\eta))=j_2,$$
$$r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-1)}(t,\eta))=j_3,\quad r(D^{\left[\substack{s\\ s+m-1}\right]\times k}(t,\eta))=j_4,$$
$$r(D^{\left[\substack{s\\ s+m}\right]\times(k-1)}(t,\eta))=j_5,\quad r(D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta))=j_6\Big\}$$

for $(j_1,j_2,j_3,j_4,j_5,j_6)\in\mathbb{N}^6$.

Proof. We define

$$\mathbb{A}=\{(t,\eta)\in\mathbb{P}^2\mid r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t,\eta))=r(D^{\left[\substack{s-1\\ s+m-1}\right]\times k}(t,\eta))=r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-1)}(t,\eta))$$
$$=r(D^{\left[\substack{s\\ s+m-1}\right]\times k}(t,\eta))=r(D^{\left[\substack{s\\ s+m}\right]\times(k-1)}(t,\eta))=r(D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta))\}.$$

and

$$\mathbb{B}=\{(t,\eta)\in\mathbb{P}^2\mid r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t,\eta))=r(D^{\left[\substack{s-1\\ s+m-1}\right]\times k}(t,\eta))=r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-1)}(t,\eta))$$
$$=r(D^{\left[\substack{s\\ s+m-1}\right]\times k}(t,\eta))=r(D^{\left[\substack{s\\ s+m}\right]\times(k-1)}(t,\eta)),$$
$$r(D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta))=r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t,\eta))+1\}.$$

By (6.10) we have, observing that $\phi(t,\eta)\cdot\theta_3^{q-1}(t,\eta)$ is constant on cosets of $\mathbb{P}_{k+s-1}\times\mathbb{P}_{k+s+m-1}$

$$\int_{\mathbb{P}\times\mathbb{P}}\phi(t,\eta)\cdot\theta_3^{q-1}(t,\eta)dt\eta=\int_{\mathbb{A}}\phi(t,\eta)\cdot\theta_3^{q-1}(t,\eta)dt\eta+\int_{\mathbb{B}}\phi(t,\eta)\cdot\theta_3^{q-1}(t,\eta)dt\eta$$

$$=\sum_{(t,\eta)\in\mathbb{A}\bigcap\left(\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\right)}2{\cdot}2^{(2s+m+k-3-r(D^{\left[\begin{smallmatrix}s-1\\ s+m-1\end{smallmatrix}\right]\times(k-1)}(t,\eta))q}\int_{\mathbb{P}_{s+k-1}}dt\int_{\mathbb{P}_{s+k+m-1}}d\eta$$

$$+\sum_{(t,\eta)\in\mathbb{B}\bigcap\left(\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\right)}-2{\cdot}2^{(2s+m+k-3-r(D^{\left[\begin{smallmatrix}s-1\\ s+m-1\end{smallmatrix}\right]\times(k-1)}(t,\eta))q}\int_{\mathbb{P}_{s+k-1}}dt\int_{\mathbb{P}_{s+k+m-1}}d\eta$$

$$=$$

$$\sum_{j=0}^{\inf(2s+m-2,k-1)}2\cdot2^{(2s+m+k-3-j)q}\,.\,\#\left(\begin{array}{c|c}j&j\\\hline j&j\\\hline j&j\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}\cdot2^{-(2k+2s+m-2)}$$

$$-\sum_{j=0}^{\inf(2s+m-2,k-1)}2\cdot2^{(2s+m+k-3-j)q}\,.\,\#\left(\begin{array}{c|c}j&j\\\hline j&j\\\hline j&j+1\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}\cdot2^{-(2k+2s+m-2)}.$$

Now by Lemma 6.13 we get for all $q\geq2$

$$\sum_{j=0}^{\inf(2s+m-2,k-1)}2^{-jq}\,.\left[\#\left(\begin{array}{c|c}j&j\\\hline j&j\\\hline j&j\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}-\#\left(\begin{array}{c|c}j&j\\\hline j&j\\\hline j&j+1\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}\right]=0.$$

□

7. RANK PROPERTIES OF SUBMATRICES OF DOUBLE PERSYMMETRIC MATRICES

Consider the following partition of the matrix $D^{\left[\begin{array}{c}s-1\\ s+m-1\\\hline \alpha_{s-}\\ \beta_{s+m-}\\ \\ \end{array}\right]\times k}(t,\eta)$

$$\left(\begin{array}{ccccc|c}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1} \\
\hline
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{array}\right).$$

By studying rank properties of submatrices of the above double persymmetric matrix, we deduce by contradiction the following rank formula for all $i \in [0, \inf(2s+m-3, k-2)]$

$$\#\left(\begin{array}{c|c} i & i+1 \\ \hline i & i+1 \\ \hline i & i+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} = 0.$$

Recall that

$$\#\left(\begin{array}{c|c} i & i+1 \\ \hline i & i+1 \\ \hline i & i+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}$$

denotes the cardinality of the following set

$$\Big\{(t,\eta) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \mid r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t,\eta)) = i, \quad r(D^{\left[\substack{s-1\\ s+m-1}\right]\times k}(t,\eta)) = i+1,$$
$$r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-1)}(t,\eta)) = i, \quad r(D^{\left[\substack{s\\ s+m-1}\right]\times k}(t,\eta)) = i+1,$$
$$r(D^{\left[\substack{s\\ s+m}\right]\times(k-1)}(t,\eta)) = i, \quad r(D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta)) = i+1\Big\}.$$

Lemma 7.1. *For all i such that $0 \leq i \leq \inf(2s+m-3, k-2)$ we have*

$$\#\left(\begin{array}{c|c} i & i+1 \\ \hline i & i+1 \\ \hline i & i+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} = 0.$$

Proof. Set $(t,\eta) = (\sum_{i\geq 1}\alpha_i T^{-i}, \sum_{i\geq 1}\beta_i T^{-i}) \in \mathbb{P}\times\mathbb{P}$ *and let* $(s,k,m,j) \in \mathbb{N}^*\times\mathbb{N}^*\times\mathbb{N}\times\mathbb{N}^*$,
where $1 \leq j \leq k-1$.

We denote by $D_{j}^{\left[\substack{s\\ s+m}\right]\times(k-j+1)}(t,\eta)$ the following $(2s+m)\times(k-j+1)$ matrix

$$\left(\begin{array}{ccccc|c}
\alpha_{j} & \alpha_{j+1} & \alpha_{j+2} & \ldots & \alpha_{k-1} & \alpha_{k}\\
\beta_{j} & \beta_{j+1} & \beta_{j+2} & \ldots & \beta_{k-1} & \beta_{k}\\
\alpha_{j+1} & \alpha_{j+2} & \alpha_{j+3} & \ldots & \alpha_{k} & \alpha_{k+1}\\
\beta_{j+1} & \beta_{j+2} & \beta_{j+3} & \ldots & \beta_{k} & \beta_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\alpha_{j+s-2} & \alpha_{j+s-1} & \alpha_{j+s} & \ldots & \alpha_{k+s-3} & \alpha_{k+s-2}\\
\beta_{j+s-2} & \beta_{j+s-1} & \beta_{j+s} & \ldots & \beta_{k+s-3} & \beta_{k+s-2}\\
\beta_{j+s-1} & \beta_{j+s} & \beta_{j+s+1} & \ldots & \beta_{k+s-2} & \beta_{k+s-1}\\
\beta_{j+s} & \beta_{j+s+1} & \beta_{j+s+2} & \ldots & \beta_{k+s-1} & \beta_{k+s}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{j+s+m-3} & \beta_{j+s+m-2} & \beta_{j+s+m-1} & \ldots & \beta_{k+s+m-4} & \beta_{k+s+m-3}\\
\beta_{j+s+m-2} & \beta_{j+s+m-1} & \beta_{j+s+m} & \ldots & \beta_{k+s+m-3} & \beta_{s+m+k-2}\\
\hline
 & & & & & \\
\alpha_{j+s-1} & \alpha_{j+s} & \alpha_{j+s+1} & \ldots & \alpha_{k+s-2} & \alpha_{k+s-1}\\
\beta_{j+s+m-1} & \beta_{j+s+m} & \beta_{j+s+m+1} & \ldots & \beta_{k+s+m-2} & \beta_{k+s+m-1}
\end{array}\right).$$

Remark that after a rearrangement of the rows in the above matrix we obtain the following double persymmetric matrix where the first s rows form a $s\times(k-j+1)$ persymmetric matrix and the last s+m rows form a $(s+m)\times(k-j+1)$ persymmetric matrix with entries in $\mathbb{F}_2$

$$\left(\begin{array}{ccccc|c}
\alpha_{j} & \alpha_{j+1} & \alpha_{j+2} & \ldots & \alpha_{k-1} & \alpha_{k}\\
\alpha_{j+1} & \alpha_{j+2} & \alpha_{j+3} & \ldots & \alpha_{k} & \alpha_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\alpha_{j+s-2} & \alpha_{j+s-1} & \alpha_{j+s} & \ldots & \alpha_{k+s-3} & \alpha_{k+s-2}\\
\alpha_{j+s-1} & \alpha_{j+s} & \alpha_{j+s+1} & \ldots & \alpha_{k+s-2} & \alpha_{k+s-1}\\
\beta_{j} & \beta_{j+1} & \beta_{j+2} & \ldots & \beta_{k-1} & \beta_{k}\\
\beta_{j+1} & \beta_{j+2} & \beta_{j+3} & \ldots & \beta_{k} & \beta_{k+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{j+s-2} & \beta_{j+s-1} & \beta_{j+s} & \ldots & \beta_{k+s-3} & \beta_{k+s-2}\\
\beta_{j+s-1} & \beta_{j+s} & \beta_{j+s+1} & \ldots & \beta_{k+s-2} & \beta_{k+s-1}\\
\beta_{j+s} & \beta_{j+s+1} & \beta_{j+s+2} & \ldots & \beta_{k+s-1} & \beta_{k+s}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{j+s+m-3} & \beta_{j+s+m-2} & \beta_{j+s+m-1} & \ldots & \beta_{k+s+m-4} & \beta_{k+s+m-3}\\
\beta_{j+s+m-2} & \beta_{j+s+m-1} & \beta_{j+s+m} & \ldots & \beta_{k+s+m-3} & \beta_{k+s+m-2}\\
\beta_{j+s+m-1} & \beta_{j+s+m} & \beta_{j+s+m+1} & \ldots & \beta_{k+s+m-2} & \beta_{k+s+m-1}
\end{array}\right).$$

Proof by contradiction.
Assume on the contrary that there exists $i_0\in[0,\inf(2s+m-3,k-2)]$ such that

$$\#\left(\begin{array}{c|c}
i_0 & i_0+1\\
\hline
i_0 & i_0+1\\
\hline
i_0 & i_0+1
\end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}}>0. \tag{7.1}$$

We are going to show that

$$\#\left(\begin{array}{c|c} i_0 & i_0+1 \\ \hline i_0 & i_0+1 \\ \hline i_0 & i_0+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} > 0 \Longrightarrow \#\left(\begin{array}{c|c} i_0-1 & i_0 \\ \hline i_0-1 & i_0 \\ \hline i_0-1 & i_0 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}_1/\mathbb{P}_{k+s-1}\times\mathbb{P}_1/\mathbb{P}_{k+s+m-1}} > 0$$

$$\Longrightarrow \ldots \Longrightarrow \#\left(\begin{array}{c|c} 0 & 1 \\ \hline 0 & 1 \\ \hline 0 & 1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}_{i_0}/\mathbb{P}_{k+s-1}\times\mathbb{P}_{i_0}/\mathbb{P}_{k+s+m-1}} > 0,$$

which obviously contradicts

$$\#\left(\begin{array}{c|c} 0 & 1 \\ \hline 0 & 1 \\ \hline 0 & 1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}_{i_0}/\mathbb{P}_{k+s-1}\times\mathbb{P}_{i_0}/\mathbb{P}_{k+s+m-1}} = 0.$$

By (7.1) there exists $(t_0,\eta_0)\in\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}$ such that

$$r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t_0,\eta_0)) = r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-1)}(t_0,\eta_0)) = r(D^{\left[\substack{s\\ s+m}\right]\times(k-1)}(t_0,\eta_0)) = i_0,$$

$$r(D^{\left[\substack{s-1\\ s+m-1}\right]\times k}(t_0,\eta_0)) = r(D^{\left[\substack{s\\ s+m-1}\right]\times k}(t_0,\eta_0)) = r(D^{\left[\substack{s\\ s+m}\right]\times k}(t_0,\eta_0)) = i_0+1.$$

Set $(t_0,\eta_0)=(\sum_{i\geq 1}\alpha_i T^{-i},\sum_{i\geq 1}\beta_i T^{-i})$, and consider the following partition of the matrix $D^{\left[\substack{s\\ s+m}\right]\times k}(t_0,\eta_0)$

$$\left(\begin{array}{c|ccccc|c}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\hline
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{k+s-3} & \alpha_{k+s-2} \\
\beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{k+s-3} & \beta_{k+s-2} \\
\beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{k+s-2} & \beta_{k+s-1} \\
\beta_{s+1} & \beta_{s+2} & \beta_{s+3} & \ldots & \beta_{k+s-1} & \beta_{k+s} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-2} & \beta_{s+m-1} & \beta_{s+m} & \ldots & \beta_{k+s+m-4} & \beta_{k+s+m-3} \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{k+s+m-3} & \beta_{k+s+m-2} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{k+s-2} & \alpha_{k+s-1} \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{k+s+m-2} & \beta_{k+s+m-1}
\end{array}\right).$$

By (7.1) we have $r(D_2^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t_0,\eta_0))\leq i_0$ *and* $r(D^{\left[\substack{s-1\\ s+m-1}\right]\times k}(t_0,\eta_0)) = i_0+1$.

It follows that $r(D_2^{\left[\substack{s-1\\ s+m-1}\right]\times(k-1)}(t_0,\eta_0)) = i_0$.

Let $a_1, a_2, \ldots, a_k$ denote the columns of $D^{\left[\overset{s-1}{s+m-1}\right]\times k}(t_0, \eta_0)$ that is,

$$a_j = \begin{pmatrix} \alpha_j \\ \beta_j \\ \alpha_{j+1} \\ \beta_{j+1} \\ \vdots \\ \alpha_{j+s-2} \\ \beta_{j+s-2} \\ \beta_{j+s-1} \\ \beta_{j+s} \\ \vdots \\ \beta_{j+s+m-3} \\ \beta_{j+s+m-2} \end{pmatrix}.$$

Since $r(D_2^{\left[\overset{s-1}{s+m-1}\right]\times (k-1)}(t_0, \eta_0)) = i_0 \; and \; r(D^{\left[\overset{s-1}{s+m-1}\right]\times k}(t_0, \eta_0)) = i_0 + 1$, we have $a_1 \notin span \left\{a_2, a_3, \ldots, a_k\right\}$, therefore $r(D_2^{\left[\overset{s-1}{s+m-1}\right]\times (k-2)}(t_0, \eta_0)) = i_0 - 1$. We have then

$$r(D_2^{\left[\overset{s-1}{s+m-1}\right]\times (k-2)}(t_0, \eta_0)) = i_0 - 1, \tag{7.2}$$

$$r(D^{\left[\overset{s-1}{s+m-1}\right]\times (k-1)}(t_0, \eta_0)) = i_0, \tag{7.3}$$

$$r(D^{\left[\overset{s}{s+m-1}\right]\times (k-1)}(t_0, \eta_0)) = i_0, \tag{7.4}$$

$$r(D^{\left[\overset{s}{s+m}\right]\times (k-1)}(t_0, \eta_0)) = i_0 \qquad \text{and, see above,} \tag{7.5}$$

$$r(D_2^{\left[\overset{s-1}{s+m-1}\right]\times (k-1)}(t_0, \eta_0)) = i_0. \tag{7.6}$$

Now we consider the matrix obtained from the matrix $D^{\left[\overset{s}{s+m}\right]\times k}(t_0,\eta_0)$ by delating the first column and replacing the last column by the first one :

$$\left(\begin{array}{cccc|c}
\alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_1 \\
\beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_1 \\
\alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_2 \\
\beta_3 & \beta_4 & \ldots & \beta_k & \beta_2 \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_s & \alpha_{s+1} & \ldots & \alpha_{k+s-3} & \alpha_{s-1} \\
\beta_s & \beta_{s+1} & \ldots & \beta_{k+s-3} & \beta_{s-1} \\
\beta_{s+1} & \beta_{s+2} & \ldots & \beta_{k+s-2} & \beta_s \\
\beta_{s+2} & \beta_{s+3} & \ldots & \beta_{k+s-1} & \beta_{s+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \ldots & \beta_{k+s+m-4} & \beta_{s+m-2} \\
\beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{k+s+m-3} & \beta_{s+m-1} \\
\hline
\alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{k+s-2} & \alpha_s \\
\beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{k+s+m-2} & \beta_{s+m}
\end{array}\right).$$

From (7.2), (7.3), (7.4) and (7.5) we obtain by elementary rank considerations

$$r(D_2^{\left[\overset{s}{s+m-1}\right]\times (k-2)}(t_0,\eta_0)) = i_0 - 1, \tag{7.7}$$

$$r(D_2^{\left[\overset{s}{s+m}\right]\times (k-2)}(t_0,\eta_0)) = i_0 - 1. \tag{7.8}$$

Consider now the matrix $D_2^{\left[\overset{s}{s+m}\right]\times (k-1)}(t_0,\eta_0)$ obtained by the matrix $D^{\left[\overset{s}{s+m}\right]\times k}(t_0,\eta_0)$ by delating the first column, that is

$$\left(\begin{array}{cccc|c}
\alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_s & \alpha_{s+1} & \ldots & \alpha_{k+s-3} & \alpha_{k+s-2} \\
\beta_s & \beta_{s+1} & \ldots & \beta_{k+s-3} & \beta_{k+s-2} \\
\beta_{s+1} & \beta_{s+2} & \ldots & \beta_{k+s-2} & \beta_{k+s-1} \\
\beta_{s+2} & \beta_{s+3} & \ldots & \beta_{k+s-1} & \beta_{k+s} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \ldots & \beta_{k+s+m-4} & \beta_{k+s+m-3} \\
\beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{k+s+m-3} & \beta_{k+s+m-2} \\
\hline
\alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{k+s-2} & \alpha_{k+s-1} \\
\beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{k+s+m-2} & \beta_{k+s+m-1}
\end{array}\right).$$

From (7.2), (7.6), (7.7) and (7.8), we obtain

$$r(D_2^{\left[\overset{s}{s+m-1}\right]\times(k-1)}(t_0,\eta_0)) = r(D_2^{\left[\overset{s}{s+m}\right]\times(k-1)}(t_0,\eta_0)) = i_0.$$

We get

$$\#\left(\begin{array}{c|c} i_0-1 & i_0 \\ \hline i_0-1 & i_0 \\ \hline i_0-1 & i_0 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}_1/\mathbb{P}_{k+s-1}\times\mathbb{P}_1/\mathbb{P}_{k+s+m-1}} > 0.$$

Recall that

$$\#\left(\begin{array}{c|c} i_0-1 & i_0 \\ \hline i_0-1 & i_0 \\ \hline i_0-1 & i_0 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}_1/\mathbb{P}_{k+s-1}\times\mathbb{P}_1/\mathbb{P}_{k+s+m-1}}$$

denotes the cardinality of the following set

$$\Big\{(t,\eta)\in\mathbb{P}_1/\mathbb{P}_{k+s-1}\times\mathbb{P}_1/\mathbb{P}_{k+s+m-1} \mid r(D_2^{\left[\overset{s-1}{s+m-1}\right]\times(k-2)}(t,\eta)) = i_0-1, \quad r(D_2^{\left[\overset{s-1}{s+m-1}\right]\times(k-1)}(t,\eta)) = i_0,$$
$$r(D_2^{\left[\overset{s}{s+m-1}\right]\times(k-2)}(t,\eta)) = i_0-1, \quad r(D_2^{\left[\overset{s}{s+m-1}\right]\times(k-1)}(t,\eta)) = i_0,$$
$$r(D_2^{\left[\overset{s}{s+m}\right]\times(k-2)}(t,\eta)) = i_0-1, \quad r(D_2^{\left[\overset{s}{s+m}\right]\times(k-1)}(t,\eta)) = i_0\Big\}.$$

We have now proved that

$$\#\left(\begin{array}{c|c} i_0 & i_0+1 \\ \hline i_0 & i_0+1 \\ \hline i_0 & i_0+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} > 0 \Longrightarrow \#\left(\begin{array}{c|c} i_0-1 & i_0 \\ \hline i_0-1 & i_0 \\ \hline i_0-1 & i_0 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}_1/\mathbb{P}_{k+s-1}\times\mathbb{P}_1/\mathbb{P}_{k+s+m-1}} > 0.$$

We repeat this procedure and obtain after finitely many steps

$$\#\left(\begin{array}{c|c} 0 & 1 \\ \hline 0 & 1 \\ \hline 0 & 1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}_{i_0}/\mathbb{P}_{k+s-1}\times\mathbb{P}_{i_0}/\mathbb{P}_{k+s+m-1}} > 0. \tag{7.9}$$

From

$$D^{\left[\substack{s\\ s+m}\right]\times(k-i_0)}_{i_0+1}(t_0,\eta_0) = \left(\begin{array}{cccccc|c}
\alpha_{i_0+1} & \alpha_{i_0+2} & \alpha_{i_0+3} & \ldots & \alpha_{k-1} & & \alpha_k \\
\beta_{i_0+1} & \beta_{i_0+2} & \beta_{i_0+3} & \ldots & \beta_{k-1} & & \beta_k \\
\alpha_{i_0+2} & \alpha_{i_0+3} & \alpha_{i_0+4} & \ldots & \alpha_k & & \alpha_{k+1} \\
\beta_{i_0+2} & \beta_{i_0+3} & \beta_{i_0+4} & \ldots & \beta_k & & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots \\
\alpha_{i_0+s-1} & \alpha_{i_0+s} & \alpha_{i_0+s+1} & \ldots & \alpha_{k+s-3} & & \alpha_{k+s-2} \\
\beta_{i_0+s} & \beta_{i_0+s+1} & \beta_{i_0+s+2} & \ldots & \beta_{k+s-3} & & \beta_{k+s-2} \\
\beta_{i_0+s+1} & \beta_{i_0+s+2} & \beta_{i_0+s+3} & \ldots & \beta_{k+s-2} & & \beta_{k+s-1} \\
\beta_{i_0+s+2} & \beta_{i_0+s+3} & \beta_{i_0+s+4} & \ldots & \beta_{k+s-1} & & \beta_{k+s} \\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots \\
\beta_{i_0+s+m-2} & \beta_{i_0+s+m-1} & \beta_{i_0+s+m} & \ldots & \beta_{k+s+m-4} & & \beta_{k+s+m-3} \\
\beta_{i_0+s+m-1} & \beta_{i_0+s+m} & \beta_{i_0+s+m+1} & \ldots & \beta_{k+s+m-3} & & \beta_{s+m+k-2} \\
\hline
\alpha_{i_0+s} & \alpha_{i_0+s+1} & \alpha_{i_0+s+2} & \ldots & \alpha_{k+s-2} & & \alpha_{k+s-1} \\
\beta_{i_0+s+m} & \beta_{i_0+s+m+1} & \beta_{i_0+s+m+2} & \ldots & \beta_{k+s+m-2} & & \beta_{k+s+m-1},
\end{array}\right).$$

we obviously get

$$\#\left(\begin{array}{c|c} 0 & 1 \\ \hline 0 & 1 \\ \hline 0 & 1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}_{i_0}/\mathbb{P}_{k+s-1}\times\mathbb{P}_{i_0}/\mathbb{P}_{k+s+m-1}} = 0$$

which clearly contradicts (7.9).

Recall that

$$\#\left(\begin{array}{c|c} 0 & 1 \\ \hline 0 & 1 \\ \hline 0 & 1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}_{i_0}/\mathbb{P}_{k+s-1}\times\mathbb{P}_{i_0}/\mathbb{P}_{k+s+m-1}}$$

denotes the cardinality of the following set

$$\Big\{(t,\eta)\in\mathbb{P}_{i_0}/\mathbb{P}_{k+s-1}\times\mathbb{P}_{i_0}/\mathbb{P}_{k+s+m-1} \mid r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-i_0-1)}(t,\eta)) = 0, \quad r(D^{\left[\substack{s-1\\ s+m-1}\right]\times(k-i_0)}(t,\eta)) = 1,$$
$$r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-i_0-1)}(t,\eta)) = 0, \quad r(D^{\left[\substack{s\\ s+m-1}\right]\times(k-i_0)}(t,\eta)) = 1,$$
$$r(D^{\left[\substack{s\\ s+m}\right]\times(k-i_0-1)}(t,\eta)) = 0, \quad r(D^{\left[\substack{s\\ s+m}\right]\times(k-i_0)}(t,\eta)) = 1\Big\}.$$

□

Lemma 7.2. *We have*

$$\#\left(\begin{array}{c|c} i & i+1 \\ \hline i & i+1 \\ \hline i & i+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} = 0 \quad \textit{if } 0\le i\le \inf(2s+m-3,k-2), \tag{7.10}$$

(7.11)

$$\#\left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} = \#\left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} \quad \textit{if } 0 \leq i \leq \inf(2s+m-2, k-1),$$

(7.12)

$$\#\left(\begin{array}{c|c} i & i \\ \hline i & i+1 \\ \hline i & i+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} = 2\cdot\#\left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} \quad \textit{if } 0 \leq i \leq \inf(2s+m-2, k-1).$$

Proof. The Lemma follows respectively from Lemmas 7.1, 6.14 and 6.10. ☐

8. STUDY OF THE REMAINDER $\Delta_i^{\left[\substack{s \\ s+m}\right]\times k}$ IN THE RECURRENT FORMULA

From the rank formulas established in sections 6 and 7, we deduce by elementary rank considerations the following formula for $1 \leq i \leq 2s+m,\ k \geq i+1$

$$\Delta_i^{\left[\substack{s \\ s+m}\right]\times k} = \sum_{j=i-2}^{i+1} a_j \cdot \Gamma_j^{\left[\substack{s-1 \\ s-1+m}\right]\times j}$$

where the $a_j \in \mathbb{Z}$ are explicitely determined.
We get

$$\Delta_i^{\left[\substack{s \\ s+m}\right]\times k} = \Delta_i^{\left[\substack{s \\ s+m}\right]\times (i+1)} \quad \textit{whenever } k \geq i+1.$$

Let $(j_1, j_2, j_3) \in \mathbb{N}^3$ recall that we define

$$\sigma_{j_1,j_2,j_3}^{\left[\begin{array}{c} \frac{\substack{s-1 \\ s+m-1}}{\substack{\alpha_{s-} \\ \beta_{s+m-}}} \\ \\ \end{array}\right]\times k}$$

to be the cardinality of the following set

$$\Big\{(t,\eta) \in \mathbb{P}/\mathbb{P}_{k+s-1} \times \mathbb{P}/\mathbb{P}_{k+s+m-1} \mid r(D^{\left[\substack{s-1 \\ s+m-1}\right]\times k}(t,\eta)) = j_1, \quad r(D^{\left[\substack{s \\ s+m-1}\right]\times k}(t,\eta)) = j_2,$$
$$r(D^{\left[\substack{s \\ s+m}\right]\times k}(t,\eta)) = j_3\Big\}.$$

Lemma 8.1. *For* $1 \leq i \leq 2s+m-3, \quad k \geq i+1$ *we have*

$$\sigma_{i,i,i}^{\left[\begin{smallmatrix} s-1 \\ \overline{s+m-1} \\ \overline{\alpha_{s-}} \\ \beta_{s+m-} \\ \\ \\ \end{smallmatrix}\right]\times k} = \sigma_{i,i,i}^{\left[\begin{smallmatrix} s-1 \\ \overline{s+m-1} \\ \overline{\alpha_{s-}} \\ \beta_{s+m-} \\ \\ \\ \end{smallmatrix}\right]\times (i+1)} . \tag{8.1}$$

Proof. The formula (8.1) is obvious for k = i+1.

We consider the following partition of the matrix

$$D^{\left[\begin{smallmatrix} s-1 \\ \overline{s+m-1} \\ \overline{\alpha_{s-}} \\ \beta_{s+m-} \\ \\ \\ \end{smallmatrix}\right]\times k}(t,\eta)$$

$$\left(\begin{array}{ccccc|c}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1} \\
\hline
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{array}\right).$$

Let $k \geq i+2$. Using Lemma 7.1, we obtain by elementary rank considerations

$$\begin{aligned}
\sigma_{i,i,i}^{\left[\begin{smallmatrix} s-1 \\ \overline{s+m-1} \\ \overline{\alpha_{s-}} \\ \beta_{s+m-} \\ \\ \\ \end{smallmatrix}\right]\times k} &= \#\left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} + \#\left(\begin{array}{c|c} i-1 & i \\ \hline i & i \\ \hline i & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} \\
&+ \#\left(\begin{array}{c|c} i-1 & i \\ \hline i-1 & i \\ \hline i & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} + \#\left(\begin{array}{c|c} i-1 & i \\ \hline i-1 & i \\ \hline i-1 & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} \\
&= \#\left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} .
\end{aligned} \tag{8.2}$$

Further, by Lemma 7.2 we have

(8.3)

$$
\begin{aligned}
4 \cdot \sigma_{i,i,i}^{\left[\begin{matrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \end{matrix}\right] \times (k-1)} &= {}^{\#}\left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} + {}^{\#}\left(\begin{array}{c|c} i & i \\ \hline i & i+1 \\ \hline i & i+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} \\
&+ {}^{\#}\left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} + {}^{\#}\left(\begin{array}{c|c} i & i+1 \\ \hline i & i+1 \\ \hline i & i+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} \\
&= 4 \cdot {}^{\#}\left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} .
\end{aligned}
$$

By (8.2), (8.3) we deduce

$$
\sigma_{i,i,i}^{\left[\begin{matrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \end{matrix}\right] \times (k-1)} = \sigma_{i,i,i}^{\left[\begin{matrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \end{matrix}\right] \times k} \qquad for\ all\ k \geq i+2. \tag{8.4}
$$

Hence by (8.4) we obtain successively

$$
\sigma_{i,i,i}^{\left[\begin{matrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \end{matrix}\right] \times k} = \sigma_{i,i,i}^{\left[\begin{matrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \end{matrix}\right] \times (k-1)} = \sigma_{i,i,i}^{\left[\begin{matrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \end{matrix}\right] \times (k-2)} = \ldots = \sigma_{i,i,i}^{\left[\begin{matrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \end{matrix}\right] \times (i+1)} .
$$

□

Lemma 8.2. *For all i such that $1 \leq i \leq 2s+m-3$ we have*

$$
\sigma_{i,i,i}^{\left[\begin{matrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \end{matrix}\right] \times (i+1)} = 4 \cdot \Gamma_{i}^{\left[\begin{matrix} s-1 \\ s-1+m \end{matrix}\right] \times i} - \Gamma_{i+1}^{\left[\begin{matrix} s-1 \\ s-1+m \end{matrix}\right] \times (i+1)} . \tag{8.5}
$$

Proof. Consider the following partition of the matrix

$$
D^{\left[\begin{matrix} s-1 \\ \underline{s+m-1} \\ \alpha_{s-} \\ \beta_{s+m-} \\ \\ \end{matrix}\right] \times (i+1)} \qquad (t, \eta)
$$

$$\left(\begin{array}{cccccc|c}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_i & & \alpha_{i+1} \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{1+1} & & \alpha_{i+2} \\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{i+s-2} & & \alpha_{i+s-1} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_i & & \beta_{i+1} \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{i+1} & & \beta_{i+2} \\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{i+m} & & \beta_{i+m+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+i-2} & & \beta_{s+m+i-1} \\
\hline
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+i-1} & & \alpha_{s+i} \\
\hline
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+i-1} & & \beta_{s+m+i}
\end{array}\right).$$

By elementary rank considerations and using Lemma 7.1 we obtain

(8.6)

$$\begin{aligned}
\sigma_{i,i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right]\times(i+1)}
&= {}^{\#}\left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{i+s}\times\mathbb{P}/\mathbb{P}_{i+s+m}}
+ {}^{\#}\left(\begin{array}{c|c} i-1 & i \\ \hline i & i \\ \hline i & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{i+s}\times\mathbb{P}/\mathbb{P}_{i+s+m}} \\
&+ {}^{\#}\left(\begin{array}{c|c} i-1 & i \\ \hline i-1 & i \\ \hline i & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{i+s}\times\mathbb{P}/\mathbb{P}_{i+s+m}}
+ {}^{\#}\left(\begin{array}{c|c} i-1 & i \\ \hline i-1 & i \\ \hline i-1 & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{i+s}\times\mathbb{P}/\mathbb{P}_{i+s+m}} \\
&= {}^{\#}\left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{i+s}\times\mathbb{P}/\mathbb{P}_{i+s+m}}.
\end{aligned}$$

Further, by Lemma 7.2 we have

(8.7)

$$\begin{aligned}
4\cdot\sigma_{i,i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right]\times i}
&= 4\cdot 4\cdot\Gamma_i^{\left[\begin{array}{c} s-1 \\ s-1+m \end{array}\right]\times i} \\
&= {}^{\#}\left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{i+s}\times\mathbb{P}/\mathbb{P}_{i+s+m}}
+ {}^{\#}\left(\begin{array}{c|c} i & i \\ \hline i & i+1 \\ \hline i & i+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{i+s}\times\mathbb{P}/\mathbb{P}_{i+s+m}} \\
&+ {}^{\#}\left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{i+s}\times\mathbb{P}/\mathbb{P}_{i+s+m}}
+ {}^{\#}\left(\begin{array}{c|c} i & i+1 \\ \hline i & i+1 \\ \hline i & i+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{i+s}\times\mathbb{P}/\mathbb{P}_{i+s+m}} \\
&= 4\cdot{}^{\#}\left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{i+s}\times\mathbb{P}/\mathbb{P}_{i+s+m}}
+ {}^{\#}\left(\begin{array}{c|c} i & i+1 \\ \hline i & i+1 \\ \hline i & i+1 \end{array}\right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{i+s}\times\mathbb{P}/\mathbb{P}_{i+s+m}}
\end{aligned}$$

$$= 4 \cdot \# \left(\begin{array}{c|c} i & i \\ \hline i & i \\ \hline i & i \end{array} \right)^{\frac{\alpha}{\beta}}_{\mathbb{P}/\mathbb{P}_{i+s} \times \mathbb{P}/\mathbb{P}_{i+s+m}} + 4 \cdot \Gamma_{i+1}^{\left[\substack{s-1 \\ s-1+m}\right] \times (i+1)}.$$

Combining (8.6), (8.7) we deduce

$$16 \cdot \Gamma_{i}^{\left[\substack{s-1 \\ s-1+m}\right] \times i} = 4 \cdot \sigma_{i,i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right] \times (i+1)} + 4 \cdot \Gamma_{i+1}^{\left[\substack{s-1 \\ s-1+m}\right] \times (i+1)} \tag{8.8}$$

and (8.5) is established. ☐

Lemma 8.3. *Let $s \geq 2$ and $m \geq 0$, we have in the following two cases :*

$\underline{\textit{The case } 1 \leq k \leq 2s+m-2}$

$$\sigma_{i,i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right] \times k} = \begin{cases} 1 & \textit{if } i = 0, \quad k \geq 1, \\ 4 \cdot \Gamma_{i}^{\left[\substack{s-1 \\ s-1+m}\right] \times i} - \Gamma_{i+1}^{\left[\substack{s-1 \\ s-1+m}\right] \times (i+1)} & \textit{if } 1 \leq i \leq k-1, \\ 4 \cdot \Gamma_{k}^{\left[\substack{s-1 \\ s-1+m}\right] \times k} & \textit{if } i = k. \end{cases} \tag{8.9}$$

$\underline{\textit{The case } k \geq 2s+m-2}$

$$\sigma_{i,i,i}^{\left[\begin{array}{c} s-1 \\ s+m-1 \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right] \times k} = \begin{cases} 1 & \textit{if } i = 0, \\ 4 \cdot \Gamma_{i}^{\left[\substack{s-1 \\ s-1+m}\right] \times i} - \Gamma_{i+1}^{\left[\substack{s-1 \\ s-1+m}\right] \times (i+1)} & \textit{if } 1 \leq i \leq 2s+m-3, \\ 4 \cdot \Gamma_{2s+m-2}^{\left[\substack{s-1 \\ s-1+m}\right] \times (2s+m-2)} & \textit{if } i = 2s+m-2. \end{cases} \tag{8.10}$$

Proof. Combining (8.1) and (8.5) and recalling the definition of $\sigma_{i,i,i}^{\left[\begin{array}{c} s \\ s+m \\ \hline \alpha_{s-} \\ \beta_{s+m-} \end{array}\right] \times k}$ we deduce easily Lemma 8.3. ☐

Lemma 8.4. *The remainder $\Delta_{i}^{\left[\substack{s \\ s+m}\right] \times k}$ in the recurrent formula is equal to*

(8.11)

$$\begin{cases}
1 & \text{if } i=0,\ k\geq 1,\\
4\cdot\Gamma_{1}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times 1}-\Gamma_{2}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times 2}-3 & \text{if } i=1,\ k\geq 2,\\
4\cdot\Gamma_{1}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times 1}-3 & \text{if } i=1,\ k=1,\\
7\cdot\Gamma_{2}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times 2}-12\cdot\Gamma_{1}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times 1}-\Gamma_{3}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times 3}+2 & \text{if } i=2,\ k\geq 3,\\
7\cdot\Gamma_{2}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times 2}-12\cdot\Gamma_{1}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times 1}+2 & \text{if } i=2,\ k=2,\\
7\cdot\Gamma_{i}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times i}-14\cdot\Gamma_{i-1}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times (i-1)}+8\cdot\Gamma_{i-2}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times (i-2)}-\Gamma_{i+1}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times (i+1)} & \text{if } 3\leq i\leq 2s+m-3, k\geq i+1,\\
7\cdot\Gamma_{i}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times i}-14\cdot\Gamma_{i-1}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times (i-1)}+8\cdot\Gamma_{i-2}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times (i-2)} & \text{if } 3\leq i\leq 2s+m-3,\ k=i,\\
7\cdot\Gamma_{2s+m-2}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times (2s+m-2)}-14\cdot\Gamma_{2s+m-3}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times (2s+m-3)}+8\cdot\Gamma_{2s+m-4}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times (2s+m-4)} & \text{if } i=2s+m-2,\ k\geq i,\\
-14\cdot\Gamma_{2s+m-2}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times (2s+m-2)}+8\cdot\Gamma_{2s+m-3}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times (2s+m-3)} & \text{if } i=2s+m-1,\ k\geq i,\\
8\cdot\Gamma_{2s+m-2}^{\left[\begin{smallmatrix}s-1\\ s-1+m\end{smallmatrix}\right]\times (2s+m-2)} & \text{if } i=2s+m,\ k\geq i.
\end{cases}$$

Proof. From (8.9) and (8.10) and recalling that $\Delta_{i}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times k}$ is equal to

$$\sigma_{i,i,i}^{\left[\begin{array}{c}s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\end{array}\right]\times k}-3\cdot\sigma_{i-1,i-1,i-1}^{\left[\begin{array}{c}s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\end{array}\right]\times k}+2\cdot\sigma_{i-2,i-2,i-2}^{\left[\begin{array}{c}s-1\\ s+m-1\\ \hline \alpha_{s-}\\ \beta_{s+m-}\end{array}\right]\times k}$$

we deduce easily Lemma 8.4. □

Lemma 8.5. *We have for all i such that $0\leq i\leq 2s+m-3$ and for all $k\geq i+1$*

$$\Delta_{i}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times k}=\Delta_{i}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times (i+1)}, \tag{8.12}$$

$$\Delta_{i}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times k}=\Delta_{i}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times i}\quad for\ i\in\{2s+m-2,2s+m-1,2s+m\},\ k\geq i. \tag{8.13}$$

Proof. Follows immediately from Lemma 8.4.

□

9. A RECURRENT FORMULA FOR THE DIFFERENCE $\Gamma_{i}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times (k+1)}-\Gamma_{i}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times k}, 1\leq i\leq 2s+m,\ k\geq i+1$

In this section we deduce from the recurrent formula in Lemma 5.14 and from the fact that the remainder $\Delta_{i}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times k}$ is independent of k if $k\geq i+1$, the following recurrent formula

$$\Gamma_{i}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times (k+1)}-\Gamma_{i}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times k}=4\cdot\left[\Gamma_{i-1}^{\left[\begin{smallmatrix}s\\ s+(m-1)\end{smallmatrix}\right]\times (k+1)}-\Gamma_{i-1}^{\left[\begin{smallmatrix}s\\ s+(m-1)\end{smallmatrix}\right]\times k}\right]$$

$+R(i,s,m,k)$ if $0\leq i\leq \inf(2s+m,k-1)$, where R(i,s,m,k) is equal to

$$\begin{cases} 0 & \text{if} \quad 1 \leq i \leq s-1,\ k \geq i+1, \\ 2^{s-1}\left[\Gamma_{j+1}^{\left[{1 \atop 1+(s+m-1)}\right]\times(k+1)} - \Gamma_{j+1}^{\left[{1 \atop 1+(s+m-1)}\right]\times k}\right] & \\ -2^{s+1}\left[\Gamma_{j}^{\left[{1 \atop 1+(s+m-2)}\right]\times(k+1)} - \Gamma_{j}^{\left[{1 \atop 1+(s+m-2)}\right]\times k}\right] & \text{if} \quad i = s+j, \quad 0 \leq j \leq \inf(s+m, k-s-1). \end{cases}$$

If i = s+j we set R(i,s,m,k) = R(j,s,m,k) where $0 \leq j \leq \inf(s+m, k-s-1)$.

Lemma 9.1. *Let* $s \geq 2, m \geq 0$ *and* $k \geq 1$, *then* $\Gamma_{i}^{\left[{s \atop s+m}\right]\times k} - 4\cdot\Gamma_{i-1}^{\left[{s \atop s+(m-1)}\right]\times k}$ *is given by*

$$\begin{cases} 5\cdot 2^{i-1} + \sum_{j=0}^{i-2} 2^{j}\Delta_{i-j}^{\left[{s-j \atop s-j+(m+j)}\right]\times k} & \text{if } 2 \leq i \leq \inf(s-1,k), \\ 2^{s-1}\left[\Gamma_{i-(s-1)}^{\left[{1 \atop 1+m+(s-1)}\right]\times k} - 4\cdot\Gamma_{i-s}^{\left[{1 \atop 1+m+(s-2)}\right]\times k}\right] + \sum_{j=0}^{s-2} 2^{j}\Delta_{i-j}^{\left[{s-j \atop s-j+(m+j)}\right]\times k} & \text{if } s \leq i \leq \inf(k, 2s+m). \end{cases} \tag{9.1}$$

Proof. The first case $2 \leq i \leq \inf(s-1,k)$

Set for $1 \leq i \leq \inf(2s+m,k)$

$$\Omega_i(s,m,k) = \Gamma_{i}^{\left[{s \atop s+m}\right]\times k} - 4\cdot\Gamma_{i-1}^{\left[{s \atop s+(m-1)}\right]\times k}$$

so

$$\Omega_{i-1}(s-1,m+1,k) = \Gamma_{i-1}^{\left[{s-1 \atop s-1+(m+1)}\right]\times k} - 4\cdot\Gamma_{i-2}^{\left[{s-1 \atop s-1+m}\right]\times k}.$$

We obtain then by Lemma 5.14

$$\Gamma_{i}^{\left[{s \atop s+m}\right]\times k} = 2\cdot\Gamma_{i-1}^{\left[{s-1 \atop s-1+(m+1)}\right]\times k} + 4\cdot\Gamma_{i-1}^{\left[{s \atop s+(m-1)}\right]\times k} - 8\cdot\Gamma_{i-2}^{\left[{s-1 \atop s-1+m}\right]\times k} + \Delta_{i}^{\left[{s \atop s+m}\right]\times k}, \tag{9.2}$$

$$\iff \Omega_i(s,m,k) = 2\cdot\Omega_{i-1}(s-1,m+1,k) + \Delta_{i}^{\left[{s \atop s+m}\right]\times k}. \tag{9.3}$$

By (9.3) we obtain succesively

$\Omega_i(s,m,k) = 2\cdot\Omega_{i-1}(s-1,m+1,k) + \Delta_{i}^{\left[{s \atop s+m}\right]\times k}$

$2\cdot\Omega_{i-1}(s-1,m+1,k) = 2\cdot\left(2\cdot\Omega_{i-2}(s-2,m+2,k) + \Delta_{i-1}^{\left[{s-1 \atop s-1+(m+1)}\right]\times k}\right)$

$$\vdots$$

$$2^j \cdot \Omega_{i-j}(s-j, m+j, k) = 2^j \cdot \left(2 \cdot \Omega_{i-(j+1)}(s-(j+1), m+(j+1), k) + \Delta_{i-j}^{\left[\substack{s-j \\ s-j+(m+j)}\right]\times k} \right)$$

$$\vdots$$

$$2^{i-2} \cdot \Omega_{i-(i-2)}(s-(i-2), m+(i-2), k) = 2^{i-2} \cdot \left(2 \cdot \Omega_{i-(i-1)}(s-(i-1), m+(i-1), k) + \Delta_{i-(i-2)}^{\left[\substack{s-(i-2) \\ s-(i-2)+(m+(i-2))}\right]\times k} \right)$$

By summing the above equations we get

$$\begin{aligned}
\sum_{j=0}^{i-2} 2^j \cdot \Omega_{i-j}(s-j, m+j, k) &= \sum_{j=0}^{i-2} 2^{j+1} \cdot \Omega_{i-(j+1)}(s-(j+1), m+(j+1), k) + \sum_{j=0}^{i-2} 2^j \cdot \Delta_{i-j}^{\left[\substack{s-j \\ s-j+(m+j)}\right]\times k} \\
&= \sum_{j=1}^{i-1} 2^j \cdot \Omega_{i-j}(s-j, m+j, k) + \sum_{j=0}^{i-2} 2^j \cdot \Delta_{i-j}^{\left[\substack{s-j \\ s-j+(m+j)}\right]\times k}.
\end{aligned} \tag{9.4}$$

By (9.4) we get after some simplifications

$$\Omega_i(s, m, k) = 2^{i-1} \cdot \Omega_1(s-(i-1), m+(i-1), k) + \sum_{j=0}^{i-2} 2^j \cdot \Delta_{i-j}^{\left[\substack{s-j \\ s-j+(m+j)}\right]\times k}. \tag{9.5}$$

By the definition of $\Omega_i(s, m, k)$ we have $\Omega_1(s-(i-1), m+(i-1), k) = \Gamma_1^{\left[\substack{s-i+1 \\ s+m}\right]\times k} - 4 \cdot \Gamma_0^{\left[\substack{s-i+1 \\ s+(m-1)}\right]\times k}$.

Recall that $D^{\left[\substack{s-i+1 \\ s+m}\right]\times k}(t, \eta) = \left[\frac{D_{(s-i+1)\times k}(t)}{D_{(s+m)\times k}(\eta)}\right]$ denotes the following $(2s+m-i+1)\times k$ matrix where the first (s -i +1) rows form a $(s-i+1)\times k$ persymmetric matrix and the last (s+m) rows form a $(s+m)\times k$ persymmetric matrix with entries in $\mathbb{F}_2$

$$\left(\begin{array}{cccccc} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-i+1} & \alpha_{s-i+2} & \alpha_{s-i+3} & \dots & \alpha_{s-i+k-1} & \alpha_{s-i+k} \\ \hline \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \end{array}\right).$$

Then obviously we have

$$\begin{aligned}
&\Gamma_1^{\left[\substack{s-i+1\\ s+m}\right]\times k} \\
&= Card\left\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-i-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\mid r(D^{\left[\substack{s-i+1\\ s+m}\right]\times k}(t,\eta))=1\right\} \\
&= Card\left\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-i-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\mid r(D_{(s-i+1)\times k}(t))=0,\quad r(D_{(s+m)\times k}(\eta))=1\right\} \\
&+ Card\left\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-i-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\mid r(D_{(s-i+1)\times k}(t))=1,\quad r(D_{(s+m)\times k}(\eta))=0\right\} \\
&+ Card\left\{(t,\eta)\in\mathbb{P}/\mathbb{P}_{k+s-i-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}\mid r(D_{(s-i+1)\times k}(t))=1,\ r(D_{(s+m)\times k}(\eta))=1,\ r(D^{\left[\substack{s-i+1\\ s+m}\right]\times k}(t,\eta))=1\right\} \\
&= 3+3+3=9.
\end{aligned}$$

And (9.1) follows in the first case.

The second case $s\leq i\leq \inf(2s+m,k)$

We proceed as in the first case.

Again by (9.3) we obtain succesively

$$\Omega_i(s,m,k)=2\cdot\Omega_{i-1}(s-1,m+1,k)+\Delta_i^{\left[\substack{s\\ s+m}\right]\times k}$$

$$2\cdot\Omega_{i-1}(s-1,m+1,k)=2\cdot\left(2\cdot\Omega_{i-2}(s-2,m+2,k)+\Delta_{i-1}^{\left[\substack{s-1\\ s-1+(m+1)}\right]\times k}\right)$$

$\vdots$

$$2^j\cdot\Omega_{i-j}(s-j,m+j,k)=2^j\cdot\left(2\cdot\Omega_{i-(j+1)}(s-(j+1),m+(j+1),k)+\Delta_{i-j}^{\left[\substack{s-j\\ s-j+(m+j)}\right]\times k}\right)$$

$\vdots$

$$2^{s-2}\cdot\Omega_{i-(s-2)}(s-(s-2),m+(s-2),k)=2^{s-2}\cdot\left(2\cdot\Omega_{i-(s-1)}(s-(s-1),m+(s-1),k)+\Delta_{i-(s-2)}^{\left[\substack{s-(s-2)\\ s+m}\right]\times k}\right).$$

By summing the above equations we obtain

$$\begin{aligned}\sum_{j=0}^{s-2}2^{j}\cdot\Omega_{i-j}(s-j,m+j,k)&=\sum_{j=0}^{s-2}2^{j+1}\cdot\Omega_{i-(j+1)}(s-(j+1),m+(j+1),k)+\sum_{j=0}^{s-2}2^{j}\cdot\Delta_{i-j}^{\left[\substack{s-j\\ s-j+(m+j)}\right]\times k}\\&=\sum_{j=1}^{s-1}2^{j}\cdot\Omega_{i-j}(s-j,m+j,k)+\sum_{j=0}^{s-2}2^{j}\cdot\Delta_{i-j}^{\left[\substack{s-j\\ s-j+(m+j)}\right]\times k}.\end{aligned}\tag{9.6}$$

By (9.6) we get after some simplifications

$$\Omega_{i}(s,m,k)=2^{s-1}\cdot\Omega_{i-(s-1)}(1,m+s-1,k)+\sum_{j=0}^{s-2}2^{j}\cdot\Delta_{i-j}^{\left[\substack{s-j\\ s-j+(m+j)}\right]\times k}.\tag{9.7}$$

By the definition of $\Omega_i(s,m,k)$ we have $\Omega_{i-(s-1)}(1,m+s-1,k)=\Gamma_{i-(s-1)}^{\left[\substack{1\\ 1+m+(s-1)}\right]\times k}-4\cdot\Gamma_{i-s}^{\left[\substack{1\\ 1+m+(s-2)}\right]\times k}$.
And (9.1) follows in the second case. □

Lemma 9.2. *We have*

$$\begin{cases}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(k+1)}-\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}=4\cdot\left[\Gamma_{i-1}^{\left[\substack{s\\ s+(m-1)}\right]\times(k+1)}-\Gamma_{i-1}^{\left[\substack{s\\ s+(m-1)}\right]\times k}\right] & \text{if } 2\leq i\leq s-1,\quad k\geq i+1,\\ \Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(k+1)}-\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}=4\cdot\left[\Gamma_{i-1}^{\left[\substack{s\\ s+(m-1)}\right]\times(k+1)}-\Gamma_{i-1}^{\left[\substack{s\\ s+(m-1)}\right]\times k}\right] & \\ +2^{s-1}\left[\Gamma_{i-(s-1)}^{\left[\substack{1\\ 1+(s+m-1)}\right]\times(k+1)}-\Gamma_{i-(s-1)}^{\left[\substack{1\\ 1+(s+m-1)}\right]\times k}\right] & \\ -2^{s+1}\left[\Gamma_{i-s}^{\left[\substack{1\\ 1+(s+m-2)}\right]\times(k+1)}-\Gamma_{i-s}^{\left[\substack{1\\ 1+(s+m-2)}\right]\times k}\right] & \text{if } s\leq i\leq 2s+m,\quad k\geq i+1.\end{cases}\tag{9.8}$$

Proof. By (8.12) we obtain

$$\sum_{j=0}^{i-2}2^{j}\Delta_{i-j}^{\left[\substack{s-j\\ s-j+(m+j)}\right]\times(k+1)}=\sum_{j=0}^{i-2}2^{j}\Delta_{i-j}^{\left[\substack{s-j\\ s-j+(m+j)}\right]\times k}\ for\ 2\leq i\leq s-1,\quad k\geq i+1$$

and equally

$$\sum_{j=0}^{s-2}2^{j}\Delta_{i-j}^{\left[\substack{s-j\\ s-j+(m+j)}\right]\times(k+1)}=\sum_{j=0}^{s-2}2^{j}\Delta_{i-j}^{\left[\substack{s-j\\ s-j+(m+j)}\right]\times k}\ for\ 2\leq s\leq i\leq 2s+m,\quad k\geq i+1.$$

Lemma 9.2 now follows from Theorem 9.1 using the above equations.

□

10. COMPUTATION OF THE REMAINDER R(j,s,m,k) IN THE RECURRENT FORMULA FOR THE DIFFERENCE $\Gamma_{s+j}^{\left[\frac{s}{s+m}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\frac{s}{s+m}\right]\times k}$ $\quad for\ 0 \leq j \leq s+m,\ k > s+j$

From a formula of $\Gamma_{i}^{\left[\frac{1}{1+m}\right]\times k}$ obtained in [4] we compute in this section the remainder R(j,s,m,k) in the recurrent formula for the difference $\Gamma_{i}^{\left[\frac{s}{s+m}\right]\times(k+1)} - \Gamma_{i}^{\left[\frac{s}{s+m}\right]\times k}$.
By definition we have for $s \leq i \leq \inf(2s+m, k)$

$$\Omega_{i-(s-1)}(1, m+s-1, k) = \Gamma_{i-(s-1)}^{\left[\frac{1}{1+m+(s-1)}\right]\times k} - 4 \cdot \Gamma_{i-s}^{\left[\frac{1}{1+m+(s-2)}\right]\times k}.$$

Set i - s = j, then

$$\Omega_{j+1}(1, m+s-1, k) = \Gamma_{j+1}^{\left[\frac{1}{1+m+(s-1)}\right]\times k} - 4 \cdot \Gamma_{j}^{\left[\frac{1}{1+m+(s-2)}\right]\times k}.$$

Lemma 10.1. *We have by Theorem 3.8 [see [4]] with* $m \to s+m-1, m \to s+m-2$

The case s+m -1 = 0, k ≥ 2

$$\Gamma_{j}^{\left[\frac{1}{1+(s+m-1)}\right]\times k} = \Gamma_{j}^{\left[\frac{1}{1}\right]\times k} = \begin{cases} 1 & if\ j = 0, \\ 3 \cdot (2^k - 1) & if\ j = 1, \\ 2^{2k} - 3 \cdot 2^k + 2 & if\ j = 2. \end{cases}$$

The case s+m -1 = 1, k ≥ 3

$$\Gamma_{j}^{\left[\frac{1}{1+(s+m-1)}\right]\times k} = \Gamma_{j}^{\left[\frac{1}{1+1}\right]\times k} = \begin{cases} 1 & if\ j = 0, \\ 2^k + 5 & if\ j = 1, \\ 11 \cdot (2^k - 1) & if\ j = 2, \\ 2^{2k+1} - 3 \cdot 2^{k+2} + 2^4 & if\ j = 3. \end{cases}$$

The case $3 \leq k \leq s+m$

$$\Gamma_{j+1}^{\left[\frac{1}{1+(s+m-1)}\right]\times k} = \begin{cases} 2^k + 5 & if\ j = 0, \\ 3 \cdot 2^{k+2j-2} + 21 \cdot 2^{3j-2} & if\ 1 \leq j \leq k-2, \\ 2^{2k+s+m-1} - 5 \cdot 2^{3k-5} & if\ j = k-1. \end{cases}$$

The case $4 \leq s+m \leq k$

$$\Gamma_{j+1}^{\left[\frac{1}{1+(s+m-1)}\right]\times k} = \begin{cases} 2^k + 5 & if\ j = 0, \\ 3 \cdot 2^{k+2j-2} + 21 \cdot 2^{3j-2} & if\ 1 \leq j \leq s+m-2, \\ 11[2^{k+2s+2m-4} - 2^{3s+3m-5}] & if\ j = s+m-1, \\ 2^{2k+s+m-1} - 3 \cdot 2^{k+2s+2m-2} + 2^{3s+3m-2} & if\ j = s+m. \end{cases}$$

The case $3 \leq k \leq s+m$

$$\Gamma_{j}^{\left[\frac{1}{1+(s+m-2)}\right]\times k} = \begin{cases} 1 & if\ j = 0, \\ 2^k + 5 & if\ j = 1, \\ 3 \cdot 2^{k+2j-4} + 21 \cdot 2^{3j-5} & if\ 2 \leq j \leq k-1, \\ 2^{2k+s+m-2} - 5 \cdot 2^{3k-5} & if\ j = k. \end{cases}$$

The case $4 \leq s+m \leq k$

$$\Gamma_{j}^{\left[\begin{smallmatrix}1\\1+(s+m-2)\end{smallmatrix}\right]\times k} = \begin{cases} 1 & \text{if } j=0,\\ 2^{k}+5 & \text{if } j=1,\\ 3\cdot 2^{k+2j-4}+21\cdot 2^{3j-5} & \text{if } 2\leq j\leq s+m-2,\\ 11[2^{k+2s+2m-6}-2^{3s-3m-8}] & \text{if } j=s+m-1,\\ 2^{2k+s+m-2}-3\cdot 2^{k+2s+2m-4}+2^{3s+3m-5} & \text{if } j=s+m. \end{cases}$$

Lemma 10.2. *For $0 \leq j \leq s+m, k \geq s+j+1$*

$$2^{s-1}\left[\Gamma_{j+1}^{\left[\begin{smallmatrix}1\\1+(s+m-1)\end{smallmatrix}\right]\times (k+1)} - \Gamma_{j+1}^{\left[\begin{smallmatrix}1\\1+(s+m-1)\end{smallmatrix}\right]\times k}\right] - 2^{s+1}\left[\Gamma_{j}^{\left[\begin{smallmatrix}1\\1+(s+m-2)\end{smallmatrix}\right]\times (k+1)} - \Gamma_{j}^{\left[\begin{smallmatrix}1\\1+(s+m-2)\end{smallmatrix}\right]\times k}\right]$$

is equal to

$$\begin{cases} 2^{k+s-1} & \text{if } j=0, \quad k\geq s+1,\\ -2^{k+s-1} & \text{if } j=1, \quad k\geq s+2,\\ 0 & \text{if } 2\leq j\leq s+m-1, \quad k\geq s+j+1,\\ -3\cdot 2^{2k+2s+m-2} & \text{if } j=s+m, \quad k\geq 2s+m+1. \end{cases}$$

Proof. From Lemma 10.1 we deduce

In the case $j=0,\ k\geq s+1$

$$2^{s-1}\left[\Gamma_{1}^{\left[\begin{smallmatrix}1\\1+(s+m-1)\end{smallmatrix}\right]\times (k+1)} - \Gamma_{1}^{\left[\begin{smallmatrix}1\\1+(s+m-1)\end{smallmatrix}\right]\times k}\right] - 2^{s+1}\left[\Gamma_{0}^{\left[\begin{smallmatrix}1\\1+(s+m-2)\end{smallmatrix}\right]\times (k+1)} - \Gamma_{0}^{\left[\begin{smallmatrix}1\\1+(s+m-2)\end{smallmatrix}\right]\times k}\right]$$
$$= 2^{s-1}\left[(2^{k+1}+5)-(2^{k}+5)\right] - 2^{s+1}\left[1-1\right] = 2^{k+s-1}.$$

In the case $j=1,\ k\geq s+2$

$$2^{s-1}\left[\Gamma_{2}^{\left[\begin{smallmatrix}1\\1+(s+m-1)\end{smallmatrix}\right]\times (k+1)} - \Gamma_{2}^{\left[\begin{smallmatrix}1\\1+(s+m-1)\end{smallmatrix}\right]\times k}\right] - 2^{s+1}\left[\Gamma_{1}^{\left[\begin{smallmatrix}1\\1+(s+m-2)\end{smallmatrix}\right]\times (k+1)} - \Gamma_{1}^{\left[\begin{smallmatrix}1\\1+(s+m-2)\end{smallmatrix}\right]\times k}\right]$$
$$= 2^{s-1}\left[(3\cdot 2^{k+1}+42)-(3\cdot 2^{k}+42)\right] - 2^{s+1}\left[(2^{k+1}+5)-(2^{k}+5)\right]$$
$$= 3\cdot 2^{k+s-1} - 2^{k+s+1} = -2^{k+s-1}.$$

In the case $2\leq j\leq m-1,\ s+j+1\leq k\leq s+m$

$$2^{s-1}\left[\Gamma_{j+1}^{\left[\begin{smallmatrix}1\\1+(s+m-1)\end{smallmatrix}\right]\times (k+1)} - \Gamma_{j+1}^{\left[\begin{smallmatrix}1\\1+(s+m-1)\end{smallmatrix}\right]\times k}\right] - 2^{s+1}\left[\Gamma_{j}^{\left[\begin{smallmatrix}1\\1+(s+m-2)\end{smallmatrix}\right]\times (k+1)} - \Gamma_{j}^{\left[\begin{smallmatrix}1\\1+(s+m-2)\end{smallmatrix}\right]\times k}\right]$$
$$= 2^{s-1}\left[(3\cdot 2^{k+2j-1}+21\cdot 2^{3j-2})-(3\cdot 2^{k+2j-2}+21\cdot 2^{3j-2})\right]$$
$$- 2^{s+1}\left[(3\cdot 2^{k+2j-3}+21\cdot 2^{3j-5})-(3\cdot 2^{k+2j-4}+21\cdot 2^{3j-5})\right]$$
$$= 3\cdot 2^{k+2j+s-3} - 3\cdot 2^{k+2j+s-3} = 0$$

In the case $2 \leq j \leq m-1 \leq s+m-2,\ k \geq s+m$

$$\begin{aligned}
&2^{s-1}\left[\Gamma_{j+1}^{\left[\substack{1 \\ 1+(s+m-1)}\right]\times(k+1)} - \Gamma_{j+1}^{\left[\substack{1 \\ 1+(s+m-1)}\right]\times k}\right] - 2^{s+1}\left[\Gamma_{j}^{\left[\substack{1 \\ 1+(s+m-2)}\right]\times(k+1)} - \Gamma_{j}^{\left[\substack{1 \\ 1+(s+m-2)}\right]\times k}\right] \\
&= 2^{s-1}\left[(3\cdot 2^{k+2j-1} + 21\cdot 2^{3j-2}) - (3\cdot 2^{k+2j-2} + 21\cdot 2^{3j-2})\right] \\
&- 2^{s+1}\left[(3\cdot 2^{k+2j-3} + 21\cdot 2^{3j-5}) - (3\cdot 2^{k+2j-4} + 21\cdot 2^{3j-5})\right] \\
&= 3\cdot 2^{k+2j+s-3} - 3\cdot 2^{k+2j+s-3} = 0.
\end{aligned}$$

In the case $2 \leq m \leq j \leq s+m-2,\ k \geq s+j+1 \geq s+m+1$

$$\begin{aligned}
&2^{s-1}\left[\Gamma_{j+1}^{\left[\substack{1 \\ 1+(s+m-1)}\right]\times(k+1)} - \Gamma_{j+1}^{\left[\substack{1 \\ 1+(s+m-1)}\right]\times k}\right] - 2^{s+1}\left[\Gamma_{j}^{\left[\substack{1 \\ 1+(s+m-2)}\right]\times(k+1)} - \Gamma_{j}^{\left[\substack{1 \\ 1+(s+m-2)}\right]\times k}\right] \\
&= 2^{s-1}\left[(3\cdot 2^{k+2j-1} + 21\cdot 2^{3j-2}) - (3\cdot 2^{k+2j-2} + 21\cdot 2^{3j-2})\right] \\
&- 2^{s+1}\left[(3\cdot 2^{k+2j-3} + 21\cdot 2^{3j-5}) - (3\cdot 2^{k+2j-4} + 21\cdot 2^{3j-5})\right] \\
&= 3\cdot 2^{k+2j+s-3} - 3\cdot 2^{k+2j+s-3} = 0.
\end{aligned}$$

In the case $j = s+m-1,\ k \geq s+j+1 = 2s+m$

$$\begin{aligned}
&2^{s-1}\left[\Gamma_{s+m}^{\left[\substack{1 \\ 1+(s+m-1)}\right]\times(k+1)} - \Gamma_{s+m}^{\left[\substack{1 \\ 1+(s+m-1)}\right]\times k}\right] - 2^{s+1}\left[\Gamma_{s+m-1}^{\left[\substack{1 \\ 1+(s+m-2)}\right]\times(k+1)} - \Gamma_{s+m-1}^{\left[\substack{1 \\ 1+(s+m-2)}\right]\times k}\right] \\
&= 2^{s-1}\left[11(2^{k+2s+2m-3} - 2^{3s+3m-5}) - 11(2^{k+2s+2m-4} - 2^{3s+3m-5})\right] \\
&- 2^{s+1}\left[11(2^{k+2s+2m-5} - 2^{3s+3m-8}) - 11(2^{k+2s+2m-6} - 2^{3s+3m-8})\right] \\
&= 11\cdot 2^{k+3s+2m-5} - 11\cdot 2^{k+3s+2m-5} = 0.
\end{aligned}$$

In the case $j = s+m,\ k \geq s+j+1 = 2s+m+1$

$$\begin{aligned}
&2^{s-1}\left[\Gamma_{s+m+1}^{\left[\substack{1 \\ 1+(s+m-1)}\right]\times(k+1)} - \Gamma_{s+m+1}^{\left[\substack{1 \\ 1+(s+m-1)}\right]\times k}\right] - 2^{s+1}\left[\Gamma_{s+m}^{\left[\substack{1 \\ 1+(s+m-2)}\right]\times(k+1)} - \Gamma_{s+m}^{\left[\substack{1 \\ 1+(s+m-2)}\right]\times k}\right] \\
&= 2^{s-1}\left[(2^{2k+s+m+1} - 3\cdot 2^{k+2s+2m-1} + 2^{3s+3m-2}) - (2^{2k+s+m-1} - 3\cdot 2^{k+2s+2m-2} + 2^{3s+3m-2})\right] \\
&- 2^{s+1}\left[(2^{2k+s+m} - 3\cdot 2^{k+2s+2m-3} + 2^{3s+3m-5}) - (2^{2k+s+m-2} - 3\cdot 2^{k+2s+2m-4} + 2^{3s+3m-5})\right] \\
&= 2^{s-1}\left[3\cdot 2^{2k+s+m-1} - 3\cdot 2^{k+2s+2m-2}\right] - 2^{s+1}\left[3\cdot 2^{2k+s+m-2} - 3\cdot 2^{k+2s+2m-4}\right] \\
&= -3\cdot 2^{2k+2s+m-2}.
\end{aligned}$$

In all the others cases the proofs are similar. ☐

Lemma 10.3. *We have for $s \geq 2, m \geq 0$ the following recurrent formula for the difference $\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times(k+1)} - \Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times k}$ where $0 \leq i \leq 2s+m, k > i$.*

(10.1)

$$\begin{cases}\Gamma_{j}^{\left[\substack{s\\ s+m}\right]\times(k+1)}-\Gamma_{j}^{\left[\substack{s\\ s+m}\right]\times k}=4\cdot\left[\Gamma_{j-1}^{\left[\substack{s\\ s+(m-1)}\right]\times(k+1)}-\Gamma_{j-1}^{\left[\substack{s\\ s+(m-1)}\right]\times k}\right] & \text{if } 1\leq j\leq s-1,\quad k\geq j+1,\\ \Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times(k+1)}-\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k}=4\cdot\left[\Gamma_{s+j-1}^{\left[\substack{s\\ s+(m-1)}\right]\times(k+1)}-\Gamma_{s+j-1}^{\left[\substack{s\\ s+(m-1)}\right]\times k}\right] & \\ +\begin{cases}2^{k+s-1} & \text{if } j=0,\quad k\geq s+1,\\ -2^{k+s-1} & \text{if } j=1,\quad k\geq s+2,\\ 0 & \text{if } 2\leq j\leq s+m-1,\quad k\geq s+j+1,\\ -3\cdot 2^{2k+2s+m-2} & \text{if } j=s+m,\quad k\geq 2s+m+1.\end{cases} & \end{cases}$$

Proof. Lemma 10.3 follows from Lemma 9.2 and lemma 10.2.

□

11. COMPUTATION OF $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}$ *for* $i\leq s-1,\ i\leq k-1,\ m\geq 0$

In this section we apply successively the recurrent formula (10.1) for the difference $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(k+1)}-\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}$ to compute $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}$ for $i\leq s-1,\ i\leq k-1,\ m\geq 0$.

Lemma 11.1. *Consider the matrix*

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-1} & \alpha_k\\ \alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_k & \alpha_{k+1}\\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+k-3} & \alpha_{s+k-2}\\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+k-2} & \alpha_{s+k-1}\\ \beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-1} & \beta_k\\ \beta_2 & \beta_3 & \beta_4 & \ldots & \beta_k & \beta_{k+1}\\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\ \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{k+m-1} & \beta_{k+m}\\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\ \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+k-3} & \beta_{s+m+k-2}\\ \beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \end{pmatrix}.$$

Let $g_{k,s,m}(t,\eta)$ *be the quadratic exponential sum in* $\mathbb{P}\times\mathbb{P}$ *defined by*

$$(t,\eta)\in\mathbb{P}\times\mathbb{P}\longmapsto\sum_{degY\leq k-1}\sum_{degZ\leq s-1}E(tYZ)\sum_{degU\leq s+m-1}E(\eta YU)\in\mathbb{Z}.$$

We have

$$\sum_{i=0}^{\inf(2s+m,k)}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}=2^{2k+2s+m-2}\tag{11.1}$$

and

$$\sum_{i=0}^{\inf(2s+m,k)} \Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}\cdot 2^{-i} = 2^{k+2s+m-2}+2^{2k-2}-2^{k-2}. \tag{11.2}$$

Proof. The proof of (11.1) is obvious.

By Lemma 4.6 we obtain, observing that $g_{k,s,m}(t,\eta)$ is constant on cosets of $\mathbb{P}_{k+s-1}\times \mathbb{P}_{k+s+m-1}$

$$\begin{aligned}
\int_{\mathbb{P}\times\mathbb{P}} g_{k,s,m}(t,\eta)dtd\eta &= \sum_{(t,\eta)\in \mathbb{P}/\mathbb{P}_{k+s-1}\times\mathbb{P}/\mathbb{P}_{k+s+m-1}} 2^{2s+m+k-r(D^{\left[\substack{s\\ s+m}\right]\times k}(t,\eta))}\int_{\mathbb{P}_{k+s-1}}dt\int_{\mathbb{P}_{k+s+m-1}}d\eta \\
&= \sum_{i=0}^{\inf(2s+m,k)} \Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}\cdot 2^{2s+m+k-i}\int_{\mathbb{P}_{k+s-1}}dt\int_{\mathbb{P}_{k+s+m-1}}d\eta \\
&= 2^{-k+2}\cdot \sum_{i=0}^{\inf(2s+m,k)} \Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times k}\cdot 2^{-i}.
\end{aligned}$$

On the other hand

$$\begin{aligned}
\int_{\mathbb{P}\times\mathbb{P}} g_{k,s,m}(t,\eta)dtd\eta &= Card\,\{(Y,Z,U), degY\leq k-1, degZ\leq s-1, degU\leq s+m-1 \mid Y\cdot Z = Y\cdot U = 0\} \\
&= 2^{2s+m}+2^{k}-1.
\end{aligned}$$

The above equations imply (11.2). □

Lemma 11.2. *We have for all* $j\geq 1$

(11.3)

$$(H_j) \qquad \Gamma_{j}^{\left[\substack{s\\ s+m}\right]\times k} = \Gamma_{j}^{\left[\substack{j\\ j}\right]\times (j+1)} \overset{def}{=} \gamma_j \quad for \quad s\geq j+1,\ k\geq j+1,\ m\geq 0.$$

Proof. The proof is by strong induction, that is:
If

- (H_1) is true, and
- for all $j\geq 1$, $\quad (H_1)\wedge(H_2)\wedge\ldots\wedge(H_j)\quad$ *implies* (H_{j+1})

then (H_j) is true for all $j\geq 1$.

$\underline{(H_1)\, is\ true}$

Indeed from (10.1) with j = 1

$$\Gamma_{1}^{\left[\substack{s\\ s+m}\right]\times (k+1)} - \Gamma_{1}^{\left[\substack{s\\ s+m}\right]\times k} = 4\cdot\left[\Gamma_{0}^{\left[\substack{s\\ s+(m-1)}\right]\times (k+1)} - \Gamma_{0}^{\left[\substack{s\\ s+(m-1)}\right]\times k}\right] \text{ for all } s\geq 2,\ k\geq 2,\ m\geq 0$$

which implies that (H_j) holds for j = 1, that is

$$\Gamma_{1}^{\left[\substack{s\\ s+m}\right]\times k} = \Gamma_{1}^{\left[\substack{s\\ s+m}\right]\times 2} = \Gamma_{1}^{\left[\substack{1\\ 1}\right]\times 2} = \gamma_1 = 9 \quad \forall\ s\geq 2,\ \forall\ k\geq 2,\ m\geq 0. \tag{11.4}$$

$\underline{(H_1)\Rightarrow(H_2)}$

From (10.1) with j = 2

$\Gamma_2^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_2^{\left[\substack{s\\ s+m}\right]\times k} = 4\cdot\left[\Gamma_1^{\left[\substack{s\\ s+(m-1)}\right]\times(k+1)} - \Gamma_1^{\left[\substack{s\\ s+(m-1)}\right]\times k}\right]$ for all $s \geq 3,\ k \geq 3,\ m \geq 0$.

Hence by (11.4) we have

$$\Gamma_2^{\left[\substack{s\\ s+m}\right]\times k} = \Gamma_2^{\left[\substack{s\\ s+m}\right]\times 3} \quad \text{for all } s \geq 3,\ k \geq 3,\ m \geq 0. \tag{11.5}$$

Consider the matrix

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 \\ \alpha_2 & \alpha_3 & \alpha_4 \\ \vdots & \vdots & \vdots \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} \\ \beta_1 & \beta_2 & \beta_3 \\ \beta_2 & \beta_3 & \beta_4 \\ \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \beta_{m+3} \\ \vdots & \vdots & \vdots \\ \beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} \end{pmatrix}.$$

We have respectively by (11.1), (11.2) with k = 3

$$\sum_{i=0}^{3}\Gamma_i^{\left[\substack{s\\ s+m}\right]\times 3} = 2^{2s+m+4}, \tag{11.6}$$

$$\sum_{i=0}^{3}\Gamma_i^{\left[\substack{s\\ s+m}\right]\times 3}\cdot 2^{-i} = 2^{2s+m+1} + 2^4 - 2. \tag{11.7}$$

From (11.6) and (11.7) using (11.4) we deduce

$$\Gamma_2^{\left[\substack{s\\ s+m}\right]\times 3} = \Gamma_2^{\left[\substack{2\\ 2}\right]\times 3} = 78 \quad \text{and} \quad \Gamma_3^{\left[\substack{s\\ s+m}\right]\times 3} = 2^{2s+m+4} - 88$$

which implies, using (11.5), that (H_j) holds for j = 2, that is

$$\Gamma_2^{\left[\substack{s\\ s+m}\right]\times k} = \Gamma_2^{\left[\substack{2\\ 2}\right]\times 3} = 78 \quad \forall\, s \geq 3,\ \forall\, k \geq 3,\ \forall\, m \geq 0. \tag{11.8}$$

$(H_1) \wedge (H_2)$ *implies* (H_3)

From (10.1) with j = 3

$\Gamma_3^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_3^{\left[\substack{s\\ s+m}\right]\times k} = 4\cdot\left[\Gamma_2^{\left[\substack{s\\ s+(m-1)}\right]\times(k+1)} - \Gamma_2^{\left[\substack{s\\ s+(m-1)}\right]\times k}\right]$ for all $s \geq 4,\ k \geq 4,\ m \geq 0$.

Hence by (11.8) we have

$$\Gamma_3^{\left[\substack{s\\s+m}\right]\times k} = \Gamma_3^{\left[\substack{s\\s+m}\right]\times 4} \quad \text{for all } s \geq 4,\ k \geq 4,\ m \geq 0. \tag{11.9}$$

Consider the matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\
\alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 \\
\vdots & \vdots & \vdots & \vdots \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \alpha_{s+3} \\
\beta_1 & \beta_2 & \beta_3 & \beta_4 \\
\beta_2 & \beta_3 & \beta_4 & \beta_5 \\
\vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \beta_{m+4} \\
\vdots & \vdots & \vdots & \vdots \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \beta_{s+m+3}
\end{pmatrix}.$$

We have respectively by (11.1), (11.2) with k = 4

$$\sum_{i=0}^{4} \Gamma_i^{\left[\substack{s\\s+m}\right]\times 4} = 2^{2s+m+6}, \tag{11.10}$$

$$\sum_{i=0}^{4} \Gamma_i^{\left[\substack{s\\s+m}\right]\times 4} \cdot 2^{-i} = 2^{2s+m+2} + 2^6 - 2^2. \tag{11.11}$$

From (11.10) and (11.11) using (11.5) and (11.9) we deduce

$$\Gamma_3^{\left[\substack{s\\s+m}\right]\times 4} = \Gamma_3^{\left[\substack{3\\3}\right]\times 4} = 648 \quad \text{and} \quad \Gamma_4^{\left[\substack{s\\s+m}\right]\times 4} = 2^{2s+m+6} - 736$$

which implies, using (11.9), that (H_j) holds for j = 3, that is

$$\Gamma_3^{\left[\substack{s\\s+m}\right]\times k} = \Gamma_3^{\left[\substack{3\\3}\right]\times 4} = 648 = \gamma_3 \quad \text{for all } s \geq 4,\ k \geq 4,\ m \geq 0. \tag{11.12}$$

$(H_1) \wedge (H_2) \wedge \ldots \wedge (H_j)$ implies (H_{j+1})

From (10.1) with j → j+1

$$\Gamma_{j+1}^{\left[\substack{s\\s+m}\right]\times (k+1)} - \Gamma_{j+1}^{\left[\substack{s\\s+m}\right]\times k} = 4 \cdot \left[\Gamma_j^{\left[\substack{s\\s+(m-1)}\right]\times (k+1)} - \Gamma_j^{\left[\substack{s\\s+(m-1)}\right]\times k} \right] \quad \text{for all } s \geq j+2,\ k \geq j+2,\ m \geq 0.$$

By (H_j) we obtain

$$\Gamma_j^{\left[\substack{s\\s+(m-1)}\right]\times (k+1)} - \Gamma_j^{\left[\substack{s\\s+(m-1)}\right]\times k} = \gamma_j - \gamma_j = 0.$$

Indeed if m = 0

$$\Gamma_{j}^{\left[\substack{s\\ s+(m-1)}\right]\times k}=\Gamma_{j}^{\left[\substack{s-1\\ s-1+1}\right]\times k}=\Gamma_{j}^{\left[\substack{j\\ j}\right]\times (j+1)}=\gamma_j$$

since

$$s-1\leq j+1,\quad k\geq j+2\geq j+1,\quad m\rightarrow 1\geq 0.$$

If $m\geq 1$

$$\Gamma_{j}^{\left[\substack{s\\ s+(m-1)}\right]\times k}=\Gamma_{j}^{\left[\substack{j\\ j}\right]\times (j+1)}=\gamma_j$$

since

$$s\geq j+2\geq j+1,\quad k\geq j+2\geq j+1,\quad m\rightarrow (m-1)\geq 0.$$

Thus, we obtain

$$\Gamma_{j+1}^{\left[\substack{s\\ s+m}\right]\times k}=\Gamma_{j+1}^{\left[\substack{s\\ s+m}\right]\times (j+2)}\quad \text{for } s\geq j+2,\ k\geq j+2,\ m\geq 0. \tag{11.13}$$

Consider the matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{j+1} & \alpha_{j+2}\\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{j+2} & \alpha_{j+3}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{s+j-1} & \alpha_{s+j}\\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{s+j} & \alpha_{s+j+1}\\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{j+1} & \beta_{j+2}\\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{j+2} & \beta_{j+3}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \ldots & \beta_{j+m+1} & \beta_{j+m+2}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \ldots & \beta_{s+m+j-1} & \beta_{s+m+j}\\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{s+m+j} & \beta_{s+m+j+1}
\end{pmatrix}.$$

We have respectively by (11.1), (11.2) with k = j+2

$$\sum_{i=0}^{j+2}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times (j+2)}=2^{2j+2s+m+2} \tag{11.14}$$

and

$$\sum_{i=0}^{j+2}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times (j+2)}\cdot 2^{-i}=2^{j+2s+m}+2^{2j+2}-2^{j}. \tag{11.15}$$

Setting s = j+1, m = 0 in (11.14), (11.15) we have

$$\sum_{i=0}^{j+2}\Gamma_{i}^{\left[\substack{j+1 \\ j+1}\right]\times(j+2)} = 2^{4j+4} \tag{11.16}$$

and

$$\sum_{i=0}^{j+2}\Gamma_{i}^{\left[\substack{j+1 \\ j+1}\right]\times(j+2)}\cdot 2^{-i} = 2^{3j+2}+2^{2j+2}-2^{j}. \tag{11.17}$$

From $(H_1)\wedge(H_2)\wedge\ldots\wedge(H_j)$, it follows that

$$\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times(j+2)} = \Gamma_{i}^{\left[\substack{j+1 \\ j+1}\right]\times(j+2)} = \Gamma_{i}^{\left[\substack{i \\ i}\right]\times(i+1)} = \gamma_i \quad \text{for} \quad 0\leq i\leq j. \tag{11.18}$$

We deduce from (11.14), (11.15)

$$\Gamma_{j+1}^{\left[\substack{s \\ s+m}\right]\times(j+2)} = 2^{3j+4}-2^{2j+2}+\sum_{i=0}^{j}\gamma_i(1-2^{j+2-i}) \tag{11.19}$$

and from (11.16), (11.17)

$$\Gamma_{j+1}^{\left[\substack{j+1 \\ j+1}\right]\times(j+2)} = 2^{3j+4}-2^{2j+2}+\sum_{i=0}^{j}\gamma_i(1-2^{j+2-i}). \tag{11.20}$$

By (11.13), (11.19) and (11.20) it follows that

$$\Gamma_{j+1}^{\left[\substack{s \\ s+m}\right]\times k} = \Gamma_{j+1}^{\left[\substack{j+1 \\ j+1}\right]\times(j+2)} = \gamma_{j+1} \quad \text{for all } s\geq j+2,\ k\geq j+2,\ m\geq 0.$$

□

Lemma 11.3. *We have*

$$\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times k} = \Gamma_{i}^{\left[\substack{i \\ i}\right]\times(i+1)} = \begin{cases} 1 & \text{if } i=0, \\ 21\cdot 2^{3i-4}-3\cdot 2^{2i-3} & \text{if } 1\leq i\leq s-1,\ k\geq i+1,\ m\geq 0. \end{cases} \tag{11.21}$$

Proof. We have respectively by (11.1), (11.2) with k = i+1, s = i and m = 0, using (11.3)

$$\sum_{j=0}^{i+1}\Gamma_{j}^{\left[\substack{i \\ i}\right]\times(i+1)} = \sum_{j=0}^{i}\gamma_j+\Gamma_{i+1}^{\left[\substack{i \\ i}\right]\times(i+1)} = 2^{4i} \tag{11.22}$$

and

$$\sum_{j=0}^{i+1}\Gamma_{j}^{\left[\begin{smallmatrix} i \\ i \end{smallmatrix}\right]\times(i+1)}\cdot 2^{-j} = \sum_{j=0}^{i}\gamma_j\cdot 2^{-j} + \Gamma_{i+1}^{\left[\begin{smallmatrix} i \\ i \end{smallmatrix}\right]\times(i+1)}\cdot 2^{-(i+1)} = 2^{3i-1}+2^{2i}-2^{i-1}. \tag{11.23}$$

From (11.22) and (11.23) we get

$$\begin{aligned} &2^{i+1}\cdot\left(\sum_{j=0}^{i}\gamma_j\cdot 2^{-j} + \Gamma_{i+1}^{\left[\begin{smallmatrix} i \\ i \end{smallmatrix}\right]\times(i+1)}\cdot 2^{-(i+1)}\right) - \left(\sum_{j=0}^{i}\gamma_j + \Gamma_{i+1}^{\left[\begin{smallmatrix} i \\ i \end{smallmatrix}\right]\times(i+1)}\right) = 2^{3i+1}-2^{2i} \\ &\Leftrightarrow \sum_{j=0}^{i}\gamma_j\cdot\left(2^{i+1-j}-1\right) = 2^{3i+1}-2^{2i}. \end{aligned} \tag{11.24}$$

Hence by (11.24) we deduce

$$\begin{aligned} &\sum_{j=0}^{i}\gamma_j\cdot\left(2^{i+1-j}-1\right) - 2\cdot\left(\sum_{j=0}^{i-1}\gamma_j\cdot\left(2^{i-j}-1\right)\right) = \left(2^{3i+1}-2^{2i}\right) - 2\cdot\left(2^{3i-2}-2^{2i-2}\right) \\ &\Leftrightarrow \sum_{j=0}^{i}\gamma_j = 3\cdot 2^{3i-1}-2^{2i-1}. \end{aligned} \tag{11.25}$$

From (11.25) we get for $i\geq 2$

$$\begin{aligned} &\sum_{j=0}^{i}\gamma_j - \sum_{j=0}^{i-1}\gamma_j = \left(3\cdot 2^{3i-1}-2^{2i-1}\right) - \left(3\cdot 2^{3i-4}-2^{2i-3}\right) \\ &\Leftrightarrow \Gamma_{i}^{\left[\begin{smallmatrix} i \\ i \end{smallmatrix}\right]\times(i+1)} = \gamma_i = 21\cdot 2^{3i-4}-3\cdot 2^{2i-3}. \end{aligned}$$

$\square$

Lemma 11.4. *We have*

$$\Gamma_{k}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = 2^{2k+2s+m-2}-3\cdot 2^{3k-4}+2^{2k-3} \quad \text{for } 1\leq k\leq s, \tag{11.26}$$

$$\sum_{j=0}^{s-1}\Gamma_{j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = 3\cdot 2^{3s-4}-2^{2s-3} \quad \text{for } k\geq s, \tag{11.27}$$

$$\sum_{j=0}^{s-1}\Gamma_{j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}\cdot 2^{-j} = 7\cdot 2^{2s-4}-3\cdot 2^{s-3} \quad \text{for } k\geq s. \tag{11.28}$$

Proof. Consider the matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\
\beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1}
\end{pmatrix}.$$

Obviously by (11.25) we have

$$\begin{aligned}
\Gamma_{k}^{\left[\substack{s\\ s+m}\right]\times k} &= 2^{2k+2s+m-2} - \sum_{j=0}^{k-1}\Gamma_{j}^{\left[\substack{s\\ s+m}\right]\times k} \\
&= 2^{2k+2s+m-2} - \sum_{j=0}^{k-1}\gamma_j \\
&= 2^{2k+2s+m-2} - (3\cdot 2^{3(k-1)-1} - 2^{2(k-1)-1}) = 2^{2k+2s+m-2} - 3\cdot 2^{3k-4} + 2^{2k-3}.
\end{aligned}$$

We deduce (11.27) from (11.25) with i = s -1.
From (11.21) we obtain after some calculations

$$\sum_{j=0}^{s-1}\Gamma_{j}^{\left[\substack{s\\ s+m}\right]\times k}\cdot 2^{-j} = 1 + \sum_{j=1}^{s-1}(21\cdot 2^{2j-4} - 3\cdot 2^{j-3}) = 7\cdot 2^{2s-4} - 3\cdot 2^{s-3}.$$

□

12. COMPUTATION OF $\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k}$ *for* $0 \le j \le s+m,\ k \ge s+j+1,\ m \ge 0$

In this section we apply successively the recurrent formula (10.1) to compute explicitly the difference $\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k}$.

Lemma 12.1. *We have*

$$(12.1)\quad \Gamma_{s+j}^{\left[\substack{s\\ s}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\substack{s\\ s}\right]\times k} = \begin{cases} 3\cdot 2^{k+s-1} & \text{if } j=0,\ k>s, \\ 21\cdot 2^{k+s+3j-4} & \text{if } 1\le j\le s-1,\ k>s+j, \\ 3\cdot 2^{2k+2s-2} - 3\cdot 2^{k+4s-4} & \text{if } j=s,\ k>2s. \end{cases}$$

Proof. From (10.1) we have the following formula

(12.2)

$$\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = 4\cdot\left[\Gamma_{s+j-1}^{\left[\begin{smallmatrix} s \\ s+(m-1)) \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+j-1}^{\left[\begin{smallmatrix} s \\ s+(m-1)) \end{smallmatrix}\right]\times k}\right]$$
$$+\begin{cases} 2^{k+s-1} & \text{if } j=0, \quad k\geq s+1, \\ -2^{k+s-1} & \text{if } j=1, \quad k\geq s+2, \\ 0 & \text{if } 2\leq j\leq s+m-1, \quad k\geq s+j+1, \\ -3\cdot 2^{2k+2s+m-2} & \text{if } j=s+m,\ k\geq 2s+m+1. \end{cases}$$

The case j = 0, $k>s$

Applying formula (12.2) we obtain using (11.21)

$$\Gamma_{s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times k} = 4\cdot\left[\Gamma_{s-1}^{\left[\begin{smallmatrix} s-1 \\ s-1+1 \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s-1}^{\left[\begin{smallmatrix} s-1 \\ s-1+1 \end{smallmatrix}\right]\times k}\right] + 2^{k+s-1}, \tag{12.3}$$

(12.4)

$$\Gamma_{s-1}^{\left[\begin{smallmatrix} s-1 \\ s-1+1 \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s-1}^{\left[\begin{smallmatrix} s-1 \\ s-1+1 \end{smallmatrix}\right]\times k} = 4\cdot\left[\Gamma_{s-2}^{\left[\begin{smallmatrix} s-1 \\ s-1 \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s-2}^{\left[\begin{smallmatrix} s-1 \\ s-1 \end{smallmatrix}\right]\times k}\right] + 2^{k+s-2} = 2^{k+s-2}.$$

From (12.3), (12.4) we get

$$\Gamma_{s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times k} = 4\cdot 2^{k+s-2} + 2^{k+s-1} = 3\cdot 2^{k+s-1}. \tag{12.5}$$

The case j = 1, $k>s+1$

We proceed as in the case j = 0 using (12.5) with $s\rightarrow s-1$

$$\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times k} = 4\cdot\left[\Gamma_{s}^{\left[\begin{smallmatrix} s-1 \\ s-1+1 \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s}^{\left[\begin{smallmatrix} s-1 \\ s-1+1 \end{smallmatrix}\right]\times k}\right] - 2^{k+s-1}, \tag{12.6}$$

$$\begin{aligned} \Gamma_{s}^{\left[\begin{smallmatrix} s-1 \\ s-1+1 \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s}^{\left[\begin{smallmatrix} s-1 \\ s-1+1 \end{smallmatrix}\right]\times k} &= 4\cdot\left[\Gamma_{s-1}^{\left[\begin{smallmatrix} s-1 \\ s-1 \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s-1}^{\left[\begin{smallmatrix} s-1 \\ s-1 \end{smallmatrix}\right]\times k}\right] - 2^{k+s-2} \\ &= 4\cdot 3\cdot 2^{k+s-2} - 2^{k+s-2} = 11\cdot 2^{k+s-2}. \end{aligned} \tag{12.7}$$

From (12.6), (12.7) we get

$$\Gamma_{s+1}^{\begin{bmatrix}s\\s\end{bmatrix}\times(k+1)} - \Gamma_{s+1}^{\begin{bmatrix}s\\s\end{bmatrix}\times k} = 4\cdot 11\cdot 2^{k+s-2} - 2^{k+s-1} = 21\cdot 2^{k+s-1}. \tag{12.8}$$

The case j $= 2,\ k > s+2$

Proceeding as before, using (12.8) with $s \to s-1$

$$\Gamma_{s+2}^{\begin{bmatrix}s\\s\end{bmatrix}\times(k+1)} - \Gamma_{s+2}^{\begin{bmatrix}s\\s\end{bmatrix}\times k} = 4\cdot\left[\Gamma_{s+1}^{\begin{bmatrix}s-1\\s-1+1\end{bmatrix}\times(k+1)} - \Gamma_{s+1}^{\begin{bmatrix}s-1\\s-1+1\end{bmatrix}\times k}\right], \tag{12.9}$$

$$\begin{aligned}\Gamma_{s-1+2}^{\begin{bmatrix}s-1\\s-1+1\end{bmatrix}\times(k+1)} - \Gamma_{s-1+2}^{\begin{bmatrix}s-1\\s-1+1\end{bmatrix}\times k} &= 4\cdot\left[\Gamma_{s-1+1}^{\begin{bmatrix}s-1\\s-1\end{bmatrix}\times(k+1)} - \Gamma_{s-1+1}^{\begin{bmatrix}s-1\\s-1\end{bmatrix}\times k}\right]\\ &= 4\cdot 21\cdot 2^{k+s-2} = 21\cdot 2^{k+s}.\end{aligned} \tag{12.10}$$

From (12.9), (12.10) we get

$$\Gamma_{s+2}^{\begin{bmatrix}s\\s\end{bmatrix}\times(k+1)} - \Gamma_{s+2}^{\begin{bmatrix}s\\s\end{bmatrix}\times k} = 4\cdot\cdot 21\cdot 2^{k+s} = 21\cdot 2^{k+s+2}. \tag{12.11}$$

The case $2 \le j \le s-1,\ k > s+j$

By (12.2) we obtain

$$\Gamma_{s+j}^{\begin{bmatrix}s\\s\end{bmatrix}\times(k+1)} - \Gamma_{s+j}^{\begin{bmatrix}s\\s\end{bmatrix}\times k} = 4\cdot\left[\Gamma_{s+j-1}^{\begin{bmatrix}s-1\\(s-1)+1\end{bmatrix}\times(k+1)} - \Gamma_{s-1+j}^{\begin{bmatrix}s-1\\(s-1)+1\end{bmatrix}\times k}\right] \quad \text{if } 2 \le j \le s-1,$$

$$\Gamma_{s-1+j}^{\begin{bmatrix}s-1\\(s-1)+1\end{bmatrix}\times(k+1)} - \Gamma_{s-1+j}^{\begin{bmatrix}s-1\\(s-1)+1\end{bmatrix}\times k} = 4\cdot\left[\Gamma_{s-1+(j-1)}^{\begin{bmatrix}s-1\\s-1\end{bmatrix}\times(k+1)} - \Gamma_{s-1+(j-1)}^{\begin{bmatrix}s-1\\s-1\end{bmatrix}\times k}\right] \quad \text{if } 2 \le j \le s-1+1-1 = s-1.$$

From the above equations we get

$$\Gamma_{s+j}^{\begin{bmatrix}s\\s\end{bmatrix}\times(k+1)} - \Gamma_{s+j}^{\begin{bmatrix}s\\s\end{bmatrix}\times k} = 4^2\cdot\left[\Gamma_{s-1+(j-1)}^{\begin{bmatrix}s-1\\s-1\end{bmatrix}\times(k+1)} - \Gamma_{s-1+(j-1)}^{\begin{bmatrix}s-1\\s-1\end{bmatrix}\times k}\right] \quad \text{if } 2 \le j \le s-1. \tag{12.12}$$

Using successively (12.12) we get

$$
\begin{aligned}
&\Gamma_{s+j}^{\begin{bmatrix} s\\ s\end{bmatrix}\times(k+1)} - \Gamma_{s+j}^{\begin{bmatrix} s\\ s\end{bmatrix}\times k} \\
&= 4^{2}\cdot\left[\Gamma_{s-1+(j-1)}^{\begin{bmatrix} s-1\\ s-1\end{bmatrix}\times(k+1)} - \Gamma_{s-1+(j-1)}^{\begin{bmatrix} s-1\\ s-1\end{bmatrix}\times k}\right] \qquad \text{if } 2\leq j\leq s-1 \\
&= 4^{4}\cdot\left[\Gamma_{s-2+(j-2)}^{\begin{bmatrix} s-2\\ s-2\end{bmatrix}\times(k+1)} - \Gamma_{s-2+(j-2)}^{\begin{bmatrix} s-2\\ s-2\end{bmatrix}\times k}\right] \qquad \text{if } 2\leq j-1\leq s-1-1 \\
&\vdots \\
&= 4^{2(l-1)}\cdot\left[\Gamma_{s-(l-1)+(j-(l-1))}^{\begin{bmatrix} s-(l-1)\\ s-(l-1)\end{bmatrix}\times(k+1)} - \Gamma_{s-(l-1)+(j-(l-1))}^{\begin{bmatrix} s-(l-1)\\ s-(l-1)\end{bmatrix}\times k}\right] \qquad \text{if } 2\leq j-(l-2)\leq s-(l-2)-1 \\
&= 4^{2l}\cdot\left[\Gamma_{s-l+(j-l)}^{\begin{bmatrix} s-l\\ s-l\end{bmatrix}\times(k+1)} - \Gamma_{s-l+(j-l)}^{\begin{bmatrix} s-l\\ s-l\end{bmatrix}\times k}\right] \qquad \text{if } 2\leq j-(l-1)\leq s-(l-1)-1 \\
&\vdots \\
&= 4^{2(j-1)}\cdot\left[\Gamma_{s-(j-1)+(j-(j-1))}^{\begin{bmatrix} s-(j-1)\\ s-(j-1)\end{bmatrix}\times(k+1)} - \Gamma_{s-(j-1)+(j-(j-1))}^{\begin{bmatrix} s-(j-1)\\ s-(j-1)\end{bmatrix}\times k}\right] \qquad \text{if } 2\leq j-((j-1)-1)\leq s-((j-1)-1)-1.
\end{aligned}
$$

From the above equations we get, using (12.8) with $s\to s-j+1$

(12.13)

$$
\begin{aligned}
\Gamma_{s+j}^{\begin{bmatrix} s\\ s\end{bmatrix}\times(k+1)} - \Gamma_{s+j}^{\begin{bmatrix} s\\ s\end{bmatrix}\times k} &= 4^{2(j-1)}\cdot\left[\Gamma_{s-j+2}^{\begin{bmatrix} s-j+1\\ s-j+1\end{bmatrix}\times(k+1)} - \Gamma_{s-j+2}^{\begin{bmatrix} s-j+1\\ s-j+1\end{bmatrix}\times k}\right] \qquad \text{if } 2\leq j\leq s-1, \\
&= 4^{2(j-1)}\cdot 21\cdot 2^{k+s-j+1-1} = 21\cdot 2^{k+s+3j-4}.
\end{aligned}
$$

The case j = s, $k>2s$

By (12.2) we obtain

$$
\Gamma_{2s}^{\begin{bmatrix} s\\ s\end{bmatrix}\times(k+1)} - \Gamma_{2s}^{\begin{bmatrix} s\\ s\end{bmatrix}\times k} = 4\cdot\left[\Gamma_{2s-1}^{\begin{bmatrix} s-1\\ (s-1)+1\end{bmatrix}\times(k+1)} - \Gamma_{2s-1}^{\begin{bmatrix} s-1\\ (s-1)+1\end{bmatrix}\times k}\right] - 3\cdot 2^{2k+2s-2},
$$

$$
\Gamma_{(s-1)+s}^{\begin{bmatrix} s-1\\ (s-1)+1\end{bmatrix}\times(k+1)} - \Gamma_{(s-1)+s}^{\begin{bmatrix} s-1\\ (s-1)+1\end{bmatrix}\times k} = 4\cdot\left[\Gamma_{2(s-1)}^{\begin{bmatrix} s-1\\ s-1\end{bmatrix}\times(k+1)} - \Gamma_{2(s-1)}^{\begin{bmatrix} s-1\\ s-1\end{bmatrix}\times k}\right] - 3\cdot 2^{2k+2(s-1)+1-2}.
$$

From the above equations we get

$$\begin{aligned}
\Gamma_{2s}^{\begin{bmatrix} s \\ s \end{bmatrix} \times (k+1)} - \Gamma_{2s}^{\begin{bmatrix} s \\ s \end{bmatrix} \times k} &= 4^2 \cdot \left[\Gamma_{2(s-1)}^{\begin{bmatrix} s-1 \\ s-1 \end{bmatrix} \times (k+1)} - \Gamma_{2(s-1)}^{\begin{bmatrix} s-1 \\ s-1 \end{bmatrix} \times k} \right] - 9 \cdot 2^{2k+2s-2} \\
\Gamma_{2(s-1)}^{\begin{bmatrix} s-1 \\ s-1 \end{bmatrix} \times (k+1)} - \Gamma_{2(s-1)}^{\begin{bmatrix} s-1 \\ s-1 \end{bmatrix} \times k} &= 4^2 \cdot \left[\Gamma_{2(s-2)}^{\begin{bmatrix} s-2 \\ s-2 \end{bmatrix} \times (k+1)} - \Gamma_{2(s-2)}^{\begin{bmatrix} s-2 \\ s-2 \end{bmatrix} \times k} \right] - 9 \cdot 2^{2k+2(s-1)-2} \\
&= \\
&\vdots \\
\Gamma_{2(s-l)}^{\begin{bmatrix} s-l \\ s-l \end{bmatrix} \times (k+1)} - \Gamma_{2(s-l)}^{\begin{bmatrix} s-l \\ s-l \end{bmatrix} \times k} &= 4^2 \cdot \left[\Gamma_{2(s-l-1)}^{\begin{bmatrix} s-l-1 \\ s-l-1 \end{bmatrix} \times (k+1)} - \Gamma_{2(s-1-1)}^{\begin{bmatrix} s-l-1 \\ s-l-1 \end{bmatrix} \times k} \right] - 9 \cdot 2^{2k+2(s-l)-2} \\
&\vdots \\
\Gamma_{2(s-(s-2))}^{\begin{bmatrix} s-(s-2) \\ s-(s-2) \end{bmatrix} \times (k+1)} - \Gamma_{2(s-(s-2))}^{\begin{bmatrix} s-(s-2) \\ s-(s-2) \end{bmatrix} \times k} &= 4^2 \cdot \left[\Gamma_{2(s-(s-2)-1)}^{\begin{bmatrix} s-(s-2)-1 \\ s-(s-2)-1 \end{bmatrix} \times (k+1)} - \Gamma_{2(s-(s-2)-1)}^{\begin{bmatrix} s-(s-2)-1 \\ s-(s-2)-1 \end{bmatrix} \times k} \right] - 9 \cdot 2^{2k+2(s-(s-2))-2}.
\end{aligned}$$

We deduce from the above equations

$$\begin{aligned}
(12.14) \qquad & \sum_{l=0}^{s-2} 4^{2l} \cdot \left[\Gamma_{2(s-l)}^{\begin{bmatrix} s-l \\ s-l \end{bmatrix} \times (k+1)} - \Gamma_{2(s-l)}^{\begin{bmatrix} s-l \\ s-l \end{bmatrix} \times k} \right] \\
&= \sum_{l=0}^{s-2} \left(4^{2(l+1)} \cdot \left[\Gamma_{2(s-l-1)}^{\begin{bmatrix} s-l-1 \\ s-l-1 \end{bmatrix} \times (k+1)} - \Gamma_{2(s-l-1)}^{\begin{bmatrix} s-l-1 \\ s-l-1 \end{bmatrix} \times k} \right] - 9 \cdot 2^{2k+2(s-l)-2} \cdot 2^{4l} \right) \\
&= \sum_{l=1}^{s-1} 4^{2l} \cdot \left[\Gamma_{2(s-l)}^{\begin{bmatrix} s-l \\ s-l \end{bmatrix} \times (k+1)} - \Gamma_{2(s-l)}^{\begin{bmatrix} s-l \\ s-l \end{bmatrix} \times k} \right] - \sum_{l=1}^{s-1} 9 \cdot 2^{2k+2s+2l-4}.
\end{aligned}$$

We get from (12.14) after some simplications

$$\begin{aligned}
\Gamma_{2s}^{\begin{bmatrix} s \\ s \end{bmatrix} \times (k+1)} - \Gamma_{2s}^{\begin{bmatrix} s \\ s \end{bmatrix} \times k} &= 4^{2(s-1)} \left[\Gamma_{2}^{\begin{bmatrix} 1 \\ 1 \end{bmatrix} \times (k+1)} - \Gamma_{2}^{\begin{bmatrix} 1 \\ 1 \end{bmatrix} \times k} \right] - 9 \cdot 2^{2k+2s-4} \cdot \sum_{l=1}^{s-1} 2^{2l} \\
&= 4^{2(s-1)} \left[(2^{2(k+1)} - 3 \cdot 2^{k+1} + 2) - (2^{2k} - 3 \cdot 2^{k} + 2) \right] - 3 \cdot 2^{2k+2s-2} \cdot (2^{2s-2} - 1) \\
&= 3 \cdot 2^{2k+2s-2} - 3 \cdot 2^{k+4s-4}.
\end{aligned}$$

□

Lemma 12.2. *We have*

(12.15)

$$\Gamma_{s+j}^{\begin{bmatrix} s \\ s+1 \end{bmatrix} \times (k+1)} - \Gamma_{s+j}^{\begin{bmatrix} s \\ s+1 \end{bmatrix} \times k} = \begin{cases} 2^{k+s-1} & \text{if } j = 0,\ k > s, \\ 11 \cdot 2^{k+s-1} & \text{if } j = 1,\ k > s+1, \\ 21 \cdot 2^{k+s+3j-5} & \text{if } 2 \leq j \leq s,\ k > s+j, \\ 3 \cdot 2^{2k+2s-1} - 3 \cdot 2^{k+4s-2} & \text{if } j = s+1,\ k > 2s+1. \end{cases}$$

Proof. We proceed as in the proof of Lemma 12.1.

The case j = 0, $k > s$

Applying formula (12.2) we obtain using (11.21)

$$\Gamma_{s}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}\times(k+1)} - \Gamma_{s}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}\times k} = 4\cdot\left[\Gamma_{s-1}^{\begin{bmatrix} s \\ s \end{bmatrix}\times(k+1)} - \Gamma_{s-1}^{\begin{bmatrix} s \\ s \end{bmatrix}\times k}\right] + 2^{k+s-1} = 2^{k+s-1}. \tag{12.16}$$

The case j = 1, $k > s+1$

From (12.2) we obtain using (12.5)

$$\begin{aligned}\Gamma_{s+1}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}\times(k+1)} - \Gamma_{s+1}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}\times k} &= 4\cdot\left[\Gamma_{s}^{\begin{bmatrix} s \\ s \end{bmatrix}\times(k+1)} - \Gamma_{s}^{\begin{bmatrix} s \\ s \end{bmatrix}\times k}\right] - 2^{k+s-1} \\ &= 4\cdot(3\cdot 2^{k+s-1}) - 2^{k+s-1} = 11\cdot 2^{k+s-1}.\end{aligned} \tag{12.17}$$

The case $2 \leq j \leq s,\ k > s+j$

From (12.2) we obtain using (12.1) with $j \to j-1$ (observing that $1 \leq j-1 \leq s-1$)

$$\begin{aligned}\Gamma_{s+j}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}\times(k+1)} - \Gamma_{s+j}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}\times k} &= 4\cdot\left[\Gamma_{s+j-1}^{\begin{bmatrix} s \\ s \end{bmatrix}\times(k+1)} - \Gamma_{s+j-1}^{\begin{bmatrix} s \\ s \end{bmatrix}\times k}\right] \\ &= 4\cdot(21\cdot 2^{k+s+3(j-1)-4}) = 21\cdot 2^{k+s+3j-5}.\end{aligned} \tag{12.18}$$

The case j = s+1

$$\begin{aligned}\Gamma_{2s+1}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}\times(k+1)} - \Gamma_{2s+1}^{\begin{bmatrix} s \\ s+1 \end{bmatrix}\times k} &= 4\cdot\left[\Gamma_{2s}^{\begin{bmatrix} s \\ s \end{bmatrix}\times(k+1)} - \Gamma_{2s}^{\begin{bmatrix} s \\ s \end{bmatrix}\times k}\right] - 3\cdot 2^{2k+2s-1} \\ &= 4\cdot[3\cdot 2^{2k+2s-2} - 3\cdot 2^{k+4s-4}] - 3\cdot 2^{2k+2s-1} \\ &= 3\cdot 2^{2k+2s-1} - 3\cdot 2^{k+4s-2}.\end{aligned} \tag{12.19}$$

□

Lemma 12.3. *We have in the case $m \geq 2$*

$$\Gamma_{s+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times(k+1)} - \Gamma_{s+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times k} = \begin{cases} 2^{k+s-1} & \text{if } j = 0,\ k > s, \\ 3\cdot 2^{k+s+2j-3} & \text{if } 1 \leq j \leq m-1,\ k > s+j, \\ 11\cdot 2^{k+s+2m-3} & \text{if } j = m,\ k > s+m, \end{cases} \tag{12.20}$$

(12.21)

$$\Gamma_{s+m+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times(k+1)} - \Gamma_{s+m+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times k} = \begin{cases} 21\cdot 2^{k+s+2m+3j-4} & \text{if } 1 \leq j \leq s-1,\ k > s+m+j, \\ 3\cdot 2^{2k+2s+m-2} - 3\cdot 2^{k+4s+2m-4} & \text{if } j = s,\ k > 2s+m. \end{cases}$$

Proof. The case s+j with $0 \leq j \leq m,\ k > s+j$

j = 0

Applying formula (12.2) we obtain using (11.21), observing that $m-1 \geq 1 \geq 0$

(12.22)
$$\Gamma_{s}^{\left[\substack{s \\ s+m}\right]\times(k+1)} - \Gamma_{s}^{\left[\substack{s \\ s+m}\right]\times k} = 4\cdot\left[\Gamma_{s-1}^{\left[\substack{s \\ s+(m-1)}\right]\times(k+1)} - \Gamma_{s-1}^{\left[\substack{s \\ s+(m-1)}\right]\times k}\right] + 2^{k+s-1} = 2^{k+s-1}.$$

j = 1

Applying formula (12.2) we obtain using (11.21), observing that $m-2 \geq 0$

$$\begin{aligned}
(12.23)\qquad & \Gamma_{s+1}^{\left[\substack{s \\ s+m}\right]\times(k+1)} - \Gamma_{s+1}^{\left[\substack{s \\ s+m}\right]\times k} \\
&= 4\cdot\left[\Gamma_{s}^{\left[\substack{s \\ s+(m-1)}\right]\times(k+1)} - \Gamma_{s}^{\left[\substack{s \\ s+(m-1)}\right]\times k}\right] - 2^{k+s-1} \\
&= 4\cdot\left(4\cdot\left[\Gamma_{s-1}^{\left[\substack{s \\ s+(m-2)}\right]\times(k+1)} - \Gamma_{s-1}^{\left[\substack{s \\ s+(m-2)}\right]\times k}\right] + 2^{k+s-1}\right) - 2^{k+s-1} \\
&= 3\cdot 2^{k+s-1}.
\end{aligned}$$

$2 \leq j \leq m-1$

Assume $0 \leq l \leq j-2$, then by (12.2) with $j \rightarrow j-l,\ m \rightarrow m-l$, we have

(12.24)
$$\Gamma_{s+j-l}^{\left[\substack{s \\ s+(m-l)}\right]\times(k+1)} - \Gamma_{s+j-l}^{\left[\substack{s \\ s+(m-l)}\right]\times k} = 4\cdot\left[\Gamma_{s+j-(l+1)}^{\left[\substack{s \\ s+(m-(l+1))}\right]\times(k+1)} - \Gamma_{s+j-(l+1)}^{\left[\substack{s \\ s+(m-(l+1))}\right]\times k}\right] \quad \text{for } l = 0,1,2,\ldots,j-2.$$

From (12.24) we deduce

(12.25)
$$\begin{aligned}
\prod_{l=0}^{j-2}\left[\Gamma_{s+j-l}^{\left[\substack{s \\ s+(m-l)}\right]\times(k+1)} - \Gamma_{s+j-l}^{\left[\substack{s \\ s+(m-l)}\right]\times k}\right] &= \prod_{l=0}^{j-2} 4\cdot\left[\Gamma_{s+j-(l+1)}^{\left[\substack{s \\ s+(m-(l+1))}\right]\times(k+1)} - \Gamma_{s+j-(l+1)}^{\left[\substack{s \\ s+(m-(l+1))}\right]\times k}\right] \\
&= \prod_{l=1}^{j-1} 4\cdot\left[\Gamma_{s+j-l}^{\left[\substack{s \\ s+(m-l)}\right]\times(k+1)} - \Gamma_{s+j-l}^{\left[\substack{s \\ s+(m-l)}\right]\times k}\right].
\end{aligned}$$

We then obtain by (12.25), (12.23), observing that $m-(j-1) \geq 2$

$$\begin{aligned}
\Gamma_{s+j}^{\left[\substack{s \\ s+m}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\substack{s \\ s+m}\right]\times k} &= 4^{j-1}\cdot\left[\Gamma_{s+1}^{\left[\substack{s \\ s+(m-(j-1))}\right]\times(k+1)} - \Gamma_{s+1}^{\left[\substack{s \\ s+(m-(j-1))}\right]\times k}\right] \\
&= 2^{2j-2}\cdot 3\cdot 2^{k+s-1} = 3\cdot 2^{k+s+2j-3}.
\end{aligned}$$

j = m

Assume $2 \leq l \leq m$ $(\leq s+m-1)$ then by (12.2) with $j \rightarrow l,\ m \rightarrow l$, we get

(12.26)

$$\Gamma_{s+l}^{\left[\substack{s\\ s+l}\right]\times(k+1)} - \Gamma_{s+l}^{\left[\substack{s\\ s+l}\right]\times k} = 4\cdot\left[\Gamma_{s+(l-1)}^{\left[\substack{s\\ s+(l-1)}\right]\times(k+1)} - \Gamma_{s+(l-1)}^{\left[\substack{s\\ s+(l-1)}\right]\times k}\right] \quad for\ l = 2,3,\ldots,m.$$

From (12.26) we deduce

$$\prod_{l=2}^{m}\left[\Gamma_{s+l}^{\left[\substack{s\\ s+l}\right]\times(k+1)} - \Gamma_{s+l}^{\left[\substack{s\\ s+l}\right]\times k}\right] = \prod_{l=2}^{m}4\cdot\left[\Gamma_{s+(l-1)}^{\left[\substack{s\\ s+(l-1)}\right]\times(k+1)} - \Gamma_{s+(l-1)}^{\left[\substack{s\\ s+(l-1)}\right]\times k}\right] \tag{12.27}$$

$$= \prod_{l=1}^{m-1}4\cdot\left[\Gamma_{s+l}^{\left[\substack{s\\ s+l}\right]\times(k+1)} - \Gamma_{s+l}^{\left[\substack{s\\ s+l}\right]\times k}\right].$$

We then obtain by (12.27) and (12.15) with j = 1

$$\Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times k} = 4^{m-1}\cdot\left[\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times(k+1)} - \Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times k}\right]$$

$$= 2^{2m-2}\cdot 11\cdot 2^{k+s-1} = 11\cdot 2^{k+s+2m-3}.$$

The case s+m+j with $1 \leq j \leq s-1$

Assume $0 \leq l \leq m-1$, then by (12.2) with $j \rightarrow (m-l)+j,\ m \rightarrow m-l$, (observing that $2 \leq m-l+j \leq s+(m-l)-1$) we have

(12.28)

$$\Gamma_{s+(m-l)+j}^{\left[\substack{s\\ s+(m-l)}\right]\times(k+1)} - \Gamma_{s+l}^{\left[\substack{s\\ s+(m-l)}\right]\times k} = 4\cdot\left[\Gamma_{s+(m-(l+1))+j}^{\left[\substack{s\\ s+(m-(l+1))}\right]\times(k+1)} - \Gamma_{s+(m-(l+1))+j}^{\left[\substack{s\\ s+(m-(l+1))}\right]\times k}\right] \quad \text{for } l = 0,1,\ldots,m-1.$$

From (12.28) we deduce

(12.29)

$$\prod_{l=0}^{m-1}\left[\Gamma_{s+(m-l)+j}^{\left[\substack{s\\ s+(m-l)}\right]\times(k+1)} - \Gamma_{s+(m-l)+j}^{\left[\substack{s\\ s+(m-l)}\right]\times k}\right] = \prod_{l=0}^{m-1}4\cdot\left[\Gamma_{s+(m-(l+1))+j}^{\left[\substack{s\\ s+(m-(l+1))}\right]\times(k+1)} - \Gamma_{s+(m-(l+1))+j}^{\left[\substack{s\\ s+(m-(l+1))}\right]\times k}\right]$$

$$= \prod_{l=1}^{m}4\cdot\left[\Gamma_{s+(m-l)+j}^{\left[\substack{s\\ s+(m-l)}\right]\times(k+1)} - \Gamma_{s+(m-l)}^{\left[\substack{s\\ s+(m-l)}\right]\times k}\right].$$

We then obtain by (12.29) and (12.1)

$$\begin{aligned}\Gamma_{s+m+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times(k+1)} - \Gamma_{s+m+j}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times k} &= 4^{m}\cdot\left[\Gamma_{s+j}^{\begin{bmatrix} s \\ s \end{bmatrix}\times(k+1)} - \Gamma_{s+j}^{\begin{bmatrix} s \\ s \end{bmatrix}\times k}\right] \\ &= 2^{2m}\cdot 21\cdot 2^{k+s+3j-4} = 21\cdot 2^{k+s+2m+3j-4}.\end{aligned}$$

The case s+m+j with j = s

Assume $0 \leq l \leq m-1$, then by (12.2) with $j \to s+(m-l),\ m \to m-l$, we have

(12.30)

$$\begin{aligned}&\Gamma_{2s+(m-l)}^{\begin{bmatrix} s \\ s+(m-l) \end{bmatrix}\times(k+1)} - \Gamma_{2s+(m-l)}^{\begin{bmatrix} s \\ s+(m-l) \end{bmatrix}\times k} \\ &= 4\cdot\left[\Gamma_{2s+(m-(l+1))}^{\begin{bmatrix} s \\ s+(m-(l+1)) \end{bmatrix}\times(k+1)} - \Gamma_{2s+(m-(l+1))}^{\begin{bmatrix} s \\ s+(m-(l+1)) \end{bmatrix}\times k}\right] - 3\cdot 2^{2k+2s+(m-l)-2} \quad \text{for } l = 0,1,\ldots,m-1.\end{aligned}$$

From (12.30) we deduce

(12.31)

$$\begin{aligned}&\sum_{l=0}^{m-1} 4^{l}\cdot\left[\Gamma_{2s+(m-l)}^{\begin{bmatrix} s \\ s+(m-l) \end{bmatrix}\times(k+1)} - \Gamma_{2s+(m-l)}^{\begin{bmatrix} s \\ s+(m-l) \end{bmatrix}\times k}\right] \\ &= \sum_{l=0}^{m-1} 4^{l}\cdot\left(4\cdot\left[\Gamma_{2s+(m-(l+1))}^{\begin{bmatrix} s \\ s+(m-(l+1)) \end{bmatrix}\times(k+1)} - \Gamma_{2s+(m-(l+1))}^{\begin{bmatrix} s \\ s+(m-(l+1)) \end{bmatrix}\times k}\right] - 3\cdot 2^{2k+2s+(m-l)-2}\right) \\ &= \sum_{l=1}^{m} 4^{l}\cdot\left[\Gamma_{2s+(m-l)}^{\begin{bmatrix} s \\ s+(m-l) \end{bmatrix}\times(k+1)} - \Gamma_{2s+(m-l)}^{\begin{bmatrix} s \\ s+(m-l) \end{bmatrix}\times k}\right] - \sum_{l=1}^{m} 3\cdot 2^{2k+2s+m+l-3}.\end{aligned}$$

We then obtain by (12.31) and (12.1) after some simplifications

$$\begin{aligned}\Gamma_{2s+m}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times(k+1)} - \Gamma_{2s+m}^{\begin{bmatrix} s \\ s+m \end{bmatrix}\times k} &= 4^{m}\cdot\left[\Gamma_{2s}^{\begin{bmatrix} s \\ s \end{bmatrix}\times(k+1)} - \Gamma_{2s}^{\begin{bmatrix} s \\ s \end{bmatrix}\times k}\right] \\ &= 2^{2m}\cdot(3\cdot 2^{2k+2s-2} - 3\cdot 2^{k+4s-4}) - (3\cdot 2^{2k+2s+2m-2} - 3\cdot 2^{2k+2s+m-2}) \\ &= 3\cdot 2^{2k+2s+m-2} - 3\cdot 2^{k+4s+2m-4}.\end{aligned}$$

□

Lemma 12.4. *We have in the case* $m \in \{0,1\}$

(12.32)

$$\Gamma_{s+j}^{\begin{bmatrix} s \\ s \end{bmatrix}\times(k+1)} - \Gamma_{s+j}^{\begin{bmatrix} s \\ s \end{bmatrix}\times k} = \begin{cases} 3\cdot 2^{k+s-1} & \text{if } j=0,\ k>s, \\ 21\cdot 2^{k+s+3j-4} & \text{if } 1\leq j\leq s-1,\ k>s+j, \\ 3\cdot 2^{2k+2s-2} - 3\cdot 2^{k+4s-4} & \text{if } j=s,\ k>2s, \end{cases}$$

(12.33)

$$\Gamma_{s+j}^{\left[\substack{s\\ s+1}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\substack{s\\ s+1}\right]\times k} = \begin{cases} 2^{k+s-1} & \text{if } j=0,\ k>s,\\ 11\cdot 2^{k+s-1} & \text{if } j=1,\ k>s+1,\\ 21\cdot 2^{k+s+3j-5} & \text{if } 2\leq j\leq s,\ k>s+j,\\ 3\cdot 2^{2k+2s-1}-3\cdot 2^{k+4s-2} & \text{if } j=s+1,\ k>2s+1. \end{cases}$$

In the case $m\geq 2$, we get

$$\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k} = \begin{cases} 2^{k+s-1} & \text{if } j=0,\ k>s,\\ 3\cdot 2^{k+s+2j-3} & \text{if } 1\leq j\leq m-1,\ k>s+j,\\ 11\cdot 2^{k+s+2m-3} & \text{if } j=m,\ k>s+m, \end{cases} \tag{12.34}$$

(12.35)

$$\Gamma_{s+m+j}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{s+m+j}^{\left[\substack{s\\ s+m}\right]\times k} = \begin{cases} 21\cdot 2^{k+s+2m+3j-4} & \text{if } 1\leq j\leq s-1,\ k>s+m+j,\\ 3\cdot 2^{2k+2s+m-2}-3\cdot 2^{k+4s+2m-4} & \text{if } j=s,\ k>2s+m. \end{cases}$$

Proof. The assertions follow from Lemmas 12.1, 12.2 and 12.3.

□

13. COMPUTATION OF $\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times k}$ *for* $j\in\{0,1\},\ k\geq s+j,\ m\geq 0$

From the equations (11.1) and (11.2) with k = s+1 we deduce $\Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times(s+1)}$ and $\Gamma_{s+1}^{\left[\substack{s\\ s+m}\right]\times(s+1)}$ since by (11.21) the terms $\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(s+1)}$ are known for $i\leq s-1$. Then from the recurrent formula (12.34) with j = 0 we compute $\Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times k}$ for $k\geq s+2$. The other results are obtained in a similar way.

Lemma 13.1. *We have in the case m = 0*

$$\Gamma_{s}^{\left[\substack{s\\ s}\right]\times s} = 2^{4s-2}-3\cdot 2^{3s-4}+2^{2s-3}, \tag{13.1}$$

$$\Gamma_{s}^{\left[\substack{s\\ s}\right]\times k} = 3\cdot 2^{k+s-1}+21\cdot 2^{3s-4}-27\cdot 2^{2s-3} \quad \text{if } k>s, \tag{13.2}$$

$$\Gamma_{s+1}^{\left[\substack{s\\ s}\right]\times(s+1)} = 2^{4s}-3\cdot 2^{3s-1}+2^{2s-1}, \tag{13.3}$$

$$\Gamma_{s+1}^{\left[\substack{s\\ s}\right]\times k} = 21\cdot[2^{k+s-1}+2^{3s-1}-5\cdot 2^{2s-1}] \quad \text{if } k>s+1. \tag{13.4}$$

Proof. We get (13.1) from (11.26) with k = s, m = 0.

To prove (13.2) and (13.3) we proceed as follows :
Consider the matrix

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_s & \alpha_{s+1} \\ \alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+1} & \alpha_{s+2} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s-2} & \alpha_{2s-1} \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s-1} & \alpha_{2s} \\ \beta_1 & \beta_2 & \beta_3 & \ldots & \beta_s & \beta_{s+1} \\ \beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+1} & \beta_{s+2} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s-2} & \beta_{2s-1} \\ \beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s-1} & \beta_{2s} \end{pmatrix}.$$

We have respectively by (11.1), (11.2), (11.27) and (11.28) with m = 0, k = s+1

$$\sum_{i=0}^{s+1}\Gamma_{i}^{\left[\substack{s \\ s}\right]\times(s+1)} = 2^{4s}, \tag{13.5}$$

$$\sum_{i=0}^{s+1}\Gamma_{i}^{\left[\substack{s \\ s}\right]\times(s+1)}\cdot 2^{-i} = 2^{3s-1}+2^{2s}-2^{s-1}, \tag{13.6}$$

$$\sum_{i=0}^{s-1}\Gamma_{i}^{\left[\substack{s \\ s}\right]\times(s+1)} = 3\cdot 2^{3s-4}-2^{2s-3} \tag{13.7}$$

and

$$\sum_{i=0}^{s-1}\Gamma_{i}^{\left[\substack{s \\ s}\right]\times(s+1)}\cdot 2^{-i} = 7\cdot 2^{2s-4}-3\cdot 2^{s-3}. \tag{13.8}$$

From (13.5), (13.6), (13.7) and (13.8) we deduce after some calculations

$$\Gamma_{s}^{\left[\substack{s \\ s}\right]\times(s+1)} = 21\cdot 2^{3s-4}-3\cdot 2^{2s-3}, \tag{13.9}$$

$$\Gamma_{s+1}^{\left[\substack{s \\ s}\right]\times(s+1)} = 2^{4s}-3\cdot 2^{3s-1}+2^{2s-1}. \tag{13.10}$$

By (12.32) with j = 0 we get

$$\Gamma_{s}^{\left[\substack{s \\ s}\right]\times(k+1)}-\Gamma_{s}^{\left[\substack{s \\ s}\right]\times k} = 3\cdot 2^{k+s-1} \quad \text{if} \quad k>s. \tag{13.11}$$

From (13.11), (13.8) we deduce

$$\sum_{j=s+1}^{k}\left(\Gamma_{s}^{\left[\substack{s \\ s}\right]\times(j+1)}-\Gamma_{s}^{\left[\substack{s \\ s}\right]\times j}\right) = \sum_{j=s+1}^{k}3\cdot 2^{j+s-1}$$

$$\Leftrightarrow \sum_{j=s+2}^{k+1} \Gamma_{s}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times j} - \sum_{j=s+1}^{k} \Gamma_{s}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times j} = 3\cdot 2^{k+s} - 3\cdot 2^{2s}$$

$$\Leftrightarrow \Gamma_{s}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times (k+1)} - \Gamma_{s}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times (s+1)} = 3\cdot 2^{k+s} - 3\cdot 2^{2s}$$

$$\Leftrightarrow \Gamma_{s}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times (k+1)} = 3\cdot 2^{k+s} - 3\cdot 2^{2s} + 21\cdot 2^{3s-4} - 3\cdot 2^{2s-3}$$

$$\Leftrightarrow \Gamma_{s}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times (k+1)} = 3\cdot 2^{k+s} + 21\cdot 2^{3s-4} - 27\cdot 2^{2s-3} \quad \text{if } k > s. \tag{13.12}$$

By (13.8) we see that (13.12) holds for $k = s$, then (13.12) holds for $k \geq s$.

To prove (13.4) we proceed as follows :
Consider the matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{s+1} & \alpha_{s+2} \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+2} & \alpha_{s+3} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s-1} & \alpha_{2s} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s} & \alpha_{2s+1} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{s+1} & \beta_{s+2} \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+2} & \beta_{s+3} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s-1} & \beta_{2s} \\
\beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s} & \beta_{2s+1}
\end{pmatrix}.$$

We have respectively by (11.1), (11.2), (11.27) and (11.28) with m = 0, k = s+2,

$$\sum_{i=0}^{s+2} \Gamma_{i}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times (s+2)} = 2^{4s+2}, \tag{13.13}$$

$$\sum_{i=0}^{s+2} \Gamma_{i}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times (s+2)} \cdot 2^{-i} = 2^{3s} + 2^{2s+2} - 2^{s}, \tag{13.14}$$

$$\sum_{i=0}^{s-1} \Gamma_{i}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times (s+1)} = 3\cdot 2^{3s-4} - 2^{2s-3} \tag{13.15}$$

and

$$\sum_{i=0}^{s-1} \Gamma_{i}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times (s+2)} \cdot 2^{-i} = 7\cdot 2^{2s-4} - 3\cdot 2^{s-3}. \tag{13.16}$$

From (13.13), (13.14), (13.15), (13.16) and (13.2) with k = s+2 we obtain

$$\Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(s+2)} + \Gamma_{s+2}^{\left[\substack{s\\s}\right]\times(s+2)}$$

(13.17)

$$= 2^{4s+2} - (3\cdot 2^{3s-4} - 2^{2s-3}) - (3\cdot 2^{s+2+s-1} + 21\cdot 2^{3s-4} - 27\cdot 2^{2s-3}),$$

$$\Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(s+2)}\cdot 2^{-(s+1)} + \Gamma_{s+2}^{\left[\substack{s\\s}\right]\times(s+2)}\cdot 2^{-(s+2)}$$

(13.18)

$$= 2^{3s} + 2^{2s+2} - 2^{s} - (7\cdot 2^{2s-4} - 3\cdot 2^{s-3}) - (3\cdot 2^{s+2+s-1} + 21\cdot 2^{3s-4} - 27\cdot 2^{2s-3})\cdot 2^{-s}.$$

Hence by (13.17), (13.18) we deduce after some calculations

$$\Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(s+2)} = 21\cdot(2^{3s-1} - 2^{2s-1}), \tag{13.19}$$

$$\Gamma_{s+2}^{\left[\substack{s\\s}\right]\times(s+2)} = 2^{4s+2} - 3\cdot 2^{3s+2} + 2^{2s+3}. \tag{13.20}$$

By (12.32), with j = 1, we get

$$\Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(k+1)} - \Gamma_{s+1}^{\left[\substack{s\\s}\right]\times k} = 21\cdot 2^{k+s-1} \quad \text{if} \quad k > s+1. \tag{13.21}$$

From (13.19), (13.21) we deduce

$$\begin{aligned}
&\sum_{j=s+2}^{k}\left(\Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(j+1)} - \Gamma_{s+1}^{\left[\substack{s\\s}\right]\times j}\right) = \sum_{j=s+2}^{k} 21\cdot 2^{j+s-1}\\
&\Leftrightarrow \sum_{j=s+3}^{k+1}\Gamma_{s+1}^{\left[\substack{s\\s}\right]\times j} - \sum_{j=s+2}^{k}\Gamma_{s+1}^{\left[\substack{s\\s}\right]\times j} = 21\cdot 2^{k+s} - 21\cdot 2^{2s+1}\\
&\Leftrightarrow \Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(k+1)} - \Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(s+2)} = 21\cdot 2^{k+s} - 21\cdot 2^{2s+1}\\
&\Leftrightarrow \Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(k+1)} = 21\cdot 2^{k+s} - 21\cdot 2^{2s+1} + 21\cdot(2^{3s-1} - 2^{2s-1})\\
&\Leftrightarrow \Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(k+1)} = 21[2^{k+s} + 2^{3s-1} - 5\cdot 2^{2s-1}] \quad \text{if } k > s+1.
\end{aligned} \tag{13.22}$$

By (13.19) we see that (13.22) holds for k = s+1, then (13.22) holds for $k \geq s+1$.

□

Lemma 13.2. *In the case m = 1, we have*

$$\Gamma_{s}^{\left[\substack{s\\s+1}\right]\times s} = 2^{4s-1} - 3\cdot 2^{3s-4} + 2^{2s-3}, \tag{13.23}$$

$$\Gamma_{s}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix}\right]\times k} = 2^{k+s-1} + 21\cdot 2^{3s-4} - 11\cdot 2^{2s-3} \quad \text{if } k > s, \tag{13.24}$$

$$\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix}\right]\times (s+1)} = 2^{4s+1} - 3\cdot 2^{3s-1} + 2^{2s-1}, \tag{13.25}$$

$$\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix}\right]\times k} = 11\cdot 2^{k+s-1} + 21\cdot 2^{3s-1} - 53\cdot 2^{2s-1} \quad \text{if } k > s+1. \tag{13.26}$$

Proof. We proceed as in the proof of Lemma 13.1.

Proof of (13.23)

Follows from (11.26) with m = 1, k = s.

Proof of (13.24),(13.25)

Consider the matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_s & \alpha_{s+1} \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+1} & \alpha_{s+2} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s-2} & \alpha_{2s-1} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s-1} & \alpha_{2s} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_s & \beta_{s+1} \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+1} & \beta_{s+2} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s-2} & \beta_{2s-1} \\
\beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s-1} & \beta_{2s} \\
\beta_{s+1} & \beta_{s+2} & \beta_{s+3} & \ldots & \beta_{2s} & \beta_{2s+1}
\end{pmatrix}.$$

We have respectively by (11.1), (11.2), (11.27) and (11.28) with m = 1, k = s+1

$$\sum_{i=0}^{s+1} \Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix}\right]\times (s+1)} = 2^{4s+1}, \tag{13.27}$$

$$\sum_{i=0}^{s+1} \Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix}\right]\times (s+1)} \cdot 2^{-i} = 2^{3s} + 2^{2s} - 2^{s-1}, \tag{13.28}$$

$$\sum_{i=0}^{s-1} \Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix}\right]\times (s+1)} = 3\cdot 2^{3s-4} - 2^{2s-3} \tag{13.29}$$

and

$$\sum_{i=0}^{s-1} \Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+1 \end{smallmatrix}\right]\times (s+1)} \cdot 2^{-i} = 7\cdot 2^{2s-4} - 3\cdot 2^{s-3}. \tag{13.30}$$

From (13.27), (13.28), (13.29) and (13.30) we deduce after some calculations

$$\Gamma_{s}^{\left[\substack{s \\ s+1}\right] \times (s+1)} = 21 \cdot 2^{3s-4} - 3 \cdot 2^{2s-3}, \tag{13.31}$$

$$\Gamma_{s+1}^{\left[\substack{s \\ s+1}\right] \times (s+1)} = 2^{4s+1} - 3 \cdot 2^{3s-1} + 2^{2s-1}. \tag{13.32}$$

By (12.33) with j = 0 we get

$$\Gamma_{s}^{\left[\substack{s \\ s+1}\right] \times (k+1)} - \Gamma_{s}^{\left[\substack{s \\ s+1}\right] \times k} = 2^{k+s-1} \quad \text{if} \quad k > s. \tag{13.33}$$

From (13.33), (13.31) we deduce

$$\begin{aligned}
&\sum_{j=s+1}^{k} \left(\Gamma_{s}^{\left[\substack{s \\ s+1}\right] \times (j+1)} - \Gamma_{s}^{\left[\substack{s \\ s+1}\right] \times j} \right) = \sum_{j=s+1}^{k} 2^{j+s-1} \\
&\Leftrightarrow \sum_{j=s+2}^{k+1} \Gamma_{s}^{\left[\substack{s \\ s+1}\right] \times j} - \sum_{j=s+1}^{k} \Gamma_{s}^{\left[\substack{s \\ s+1}\right] \times j} = 2^{k+s} - 2^{2s} \\
&\Leftrightarrow \Gamma_{s}^{\left[\substack{s \\ s+1}\right] \times (k+1)} - \Gamma_{s}^{\left[\substack{s \\ s+1}\right] \times (s+1)} = 2^{k+s} - 2^{2s} \\
&\Leftrightarrow \Gamma_{s}^{\left[\substack{s \\ s+1}\right] \times (k+1)} = 2^{k+s} - 2^{2s} + 21 \cdot 2^{3s-4} - 3 \cdot 2^{2s-3}
\end{aligned}$$

$$\Leftrightarrow \Gamma_{s}^{\left[\substack{s \\ s+1}\right] \times (k+1)} = 2^{k+s} + 21 \cdot 2^{3s-4} - 11 \cdot 2^{2s-3} \quad \text{if } k > s. \tag{13.34}$$

By (13.31) we see that (13.34) holds for k = s, then (13.34) holds for $k \geq s$.

Proof of (13.26)

Consider the matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{s+1} & \alpha_{s+2} \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+2} & \alpha_{s+3} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s-1} & \alpha_{2s} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s} & \alpha_{2s+1} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{s+1} & \beta_{s+2} \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+2} & \beta_{s+3} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s-1} & \beta_{2s} \\
\beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s} & \beta_{2s+1} \\
\beta_{s+1} & \beta_{s+2} & \beta_{s+3} & \ldots & \beta_{2s+1} & \beta_{2s+2}
\end{pmatrix}.$$

We have respectively by (11.1), (11.2), (11.27) and (11.28) with m = 1 and k = s+2

$$\sum_{i=0}^{s+2} \Gamma_{i}^{\left[\substack{s \\ s+1}\right] \times (s+2)} = 2^{4s+3}, \tag{13.35}$$

$$\sum_{i=0}^{s+2}\Gamma_{i}^{\left[\substack{s\\ s+1}\right]\times(s+2)}\cdot 2^{-i}=2^{3s+1}+2^{2s+2}-2^{s}, \tag{13.36}$$

$$\sum_{i=0}^{s-1}\Gamma_{i}^{\left[\substack{s\\ s+1}\right]\times(s+2)}=3\cdot 2^{3s-4}-2^{2s-3} \tag{13.37}$$

and

$$\sum_{i=0}^{s-1}\Gamma_{i}^{\left[\substack{s\\ s+1}\right]\times(s+2)}\cdot 2^{-i}=7\cdot 2^{2s-4}-3\cdot 2^{s-3}. \tag{13.38}$$

From (13.35), (13.36), (13.37), (13.38) and (13.24) with k = s+2 we obtain

$$\begin{aligned}&\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times(s+2)}+\Gamma_{s+2}^{\left[\substack{s\\ s+1}\right]\times(s+2)}\\ &=2^{4s+3}-(3\cdot 2^{3s-4}-2^{2s-3})-(2^{s+2+s-1}+21\cdot 2^{3s-4}-11\cdot 2^{2s-3}),\end{aligned} \tag{13.39}$$

$$\begin{aligned}&\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times(s+2)}\cdot 2^{-(s+1)}+\Gamma_{s+2}^{\left[\substack{s\\ s+1}\right]\times(s+2)}\cdot 2^{-(s+2)}\\ &=2^{3s+1}+2^{2s+2}-2^{s}-(7\cdot 2^{2s-4}-3\cdot 2^{s-3})-(2^{s+2+s-1}+21\cdot 2^{3s-4}-11\cdot 2^{2s-3})\cdot 2^{-s}.\end{aligned} \tag{13.40}$$

Hence by (13.39), (13.40) we deduce after some calculations

$$\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times(s+2)}=21\cdot 2^{3s-1}-9\cdot 2^{2s-1}, \tag{13.41}$$

$$\Gamma_{s+2}^{\left[\substack{s\\ s+1}\right]\times(s+2)}=2^{4s+3}-3\cdot 2^{3s+2}+2^{2s+2}. \tag{13.42}$$

By (12.33) with j = 1 we get

$$\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times(k+1)}-\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times k}=11\cdot 2^{k+s-1}\quad\text{if}\quad k>s+1. \tag{13.43}$$

From (13.43), (13.41) we deduce

$$\begin{aligned}&\sum_{j=s+2}^{k}\left(\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times(j+1)}-\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times j}\right)=\sum_{j=s+2}^{k}11\cdot 2^{j+s-1}\\ \Leftrightarrow&\sum_{j=s+3}^{k+1}\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times j}-\sum_{j=s+2}^{k}\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times j}=11\cdot 2^{k+s}-11\cdot 2^{2s+1}\\ \Leftrightarrow&\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times(k+1)}-\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times(s+2)}=11\cdot 2^{k+s}-11\cdot 2^{2s+1}\end{aligned}$$

$$\Leftrightarrow \Gamma_{s+1}^{\left[\substack{s\\s+1}\right]\times(k+1)} = 11\cdot 2^{k+s} - 11\cdot 2^{2s+1} + 21\cdot 2^{3s-1} - 9\cdot 2^{2s-1}$$

$$\Leftrightarrow \Gamma_{s+1}^{\left[\substack{s\\s+1}\right]\times(k+1)} = 11\cdot 2^{k+s} + 21\cdot 2^{3s-1} - 53\cdot 2^{2s-1} \quad \text{if } k > s+1. \tag{13.44}$$

By (13.41) we see that (13.44) holds for k = s+1, then (13.44) holds for $k \geq s+1$.

□

Lemma 13.3. *In the case $m \geq 2$, we have*

$$\Gamma_{s}^{\left[\substack{s\\s+m}\right]\times s} = 2^{4s+m-2} - 3\cdot 2^{3s-4} + 2^{2s-3}, \tag{13.45}$$

$$\Gamma_{s}^{\left[\substack{s\\s+m}\right]\times k} = 2^{k+s-1} + 21\cdot 2^{3s-4} - 11\cdot 2^{2s-3} \quad \textit{if } k > s, \tag{13.46}$$

$$\Gamma_{s+1}^{\left[\substack{s\\s+m}\right]\times(s+1)} = 2^{4s+m} - 3\cdot 2^{3s-1} + 2^{2s-1}, \tag{13.47}$$

$$\Gamma_{s+1}^{\left[\substack{s\\s+m}\right]\times k} = 3\cdot 2^{k+s-1} + 21\cdot [2^{3s-1} - 2^{2s-1}] \quad \textit{if } k > s+1, \tag{13.48}$$

$$\Gamma_{s+2}^{\left[\substack{s\\s+m}\right]\times(s+2)} = 2^{4s+m+2} - 3\cdot 2^{3s+2} + 2^{2s+2}. \tag{13.49}$$

Proof. We proceed as in the proof of Lemma 13.2.

Proof of (13.45)

Follows from (11.26) with $m \geq 2$, k = s.

Proof of (13.46), (13.47)

Consider the matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_s & \alpha_{s+1} \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+1} & \alpha_{s+2} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s-2} & \alpha_{2s-1} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s-1} & \alpha_{2s} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_s & \beta_{s+1} \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+1} & \beta_{s+2} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s-2} & \beta_{2s-1} \\
\beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s-1} & \beta_{2s} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{2s+m-1} & \beta_{2s+m}
\end{pmatrix}.$$

We have respectively by (11.1), (11.2), (11.27) and (11.28) with $m \geq 2$, k = s+1

$$\sum_{i=0}^{s+1}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(s+1)} = 2^{4s+m}, \tag{13.50}$$

$$\sum_{i=0}^{s+1}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(s+1)}\cdot 2^{-i} = 2^{3s+m-1}+2^{2s}-2^{s-1}, \tag{13.51}$$

$$\sum_{i=0}^{s-1}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(s+1)} = 3\cdot 2^{3s-4}-2^{2s-3} \tag{13.52}$$

and

$$\sum_{i=0}^{s-1}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(s+1)}\cdot 2^{-i} = 7\cdot 2^{2s-4}-3\cdot 2^{s-3}. \tag{13.53}$$

From (13.50), (13.51), (13.52) and (13.53) we deduce after some calculations

$$\Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times(s+1)} = 21\cdot 2^{3s-4}-3\cdot 2^{2s-3}, \tag{13.54}$$

$$\Gamma_{s+1}^{\left[\substack{s\\ s+m}\right]\times(s+1)} = 2^{4s+m}-3\cdot 2^{3s-1}+2^{2s-1}. \tag{13.55}$$

By (12.34) with j = 0, $m \geq 2$ we get

$$\Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times k} = 2^{k+s-1} \quad \text{if} \quad k > s. \tag{13.56}$$

From (13.56), (13.54) we deduce

$$\begin{aligned}
&\sum_{j=s+1}^{k}\left(\Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times(j+1)} - \Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times j}\right) = \sum_{j=s+1}^{k} 2^{j+s-1}\\
&\Leftrightarrow \sum_{j=s+2}^{k+1}\Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times j} - \sum_{j=s+1}^{k}\Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times j} = 2^{k+s}-2^{2s}\\
&\Leftrightarrow \Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times(s+1)} = 2^{k+s}-2^{2s}\\
&\Leftrightarrow \Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times(k+1)} = 2^{k+s}-2^{2s}+21\cdot 2^{3s-4}-3\cdot 2^{2s-3}
\end{aligned}$$

$$\Leftrightarrow \Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times(k+1)} = 2^{k+s}+21\cdot 2^{3s-4}-11\cdot 2^{2s-3} \quad \text{if } k > s. \tag{13.57}$$

By (13.54) we see that (13.57) holds for k = s, then (13.57) holds for $k \geq s$.

Proof of (13.48),(13.49)

Consider the matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{s+1} & \alpha_{s+2} \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+2} & \alpha_{s+3} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s-1} & \alpha_{2s} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s} & \alpha_{2s+1} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{s+1} & \beta_{s+2} \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+2} & \beta_{s+3} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s-1} & \beta_{2s} \\
\beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s} & \beta_{2s+1} \\
\beta_{s+1} & \beta_{s+2} & \beta_{s+3} & \ldots & \beta_{2s+1} & \beta_{2s+2} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{2s+m} & \beta_{2s+m+1}
\end{pmatrix}.$$

We have respectively by (11.1), (11.2), (11.27) and (11.28) with $m \geq 2$, k = s+2

$$\sum_{i=0}^{s+2} \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+2)} = 2^{4s+m+2}, \tag{13.58}$$

$$\sum_{i=0}^{s+2} \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+2)} \cdot 2^{-i} = 2^{3s+m} + 2^{2s+2} - 2^s, \tag{13.59}$$

$$\sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+2)} = 3 \cdot 2^{3s-4} - 2^{2s-3} \tag{13.60}$$

and

$$\sum_{i=0}^{s-1} \Gamma_i^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+2)} \cdot 2^{-i} = 7 \cdot 2^{2s-4} - 3 \cdot 2^{s-3}. \tag{13.61}$$

From (13.58), (13.59), (13.60), (13.61) and (13.46) with k = s+2 we obtain

$$\begin{aligned}
&\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+2)} + \Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+2)} \\
&= 2^{4s+m+2} - (3 \cdot 2^{3s-4} - 2^{2s-3}) - (2^{s+2+s-1} + 21 \cdot 2^{3s-4} - 11 \cdot 2^{2s-3}),
\end{aligned} \tag{13.62}$$

$$\begin{aligned}
&\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+2)} \cdot 2^{-(s+1)} + \Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+2)} \cdot 2^{-(s+2)} \\
&= 2^{3s+m} + 2^{2s+2} - 2^s - (7 \cdot 2^{2s-4} - 3 \cdot 2^{s-3}) - (2^{s+2+s-1} + 21 \cdot 2^{3s-4} - 11 \cdot 2^{2s-3}) \cdot 2^{-s}.
\end{aligned} \tag{13.63}$$

Hence by (13.62),(13.63) we deduce after some calculations

$$\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+2)} = 21\cdot 2^{3s-1} - 9\cdot 2^{2s-1}, \tag{13.64}$$

$$\Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+2)} = 2^{4s+m+2} - 3\cdot 2^{3s+2} + 2^{2s+2}. \tag{13.65}$$

By (12.34) with $j = 1$, $m \geq 2$ we get

$$\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = 3\cdot 2^{k+s-1} \quad \text{if} \quad k > s+1. \tag{13.66}$$

From (13.66), (13.64) we deduce

$$\begin{aligned}
&\sum_{j=s+2}^{k}\left(\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(j+1)} - \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times j}\right) = \sum_{j=s+2}^{k} 3\cdot 2^{j+s-1} \\
&\Leftrightarrow \sum_{j=s+3}^{k+1}\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times j} - \sum_{j=s+2}^{k}\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times j} = 3\cdot 2^{k+s} - 3\cdot 2^{2s+1} \\
&\Leftrightarrow \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+2)} = 3\cdot 2^{k+s} - 3\cdot 2^{2s+1} \\
&\Leftrightarrow \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k+1)} = 3\cdot 2^{k+s} - 3\cdot 2^{2s+1} + 21\cdot 2^{3s-1} - 9\cdot 2^{2s-1} \\
&\Leftrightarrow \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k+1)} = 3\cdot 2^{k+s} + 21\cdot 2^{3s-1} - 21\cdot 2^{2s-1} \quad \text{if } k > s+1.
\end{aligned} \tag{13.67}$$

By (13.64) we see that (13.67) holds for k = s+1, then (13.67) holds for $k \geq s+1$.

□

14. COMPUTATION OF $\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k}$ *for* $j \in \{2,3\}$, $k \geq s+j$, $j \leq m$

In this section we show the following reduction formulas to be needed in the induction proof in the next section

$$\Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = 8\cdot\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+(m-1) \end{smallmatrix}\right]\times(k-1)} \qquad \text{if } k \geq s+2,\ m \geq 2,$$

$$\Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = 8\cdot\Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+(m-2) \end{smallmatrix}\right]\times(k-2)} \qquad \text{if } k \geq s+3,\ m \geq 3.$$

We recall that the right hand sides in the above equations have already been computed in section 13.
In fact we have

$$\Gamma_{s+1}^{\left[\substack{s\\ s+(m-(j-1))}\right]\times(k-(j-1))} = \begin{cases} 2^{4s+1}-3\cdot 2^{3s-1}+2^{2s-1} & \text{if} \quad j=2,\ k=s+2,\ m=2,\\ 11\cdot 2^{k+s-2}+21\cdot 2^{3s-1}-53\cdot 2^{2s-1} & \text{if} \quad j=2,\ k>s+2,\ m=2,\\ 2^{4s+m-1}-3\cdot 2^{3s-1}+2^{2s-1} & \text{if} \quad j=2,\ k=s+2,\ m>2,\\ 3\cdot 2^{k+s-2}+21\cdot[2^{3s-1}-2^{2s-1}] & \text{if} \quad j=2,\ k>s+2,\ m>2,\\ 2^{4s+1}-3\cdot 2^{3s-1}+2^{2s-1} & \text{if} \quad j=3,\ k=s+3,\ m=3,\\ 11\cdot 2^{k+s-3}+21\cdot 2^{3s-1}-53\cdot 2^{2s-1} & \text{if} \quad j=3,\ k>s+3,\ m=3,\\ 2^{4s+m-2}-3\cdot 2^{3s-1}+2^{2s-1} & \text{if} \quad j=3,\ k=s+3,\ m>3,\\ 3\cdot 2^{k+s-3}+21\cdot[2^{3s-1}-2^{2s-1}] & \text{if} \quad j=3,\ k>s+3,\ m>3. \end{cases}$$

Lemma 14.1. *We have*

(14.1)
$$\Gamma_{s+2}^{\left[\substack{s\\ s+m}\right]\times(s+2)} = 8\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+(m-1)}\right]\times(s+1)} = 2^{4s+m+2}-3\cdot 2^{3s+2}+2^{2s+2} \qquad \textit{if}\ \ m\geq 2,$$

(14.2)
$$\Gamma_{s+2}^{\left[\substack{s\\ s+m}\right]\times k} = 8\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+(m-1)}\right]\times(k-1)} = 3\cdot 2^{k+s+1}+21\cdot 2^{3s+2}-21\cdot 2^{2s+2} \quad \textit{if}\ \ k>s+2, \quad m\geq 3,$$

(14.3)
$$\Gamma_{s+2}^{\left[\substack{s\\ s+2}\right]\times k} = 8\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times(k-1)} = 11\cdot 2^{k+s+1}+21\cdot 2^{3s+2}-53\cdot 2^{2s+2} \qquad \textit{if}\ \ k>s+2..$$

Proof. Proof of (14.1)

By (13.65),(13.47) and (13.25) we get

$$\Gamma_{s+2}^{\left[\substack{s\\ s+m}\right]\times(s+2)} = 2^{4s+m+2}-3\cdot 2^{3s+2}+2^{2s+2} \quad \text{if } m\geq 2,$$

$$\begin{aligned} 8\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+(m-1)}\right]\times(s+1)} &= 8\cdot[2^{4s+(m-1)}-3\cdot 2^{3s-1}+2^{2s-1}]\\ &= 2^{4s+m+2}-3\cdot 2^{3s+2}+2^{2s+2} \quad \text{if } m\geq 2. \end{aligned}$$

Proof of (14.2)
Consider the matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{s+2} & \alpha_{s+3} \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+3} & \alpha_{s+4} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s} & \alpha_{2s+1} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s+1} & \alpha_{2s+2} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{s+2} & \beta_{s+3} \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+3} & \beta_{s+4} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s} & \beta_{2s+1} \\
\beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s+1} & \beta_{2s+2} \\
\beta_{s+1} & \beta_{s+2} & \beta_{s+3} & \ldots & \beta_{2s+2} & \beta_{2s+3} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{2s+m+1} & \beta_{2s+m+2}
\end{pmatrix}.$$

We have respectively by (11.1), (11.2), (11.27) and (11.28) with $m \geq 2$, k = s+3

$$\sum_{i=0}^{s+3} \Gamma_i^{\left[\substack{s \\ s+m}\right]\times(s+3)} = 2^{4s+m+4}, \tag{14.4}$$

$$\sum_{i=0}^{s+3} \Gamma_i^{\left[\substack{s \\ s+m}\right]\times(s+3)} \cdot 2^{-i} = 2^{3s+m+1} + 2^{2s+4} - 2^{s+1}, \tag{14.5}$$

$$\sum_{i=0}^{s-1} \Gamma_i^{\left[\substack{s \\ s+m}\right]\times(s+3)} = 3 \cdot 2^{3s-4} - 2^{2s-3} \tag{14.6}$$

and

$$\sum_{i=0}^{s-1} \Gamma_i^{\left[\substack{s \\ s+m}\right]\times(s+3)} \cdot 2^{-i} = 7 \cdot 2^{2s-4} - 3 \cdot 2^{s-3}. \tag{14.7}$$

From (14.4), (14.5), (14.6), (14.7), (13.46) and (13.48) with k = s+3 we obtain

(14.8)
$$\Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times(s+3)} + \Gamma_{s+3}^{\left[\substack{s \\ s+m}\right]\times(s+3)}$$
$$= 2^{4s+m+4} - (3 \cdot 2^{3s-4} - 2^{2s-3}) - (2^{s+3+s-1} + 21 \cdot 2^{3s-4} - 11 \cdot 2^{2s-3})$$
$$- (3 \cdot 2^{s+3+s-1} + 21 \cdot 2^{3s-1} - 21 \cdot 2^{2s-1}),$$

(14.9)
$$\Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times(s+3)} \cdot 2^{-(s+2)} + \Gamma_{s+3}^{\left[\substack{s \\ s+m}\right]\times(s+3)} \cdot 2^{-(s+3)}$$
$$= 2^{3s+m+1} + 2^{2s+4} - 2^{s+1} - (7 \cdot 2^{2s-4} - 3 \cdot 2^{s-3})$$
$$- (2^{s+3+s-1} + 21 \cdot 2^{3s-4} - 11 \cdot 2^{2s-3}) \cdot 2^{-(s+1)} - (3 \cdot 2^{s+3+s-1} + 21 \cdot 2^{3s-1} - 21 \cdot 2^{2s-1}) \cdot 2^{-(s+2)}.$$

Hence by (14.8), (14.9) we deduce after some calculations

$$\Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times(s+3)} = 21\cdot 2^{3s+2} - 9\cdot 2^{2s+2}, \tag{14.10}$$

$$\Gamma_{s+3}^{\left[\substack{s \\ s+m}\right]\times(s+3)} = 2^{4s+m+4} - 3\cdot 2^{3s+5} + 2^{2s+5}. \tag{14.11}$$

By (12.34) with j = 2, $m \geq 3$ we get

$$\Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times(k+1)} - \Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times k} = 3\cdot 2^{k+s+1} \quad \text{if} \quad k > s+2. \tag{14.12}$$

From (14.12), (14.10) we deduce

$$\sum_{j=s+3}^{k}\left(\Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times(j+1)} - \Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times j}\right) = \sum_{j=s+3}^{k} 3\cdot 2^{j+s+1}$$

$$\Leftrightarrow \sum_{j=s+4}^{k+1}\Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times j} - \sum_{j=s+3}^{k}\Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times j} = 3\cdot 2^{k+s+2} - 3\cdot 2^{2s+4}$$

$$\Leftrightarrow \Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times(k+1)} - \Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times(s+3)} = 3\cdot 2^{k+s+2} - 3\cdot 2^{2s+4}$$

$$\Leftrightarrow \Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times(k+1)} = 3\cdot 2^{k+s+2} - 3\cdot 2^{2s+4} + 21\cdot 2^{3s+2} - 9\cdot 2^{2s+2}$$

$$\Leftrightarrow \Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times(k+1)} = 3\cdot 2^{k+s+2} + 21\cdot 2^{3s+2} - 21\cdot 2^{2s+2} \quad \text{if } k > s+3. \tag{14.13}$$

By (14.10) we see that (14.13) holds for k = s+3, then (14.13) holds for $k \geq s+3$.

From (13.48) with $m \to m-1$, $k \to k-1$, $m \geq 3$, $k-1 > s+1$ and (14.13) we get

$$8\cdot\Gamma_{s+1}^{\left[\substack{s \\ s+(m-1)}\right]\times(k-1)} = 8\cdot[3\cdot 2^{k+s-2} + 21\cdot(2^{3s-1} - 2^{2s-1})], \tag{14.14}$$

$$\Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times k} = 3\cdot 2^{k+s+1} + 21\cdot 2^{3s+2} - 21\cdot 2^{2s+2}. \tag{14.15}$$

By (14.14), (14.15) we obtain for $m \geq 3$, $k \geq s+3$

$$\Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times k} = 8\cdot\Gamma_{s+1}^{\left[\substack{s \\ s+(m-1)}\right]\times(k-1)}.$$

Proof of (14.3)

By (12.34) with $j = 2$, $m = 2$ we get

$$\Gamma_{s+2}^{\left[\substack{s \\ s+2}\right]\times(k+1)} - \Gamma_{s+2}^{\left[\substack{s \\ s+2}\right]\times k} = 11\cdot 2^{k+s+1} \quad \text{if} \quad k > s+2. \tag{14.16}$$

From (14.16), (14.10) we deduce

$$\begin{aligned}
&\sum_{j=s+3}^{k}\left(\Gamma_{s+2}^{\left[\substack{s\\ s+2}\right]\times(j+1)}-\Gamma_{s+2}^{\left[\substack{s\\ s+2}\right]\times j}\right)=\sum_{j=s+3}^{k}11\cdot 2^{j+s+1}\\
&\Leftrightarrow \sum_{j=s+4}^{k+1}\Gamma_{s+2}^{\left[\substack{s\\ s+2}\right]\times j}-\sum_{j=s+3}^{k}\Gamma_{s+2}^{\left[\substack{s\\ s+2}\right]\times j}=11\cdot 2^{k+s+2}-11\cdot 2^{2s+4}\\
&\Leftrightarrow \Gamma_{s+2}^{\left[\substack{s\\ s+2}\right]\times(k+1)}-\Gamma_{s+2}^{\left[\substack{s\\ s+2}\right]\times(s+3)}=11\cdot 2^{k+s+2}-11\cdot 2^{2s+4}\\
&\Leftrightarrow \Gamma_{s+2}^{\left[\substack{s\\ s+2}\right]\times(k+1)}=11\cdot 2^{k+s+2}-11\cdot 2^{2s+4}+21\cdot 2^{3s+2}-9\cdot 2^{2s+2}
\end{aligned}$$

$$\Leftrightarrow \Gamma_{s+2}^{\left[\substack{s\\ s+2}\right]\times(k+1)}=11\cdot 2^{k+s+2}+21\cdot 2^{3s+2}-53\cdot 2^{2s+2}\quad \text{if } k>s+3. \tag{14.17}$$

By (14.10) we see that (14.17) holds for k = s+3, then (14.17) holds for $k\geq s+3$.

From (13.26) (with $k\to k-1,\quad k-1>s+1$) and (14.17) we have

$$\begin{aligned}
\Gamma_{s+2}^{\left[\substack{s\\ s+2}\right]\times k}&=11\cdot 2^{k+s+1}+21\cdot 2^{3s+2}-53\cdot 2^{2s+2}\quad \text{if } k\geq s+3,\\
8\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times(k-1)}&=8\cdot[11\cdot 2^{k+s-2}+21\cdot 2^{3s-1}-53\cdot 2^{2s-1}]\quad \text{if } k-1>s+1.
\end{aligned}$$

□

Lemma 14.2. *We have*

(14.18)
$$\Gamma_{s+3}^{\left[\substack{s\\ s+m}\right]\times(s+3)}=8^2\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+(m-2)}\right]\times(s+1)}=8^2[2^{4s+m-2}-3\cdot 2^{3s-1}+2^{2s-1}]\qquad \textit{if }\ m\geq 3,$$

(14.19)
$$\Gamma_{s+3}^{\left[\substack{s\\ s+m}\right]\times k}=8^2\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+(m-2)}\right]\times(k-2)}=8^2[3\cdot 2^{k+s-3}+21\cdot 2^{3s-1}-21\cdot 2^{2s-1}]\quad \textit{if }\ k>s+3,\quad m\geq 4,$$

(14.20)
$$\Gamma_{s+3}^{\left[\substack{s\\ s+3}\right]\times k}=8^2\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times(k-2)}=8^2[11\cdot 2^{k+s-3}+21\cdot 2^{3s-1}-53\cdot 2^{2s-1}]\qquad \textit{if }\ k>s+3.$$

Proof. Proof of (14.18)

By (14.11), (13.47) and (13.25) we get

$$\Gamma_{s+3}^{\left[\substack{s\\ s+m}\right]\times(s+3)}=2^{4s+m+4}-3\cdot 2^{3s+5}+2^{2s+5}\quad \text{if}\quad m\geq 3,$$

$$\Gamma_{s+1}^{\left[\substack{s \\ s+(m-2)}\right]\times(s+1)} = 2^{4s+m-2} - 3\cdot 2^{3s-1} + 2^{2s-1} \quad \text{if} \quad m-2 \geq 1.$$

Proof of (14.19)

Consider the matrix

$$\left(\begin{array}{cccccc}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{s+3} & \alpha_{s+4} \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+4} & \alpha_{s+5} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s+1} & \alpha_{2s+2} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s+2} & \alpha_{2s+3} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{s+3} & \beta_{s+4} \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+4} & \beta_{s+5} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s+1} & \beta_{2s+2} \\
\beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s+2} & \beta_{2s+3} \\
\beta_{s+1} & \beta_{s+2} & \beta_{s+3} & \ldots & \beta_{2s+3} & \beta_{2s+4} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{2s+m+2} & \beta_{2s+m+3}
\end{array}\right).$$

We have for $m \geq 3$, k = s+4

$$\sum_{i=0}^{s+4}\Gamma_i^{\left[\substack{s \\ s+m}\right]\times(s+4)} = 2^{4s+m+6}, \quad \text{by} \quad (11.1), \tag{14.21}$$

$$\sum_{i=0}^{s+4}\Gamma_i^{\left[\substack{s \\ s+m}\right]\times(s+4)}\cdot 2^{-i} = 2^{3s+m+2} + 2^{2s+6} - 2^{s+2} \quad \text{by} \quad (11.2), \tag{14.22}$$

$$\sum_{i=0}^{s-1}\Gamma_i^{\left[\substack{s \\ s+m}\right]\times(s+4)} = 3\cdot 2^{3s-4} - 2^{2s-3} \quad \text{by} \quad (11.27), \tag{14.23}$$

$$\sum_{i=0}^{s-1}\Gamma_i^{\left[\substack{s \\ s+m}\right]\times(s+4)}\cdot 2^{-i} = 7\cdot 2^{2s-4} - 3\cdot 2^{s-3} \quad \text{by} \quad (11.28), \tag{14.24}$$

$$\Gamma_s^{\left[\substack{s \\ s+m}\right]\times(s+4)} = 21\cdot 2^{3s-4} + 53\cdot 2^{2s-3} \quad \text{by} \quad (13.46) \text{ with k = s+4}, \tag{14.25}$$

$$\Gamma_{s+1}^{\left[\substack{s \\ s+m}\right]\times(s+4)} = 21\cdot 2^{3s-1} + 27\cdot 2^{2s-1} \quad \text{by} \quad (13.48) \text{ with k = s+4}, \tag{14.26}$$

$$\Gamma_{s+2}^{\left[\substack{s \\ s+m}\right]\times(s+4)} = 21\cdot 2^{3s+2} + 3\cdot 2^{2s+2} \quad \text{by} \quad (14.2) \text{ with k = s+4}. \tag{14.27}$$

From (14.21), ..., (14.27) we get

(14.28)

$$\begin{aligned}
&\Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)} + \Gamma_{s+4}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)} \\
&= \sum_{i=0}^{s+4}\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)} - \sum_{i=0}^{s-1}\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)} - \Gamma_{s}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)} - \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)} - \Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)} \\
&= 2^{4s+m+6} - \big(3\cdot 2^{3s-4} - 2^{2s-3} + 21\cdot 2^{3s-4} + 53\cdot 2^{2s-3} + 21\cdot 2^{3s-1} + 27\cdot 2^{2s-1} + 21\cdot 2^{3s+2} + 3\cdot 2^{2s+2}\big),
\end{aligned}$$

(14.29)

$$\begin{aligned}
&\Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)}\cdot 2^{-(s+3)} + \Gamma_{s+4}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)}\cdot 2^{-(s+4)} \\
&= \sum_{i=0}^{s+4}\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)}\cdot 2^{-i} - \sum_{i=0}^{s-1}\Gamma_{i}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)}\cdot 2^{-i} - \Gamma_{s}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)}\cdot 2^{-s} - \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)}\cdot 2^{-(s+1)} \\
&- \Gamma_{s+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)}\cdot 2^{-(s+2)} \\
&= 2^{3s+m+2} + 2^{2s+6} - 2^{s+2} - \big(7\cdot 2^{2s-4} - 3\cdot 2^{s-3} + (21\cdot 2^{3s-4} + 53\cdot 2^{2s-3})\cdot 2^{-s} \\
&+ (21\cdot 2^{3s-1} + 27\cdot 2^{2s-1})\cdot 2^{-(s+1)} + (21\cdot 2^{3s+2} + 3\cdot 2^{2s+2})\cdot 2^{-(s+2)}\big) \\
&= 2^{3s+m+2} + 9\cdot 2^{2s+2} - 5\cdot 2^{s+2}.
\end{aligned}$$

Hence by (14.28), (14.29) we deduce after some calculations

$$\Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)} = 21\cdot 2^{3s+5} - 9\cdot 2^{2s+5} \qquad \text{if } m \geq 3, \tag{14.30}$$

$$\Gamma_{s+4}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)} = 2^{4s+m+6} - 3\cdot 2^{3s+8} + 2^{2s+8} \qquad \text{if } m \geq 3. \tag{14.31}$$

By (12.34) with j = 3, $m \geq 4$ we get

$$\Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times k} = 3\cdot 2^{k+s+3} \quad \text{if} \quad k > s+3. \tag{14.32}$$

From (14.32), (14.30) we deduce

$$\begin{aligned}
&\sum_{j=s+4}^{k}\left(\Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(j+1)} - \Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times j}\right) = \sum_{j=s+4}^{k} 3\cdot 2^{j+s+3} \\
&\Leftrightarrow \sum_{j=s+5}^{k+1}\Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times j} - \sum_{j=s+4}^{k}\Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times j} = 3\cdot 2^{k+s+4} - 3\cdot 2^{2s+7} \\
&\Leftrightarrow \Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+4)} = 3\cdot 2^{k+s+4} - 3\cdot 2^{2s+7} \\
&\Leftrightarrow \Gamma_{s+3}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(k+1)} = 3\cdot 2^{k+s+4} - 3\cdot 2^{2s+7} + 21\cdot 2^{3s+5} - 9\cdot 2^{2s+5}
\end{aligned}$$

$$\Leftrightarrow \Gamma_{s+3}^{\left[\substack{s\\s+m}\right]\times(k+1)} = 3\cdot 2^{k+s+4} + 21\cdot 2^{3s+5} - 21\cdot 2^{2s+5} \quad \text{if } k > s+4. \tag{14.33}$$

By (14.30) we see that (14.33) holds for k = s+3, then (14.33) holds for $k \geq s+3$.

From (13.48) with $m \to m-2, \quad k \to k-2, \quad m \geq 4, \quad k-2 > s+1$ and (14.33) we get

$$8^2 \cdot \Gamma_{s+1}^{\left[\substack{s\\s+(m-2)}\right]\times(k-2)} = 8^2 \cdot [3\cdot 2^{k+s-3} + 21\cdot(2^{3s-1} - 2^{2s-1})], \tag{14.34}$$

$$\Gamma_{s+3}^{\left[\substack{s\\s+m}\right]\times k} = 3\cdot 2^{k+s+3} + 21\cdot 2^{3s+5} - 21\cdot 2^{2s+5}. \tag{14.35}$$

By (14.34), (14.35) we obtain for $m \geq 4, \quad k \geq s+4$

$$\Gamma_{s+3}^{\left[\substack{s\\s+m}\right]\times k} = 8^2 \cdot \Gamma_{s+1}^{\left[\substack{s\\s+(m-2)}\right]\times(k-2)}.$$

Proof of (14.20)

We have

$$\Gamma_{s+3}^{\left[\substack{s\\s+3}\right]\times(s+4)} = 21\cdot 2^{3s+5} - 9\cdot 2^{2s+5} \quad \text{by (14.30) with m = 3 ,} \tag{14.36}$$

(14.37)

$$\Gamma_{s+3}^{\left[\substack{s\\s+3}\right]\times(k+1)} - \Gamma_{s+3}^{\left[\substack{s\\s+3}\right]\times k} = 11\cdot 2^{k+s+3} \qquad \text{by (12.34) with j = m = 3, } k > s+3.$$

From (14.36), (14.37) we deduce

$$\begin{aligned}
&\sum_{j=s+4}^{k}\left(\Gamma_{s+3}^{\left[\substack{s\\s+3}\right]\times(j+1)} - \Gamma_{s+4}^{\left[\substack{s\\s+3}\right]\times j}\right) = \sum_{j=s+4}^{k} 11\cdot 2^{j+s+3}\\
&\Leftrightarrow \sum_{j=s+5}^{k+1} \Gamma_{s+3}^{\left[\substack{s\\s+3}\right]\times j} - \sum_{j=s+4}^{k} \Gamma_{s+3}^{\left[\substack{s\\s+3}\right]\times j} = 11\cdot 2^{k+s+4} - 11\cdot 2^{2s+7}\\
&\Leftrightarrow \Gamma_{s+3}^{\left[\substack{s\\s+3}\right]\times(k+1)} - \Gamma_{s+3}^{\left[\substack{s\\s+3}\right]\times(s+4)} = 11\cdot 2^{k+s+4} - 11\cdot 2^{2s+7}\\
&\Leftrightarrow \Gamma_{s+3}^{\left[\substack{s\\s+3}\right]\times(k+1)} = 11\cdot 2^{k+s+4} - 11\cdot 2^{2s+7} + 21\cdot 2^{3s+5} - 9\cdot 2^{2s+5}
\end{aligned}$$

$$\Leftrightarrow \Gamma_{s+3}^{\left[\substack{s\\s+3}\right]\times(k+1)} = 11\cdot 2^{k+s+4} + 21\cdot 2^{3s+5} - 53\cdot 2^{2s+5} \quad \text{if } k > s+4. \tag{14.38}$$

By (14.36) we see that (14.38) holds for k = s+3, then (14.38) holds for $k \geq s+3$.

From (13.26) (with $k \to k-2, \quad k-2 > s+1$) and (14.38) we have

$$\Gamma_{s+3}^{\left[\begin{smallmatrix} s\\ s+3\end{smallmatrix}\right]\times k} = 11\cdot 2^{k+s+3} + 21\cdot 2^{3s+5} - 53\cdot 2^{2s+5} \quad \text{if } k\geq s+4,$$

$$8^{2}\cdot\Gamma_{s+1}^{\left[\begin{smallmatrix} s\\ s+1\end{smallmatrix}\right]\times (k-2)} = 8^{2}\cdot[11\cdot 2^{k+s-3} + 21\cdot 2^{3s-1} - 53\cdot 2^{2s-1}] \quad \text{if } k-2> s+1.$$

$\square$

15. A REDUCTION FORMULA FOR $\Gamma_{s+j}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k}$ in the case $1\leq j\leq m,\ k\geq s+j$

In this section we prove by induction on j the following reduction formula

$$\Gamma_{s+j}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k} = 8^{j-1}\cdot\Gamma_{s+1}^{\left[\begin{smallmatrix} s\\ s+(m-(j-1))\end{smallmatrix}\right]\times (k-(j-1))} \qquad \text{if } 1\leq j\leq m,\quad k\geq s+j.$$

Recall that the right hand side in the above equation has been computed in section 13.
In fact we have

$$\Gamma_{s+1}^{\left[\begin{smallmatrix} s\\ s+(m-(j-1))\end{smallmatrix}\right]\times (k-(j-1))} = \begin{cases} 3\cdot 2^{k-j+s} + 21\cdot[2^{3s-1} - 2^{2s-1}] & \text{if } 1\leq j\leq m-1,\ k>s+j,\\ 2^{4s+(m-(j-1))} - 3\cdot 2^{3s-1} + 2^{2s-1} & \text{if } 1\leq j\leq m,\ k=s+j,\\ 11\cdot 2^{k-m+s} + 21\cdot 2^{3s-1} - 53\cdot 2^{2s-1} & \text{if } j=m,\ k>s+m. \end{cases}$$

Lemma 15.1. *We have*

$$\Gamma_{s+l}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times (s+l)} = 8^{l-1}\cdot\Gamma_{s+1}^{\left[\begin{smallmatrix} s\\ s+(m-(l-1))\end{smallmatrix}\right]\times (s+1)} \qquad \textit{if } 1\leq l\leq m, \tag{15.1}$$

$$\Gamma_{s+1}^{\left[\begin{smallmatrix} s\\ s+(m-(l-1))\end{smallmatrix}\right]\times (s+1)} = 2^{4s+(m-(l-1))} - 3\cdot 2^{3s-1} + 2^{2s-1} \qquad \textit{if } 1\leq l\leq m, \tag{15.2}$$

$$\Gamma_{s+l}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k} = 8^{l-1}\cdot\Gamma_{s+1}^{\left[\begin{smallmatrix} s\\ s+(m-(l-1))\end{smallmatrix}\right]\times (k-(l-1))} \qquad \textit{if } 1\leq l\leq m,\ k>s+l, \tag{15.3}$$

$$\Gamma_{s+1}^{\left[\begin{smallmatrix} s\\ s+(m-(l-1))\end{smallmatrix}\right]\times (k-(l-1))} = 3\cdot 2^{k-l+s} + 21\cdot[2^{3s-1} - 2^{2s-1}] \qquad \textit{if } 1\leq l\leq m-1,\ k>s+l, \tag{15.4}$$

$$\Gamma_{s+1}^{\left[\begin{smallmatrix} s\\ s+1\end{smallmatrix}\right]\times (k-(m-1))} = 11\cdot 2^{k-m+s} + 21\cdot 2^{3s-1} - 53\cdot 2^{2s-1} \quad \textit{if } l=m,\ k>s+m. \tag{15.5}$$

Proof. Let l be a rational integer such that $2 \leq l \leq m-1$; we shall prove lemma 15.1 by strong induction on l.

Assume
(15.6)

$$(H_{l-1}) \quad \Gamma_{s+j}^{\left[\substack{s \\ s+m}\right]\times k} = 8^{j-1}\cdot\Gamma_{s+1}^{\left[\substack{s \\ s+(m-(j-1))}\right]\times(k-(j-1))} \qquad for \quad 1 \leq j \leq l-1 \quad k \geq s+j.$$

We are going to show that for $2 \leq l \leq m-1$,

$$(H_{l-1}) \quad \text{implies} \quad \Gamma_{s+l}^{\left[\substack{s \\ s+m}\right]\times k} = 8^{l-1}\cdot\Gamma_{s+1}^{\left[\substack{s \\ s+(m-(l-1))}\right]\times(k-(l-1))} \text{for} \quad k \geq s+l.$$

By Lemmas $14.1, 14.2$ $\quad (H_{l-1})$ holds for l = 3,4.

The case $k = s+l$

We have

(15.7)

$$\Gamma_{j}^{\left[\substack{s \\ s+m}\right]\times(s+l)} = 21\cdot 2^{3j-4} - 3\cdot 2^{2j-3} \quad \text{if} \quad 1 \leq j \leq s-1 \quad \text{by (11.21)},$$

(15.8)

$$\Gamma_{s}^{\left[\substack{s \\ s+m}\right]\times(s+l)} = 2^{2s+l-1} + 21\cdot 2^{3s-4} - 11\cdot 2^{2s-3} \quad \text{by (13.46)},$$

(15.9)

$$\Gamma_{s+1}^{\left[\substack{s \\ s+m}\right]\times k} = 3\cdot 2^{k+s-1} + 21[2^{3s-1} - 2^{2s-1}] \quad \text{if} \quad k > s+1 \text{ by (13.48)},$$

(15.10)

$$\Gamma_{s+j}^{\left[\substack{s \\ s+m}\right]\times(s+l)} = 8^{j-1}\cdot\Gamma_{s+1}^{\left[\substack{s \\ s+(m-(j-1))}\right]\times(s+l-(j-1))}$$
$$= 3\cdot 2^{2s+l+2j-3} + 21\cdot[2^{3j+3s-4} - 2^{2s+3j-4}] \quad \text{if } 1 \leq j \leq l-1$$

by (H_{l-1}) and ((13.48) with $m \rightarrow m-(j-1) \geq 2,\ k \rightarrow s+l-(j-1),\ s+l-(j-1) > s+1)$.

Consider the matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{s+l-1} & \alpha_{s+l} \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+l} & \alpha_{s+l+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s+l-3} & \alpha_{2s+l-2} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s+l-2} & \alpha_{2s+l-1} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{s+l-1} & \beta_{s+l} \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+l} & \beta_{s+l+2} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s+l-3} & \beta_{2s+l-2} \\
\beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s+l-2} & \beta_{2s+l-1} \\
\beta_{s+1} & \beta_{s+2} & \beta_{s+3} & \ldots & \beta_{2s+l-1} & \beta_{2s+l} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{2s+m+l-2} & \beta_{2s+m+l-1}
\end{pmatrix}.$$

We have respectively by (11.1) and (11.27) with k = s+l

$$\sum_{i=0}^{s+l}\Gamma_i^{\left[\substack{s\\ s+m}\right]\times(s+l)} = 2^{4s+m+2l-2}, \tag{15.11}$$

$$\sum_{i=0}^{s-1}\Gamma_i^{\left[\substack{s\\ s+m}\right]\times(s+l)} = 3\cdot 2^{3s-4}-2^{2s-3}. \tag{15.12}$$

From $(15.8), (15.9), \ldots, (15.12)$ we get

(15.13)

$$\begin{aligned}
\Gamma_{s+l}^{\left[\substack{s\\ s+m}\right]\times(s+l)} &= 2^{4s+m+2l-2} - \sum_{j=0}^{s-1}\Gamma_j^{\left[\substack{s\\ s+m}\right]\times(s+l)} - \Gamma_s^{\left[\substack{s\\ s+m}\right]\times(s+l)} - \sum_{j=1}^{l-1}\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times(s+l)} \\
&= 2^{4s+m+2l-2} - (3\cdot 2^{3s-4}-2^{2s-3}) - (2^{2s+l-1}+21\cdot 2^{3s-4}-11\cdot 2^{2s-3}) \\
&\quad - \sum_{j=1}^{l-1}(3\cdot 2^{2s+l+2j-3}+21\cdot[2^{3j+3s-4}-2^{2s+3j-4}]) \\
&= 2^{4s+m+2l-2} - 3\cdot 2^{3s+3l-4} + 2^{2s+3l-4},
\end{aligned}$$

(15.14)

$$\Gamma_{s+1}^{\left[\substack{s\\ s+(m-(l-1))}\right]\times(s+1)} = 2^{4s+(m-(l-1))} - 3\cdot 2^{3s-1} + 2^{2s-1} \quad \text{by (13.47) since } m-(l-1)\geq 2.$$

From (15.13), (15.14) we get

$$\Gamma_{s+l}^{\left[\substack{s\\ s+m}\right]\times(s+l)} = 8^{l-1}\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+(m-(l-1))}\right]\times(s+1)} \quad \text{if } \ 1\leq l\leq m-1,\ k=s+l. \tag{15.15}$$

The case $k > s+l$
Consider the matrix

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{s+l} & \alpha_{s+l+1} \\ \alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+l+1} & \alpha_{s+l+2} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s+l-2} & \alpha_{2s+l-1} \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s+l-1} & \alpha_{2s+l} \\ \beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{s+l} & \beta_{s+l+1} \\ \beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+l+1} & \beta_{s+l+2} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s+l-2} & \beta_{2s+l-1} \\ \beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s+l-1} & \beta_{2s+l} \\ \beta_{s+1} & \beta_{s+2} & \beta_{s+3} & \ldots & \beta_{2s+l} & \beta_{2s+l+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{2s+m+l-1} & \beta_{2s+m+l} \end{pmatrix}.$$

We get

(15.16)
$$\Gamma_{s}^{\left[\substack{s \\ s+m}\right]\times(s+l+1)} = 2^{2s+l} + 21\cdot 2^{3s-4} - 11\cdot 2^{2s-3} \quad \text{by (13.46)}\ ,$$

(15.17)
$$\Gamma_{s+j}^{\left[\substack{s \\ s+m}\right]\times(s+l+1)} = 8^{j-1}\cdot\Gamma_{s+1}^{\left[\substack{s \\ s+(m-(j-1))}\right]\times(s+l+1-(j-1))}$$
$$= 3\cdot 2^{2s+l+2j-2} + 21\cdot\left[2^{3j+3s-4} - 2^{2s+3j-4}\right] \quad \text{if } 1\leq j\leq l-1$$
by (H) and ((13.48) with $m \rightarrow m-(j-1)\geq 2,\ k\rightarrow s+l+1-(j-1),\ s+l+1-(j-1) > s+1$).

We have respectively by (11.1), (11.2), (11.27) and (11.28) with $m\geq 2$, k = s + l +1

$$\sum_{i=0}^{s+l+1}\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times(s+l+1)} = 2^{4s+m+2l}, \tag{15.18}$$

$$\sum_{i=0}^{s+l+1}\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times(s+l+1)}\cdot 2^{-i} = 2^{3s+m+l-1} + 2^{2s+2l} - 2^{s+l-1}, \tag{15.19}$$

$$\sum_{i=0}^{s-1}\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times(s+l+1)} = 3\cdot 2^{3s-4} - 2^{2s-3} \tag{15.20}$$

and

$$\sum_{i=0}^{s-1}\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times(s+l+1)}\cdot 2^{-i} = 7\cdot 2^{2s-4} - 3\cdot 2^{s-3}. \tag{15.21}$$

From $(15.16),\ldots,(15.21)$ we obtain

(15.22)

$$
\begin{aligned}
&\Gamma_{s+l}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times(s+l+1)} + \Gamma_{s+l+1}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times(s+l+1)}\\
&= 2^{4s+m+2l} - \sum_{j=0}^{s-1}\Gamma_{j}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times(s+l+1)} - \Gamma_{s}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times(s+l+1)} - \sum_{j=1}^{l-1}\Gamma_{s+j}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times(s+l+1)}\\
&= 2^{4s+m+2l} - (3\cdot 2^{3s-4} - 2^{2s-3}) - (2^{2s+l} + 21\cdot 2^{3s-4} - 11\cdot 2^{2s-3})\\
&- \sum_{j=1}^{l-1}(3\cdot 2^{2s+l+2j-2} + 21\cdot[2^{3j+3s-4} - 2^{2s+3j-4}])\\
&= 2^{4s+m+2l} - 3\cdot 2^{3s+3l-4} - 2^{2s+3l-4},
\end{aligned}
$$

(15.23)

$$
\begin{aligned}
&\Gamma_{s+l}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times(s+l+1)}\cdot 2^{-(s+l)} + \Gamma_{s+l+1}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times(s+l+1)}\cdot 2^{-(s+l+1)}\\
&= \sum_{j=0}^{s+l+1}\Gamma_{j}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times(s+l+1)}\cdot 2^{-j} - \sum_{j=0}^{s-1}\Gamma_{j}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times(s+l+1)}\cdot 2^{-j} - \Gamma_{s}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times(s+l+1)}\cdot 2^{-s} - \sum_{j=1}^{l-1}\Gamma_{s+j}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times(s+l+1)}\cdot 2^{-(s+j)}\\
&= 2^{3s+m+l-1} + 2^{2s+2l} - 2^{s+l-1} - (7\cdot 2^{2s-4} - 3\cdot 2^{s-3}) - (2^{s+l} + 21\cdot 2^{2s-4} - 11\cdot 2^{s-3})\\
&- \sum_{j=1}^{l-1}(3\cdot 2^{2s+l+2j-2} + 21\cdot[2^{3j+3s-4} - 2^{2s+3j-4}])\cdot 2^{-(s+j)}\\
&= 2^{4s+m+2l} + 9\cdot 2^{3s+3l-3} - 5\cdot 2^{2s+3l-3}.
\end{aligned}
$$

Hence by (15.22), (15.23) we deduce after some calculations

$$
\Gamma_{s+l}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times(s+l+1)} = 21\cdot 2^{3s+3l-4} - 9\cdot 2^{2s+3l-4}, \tag{15.24}
$$

$$
\Gamma_{s+l+1}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times(s+l+1)} = 2^{4s+m+2l} - 3\cdot 2^{3s+3l-1} + 2^{2s+3l-1}. \tag{15.25}
$$

By (12.34) with $j = l$, $m \geq l+1$ we get

$$
\Gamma_{s+l}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+l}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times k} = 3\cdot 2^{k+s+2l-3} \quad \text{if} \quad k > s+l. \tag{15.26}
$$

From (15.26), (15.24) we deduce

$$
\begin{aligned}
&\sum_{j=s+l+1}^{k}\left(\Gamma_{s+l}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times(j+1)} - \Gamma_{s+l}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times j}\right) = \sum_{j=s+l+1}^{k} 3\cdot 2^{j+s+2l-3}\\
&\Leftrightarrow \sum_{j=s+l+2}^{k+1}\Gamma_{s+l}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times j} - \sum_{j=s+l+1}^{k}\Gamma_{s+l}^{\left[\begin{smallmatrix}s\\s+m\end{smallmatrix}\right]\times j} = 3\cdot 2^{k+s+2l+2} - 3\cdot 2^{2s+3l-2}
\end{aligned}
$$

$$\Leftrightarrow \Gamma_{s+l}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{s+l}^{\left[\substack{s\\ s+m}\right]\times(s+l+1)} = 3\cdot 2^{k+s+2l+2} - 3\cdot 2^{2s+3l-2}$$

$$\Leftrightarrow \Gamma_{s+l}^{\left[\substack{s\\ s+m}\right]\times(k+1)} = 3\cdot 2^{k+s+2l+2} - 3\cdot 2^{2s+3l-2} + 21\cdot 2^{3s+3l-4} - 9\cdot 2^{2s+3l-4}$$

$$\Leftrightarrow \Gamma_{s+l}^{\left[\substack{s\\ s+m}\right]\times(k+1)} = 3\cdot 2^{k+s+2l-2} + 21\cdot 2^{3s+3l-4} - 21\cdot 2^{2s+3l-4} \quad \text{if } k > s+l. \tag{15.27}$$

By (15.24) we see that (15.27) holds for $k = s+l$, then (15.27) holds for $k \geq s+l$.

From (13.48) with $m \to m-(l-1)$, $k \to k-(l-1)$, $m \geq l+1$, $k-(l-1) > s+1$ and (15.27) we get

(15.28)

$$8^{l-1}\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+(m-(l-1))}\right]\times(k-(l-1))} = 8^{l-1}\cdot[3\cdot 2^{k-(l-1)+s-1} + 21\cdot(2^{3s-1} - 2^{2s-1})],$$

$$\Gamma_{s+l}^{\left[\substack{s\\ s+m}\right]\times k} = 3\cdot 2^{k+s+2l-3} + 21\cdot 2^{3s+3l-4} - 21\cdot 2^{2s+3l-4}. \tag{15.29}$$

By (15.28), (15.29) we obtain

$$\Gamma_{s+l}^{\left[\substack{s\\ s+m}\right]\times k} = 8^{l-1}\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+(m-(l-1))}\right]\times(k-(l-1))} \quad \text{for} \quad l \leq m-1, \quad k > s+l. \tag{15.30}$$

Observe that in view of (15.30) the formulas (15.24) and (15.25) holds for l = m, it suffices to reproduce carefully the proof of the above formulas with $k \to s+m+1$ instead of $k \to s+l+1$.
In fact we have

$$\Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times(s+m+1)} = 21\cdot 2^{3s+3m-4} - 9\cdot 2^{2s+3m-4}, \tag{15.31}$$

$$\Gamma_{s+m+1}^{\left[\substack{s\\ s+m}\right]\times(s+m+1)} = 2^{4s+3m} - 3\cdot 2^{3s+3m-1} + 2^{2s+3m-1}. \tag{15.32}$$

The case $k = s+m,\ l = m$
We have

(15.33)

$$\Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times(s+m)} = 2^{4s+3m-2} - 3\cdot 2^{3s+3m-4} + 2^{2s+3m-4} \quad \text{by (15.25) with } l \to m-1\ ,$$

(15.34)

$$\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times(s+1)} = 2^{4s+1} - 3\cdot 2^{3s-1} + 2^{2s-1} \quad \text{by (13.25).}$$

From (15.33), (15.34) we obtain

$$\Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times(s+m)} = 8^{m-1}\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times(s+1)}.$$

The case $k > s+m,\ l = m$
We have

(15.35)
$$\Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times(s+m+1)} = 21\cdot 2^{3s+3m-4} - 9\cdot 2^{2s+3m-4} \quad \text{by (15.31)},$$

(15.36)
$$\Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times k} = 11\cdot 2^{k+s+2m-3} \qquad \text{by (12.34) with j = m ,}\ \ k > s+m.$$

From (15.35), (15.36) we deduce

$$\begin{aligned}
&\sum_{j=s+m+1}^{k}\left(\Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times(j+1)} - \Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times j}\right) = \sum_{j=s+m+1}^{k} 11\cdot 2^{j+s+2m-3}\\
&\Leftrightarrow \sum_{j=s+m+2}^{k+1}\Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times j} - \sum_{j=s+m+1}^{k}\Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times j} = 11\cdot 2^{k+s+2m-2} - 11\cdot 2^{2s+3m-2}\\
&\Leftrightarrow \Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times(s+m+1)} = 11\cdot 2^{k+s+2m-2} - 11\cdot 2^{2s+3m-2}\\
&\Leftrightarrow \Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times(k+1)} = 11\cdot 2^{k+s+2m-2} - 11\cdot 2^{2s+3m-2} + 21\cdot 2^{3s+3m-4} - 9\cdot 2^{2s+3m-4}
\end{aligned}$$

(15.37)
$$\Leftrightarrow \Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times(k+1)} = 11\cdot 2^{k+s+2m-2} + 21\cdot 2^{3s+3m-4} - 53\cdot 2^{2s+3m-4} \quad \text{if } k > s+m.$$

By (15.31) we see that (15.37) holds for k = s + m, then (15.37) holds for $k \geq s+m$.
From (13.26) (with $k \rightarrow k-(m-1)$, $\quad k-(m-1) > s+1$) and (15.37) with $k+1 \rightarrow k$
we have

$$\begin{aligned}
\Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times k} &= 11\cdot 2^{k+s+2m-3} + 21\cdot 2^{3s+3m-4} - 53\cdot 2^{2s+3m-4} \quad \text{if } k > s+m,\\
8^{m-1}\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+1}\right]\times(k-(m-1))} &= 8^{m-1}\cdot[11\cdot 2^{k-(m-1)+s-1} + 21\cdot 2^{3s-1} - 53\cdot 2^{2s-1}] \quad \text{if } k-(m-1) > s+1.
\end{aligned}$$

□

Lemma 15.2. *We have*

(15.38)
$$\Gamma_{s+l}^{\left[\substack{s\\ s+m}\right]\times k} = 8^{l-1}\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+(m-(l-1))}\right]\times(k-(l-1))} \qquad \textit{if} \quad 1 \leq l \leq m,\ k \geq s+l,$$

(15.39)
$$\Gamma_{s+l}^{\left[\substack{s\\ s+m}\right]\times(s+l)} = 2^{4s+2l+m-2} - 3\cdot 2^{3s+3l-4} + 2^{2s+3l-4} \qquad \textit{if}\ \ 1\leq l\leq m,$$

(15.40)
$$\Gamma_{s+l}^{\left[\substack{s\\ s+m}\right]\times k} = 3\cdot 2^{k+2l+s-3} + 21\cdot 2^{3s+3l-4} - 21\cdot 2^{2s+3l-4} \qquad \textit{if}\ \ 1\leq l\leq m-1,\ k>s+l,$$

(15.41)
$$\Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times k} = 11\cdot 2^{k+2m+s-3} + 21\cdot 2^{3s+3m-4} - 53\cdot 2^{2s+3m-4} \quad \textit{if}\ \ k>s+m.$$

Proof. Lemma 15.2 follows from Lemma 15.1. □

16. A REDUCTION FORMULA FOR $\Gamma_{s+j+1}^{\left[\substack{s\\ s}\right]\times k}$ IN THE CASE $1\leq j\leq s-1,\ k\geq s+j+1$

In this section we prove the following reduction formula by induction on j

$$\Gamma_{s+j+1}^{\left[\substack{s\\ s}\right]\times k} = 8^{2j}\cdot\Gamma_{s-j+1}^{\left[\substack{s-j\\ s-j}\right]\times(k-2j)} \qquad \text{if}\ \ 0\leq j\leq s-1,\ k\geq s+j+1.$$

We recall once more that the right hand side in the above equation has been computed in section 13.

In fact we have

$$\Gamma_{s-j+1}^{\left[\substack{s-j\\ s-j}\right]\times(k-2j)} = \begin{cases} 2^{4s-4j} - 3\cdot 2^{3s-3j-1} + 2^{2s-2j-1} & \text{if} \quad 0\leq j\leq s-1,\ k=s+j+1,\\ 21\cdot[2^{k+s-3j-1} + 2^{3s-3j-1} - 5\cdot 2^{2s-2j-1}] & \text{if} \quad 0\leq j\leq s-2,\ k>s+j+1,\\ 2^{2k-4s+4} - 3\cdot 2^{k-2s+2} + 2 & \text{if} \quad j=s-1,\ k>2s. \end{cases}$$

Lemma 16.1. *We have*

$$\Gamma_{s+2}^{\left[\substack{s\\ s}\right]\times k} = 8^2\cdot\Gamma_{s}^{\left[\substack{s-1\\ s-1}\right]\times(k-2)} \qquad \textit{if}\quad k\geq s+2, \tag{16.1}$$

$$\Gamma_{s+2}^{\left[\substack{s\\ s}\right]\times(s+2)} = 2^{4s+2} - 3\cdot 2^{3s+2} + 2^{2s+3} \qquad \textit{if}\quad k\ =\ s{+}2\ , \tag{16.2}$$

$$\Gamma_{s+2}^{\left[\substack{s\\ s}\right]\times k} = 21\cdot[2^{k+s+2} + 2^{3s+2} - 5\cdot 2^{2s+3}] \qquad \textit{if}\ \ k>s+2. \tag{16.3}$$

Proof. Proof of (16.1) with k = s+2

From (13.20) and (13.3) (with $s\to s-1$) we get

$$\Gamma_{s+2}^{\left[\substack{s\\s}\right]\times(s+2)} = 2^{4s+2} - 3\cdot 2^{3s+2} + 2^{2s+3},$$

$$8^2\cdot\Gamma_{s}^{\left[\substack{s-1\\s-1}\right]\times s} = 8^2\cdot[2^{4s-4} - 3\cdot 2^{3s-4} + 2^{2s-3}].$$

Proof of (16.1) with $k > s+2$

Consider the matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{s+2} & \alpha_{s+3}\\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+3} & \alpha_{s+4}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s} & \alpha_{2s+1}\\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s+1} & \alpha_{2s+2}\\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{s+2} & \beta_{s+3}\\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+3} & \beta_{s+4}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s} & \beta_{2s+1}\\
\beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s+1} & \beta_{2s+2}
\end{pmatrix}.$$

We have

$$\sum_{i=0}^{s+3}\Gamma_{i}^{\left[\substack{s\\s}\right]\times(s+3)} = 2^{4s+4} \qquad \text{by (11.1) with m = 0, k = s+3,} \tag{16.4}$$

$$\sum_{i=0}^{s+3}\Gamma_{i}^{\left[\substack{s\\s}\right]\times(s+3)}\cdot 2^{-i} = 2^{3s+1} + 2^{2s+4} - 2^{s+1} \qquad \text{by (11.2) with m = 0, k = s+3,} \tag{16.5}$$

$$\sum_{i=0}^{s-1}\Gamma_{i}^{\left[\substack{s\\s}\right]\times(s+3)} = 3\cdot 2^{3s-4} - 2^{2s-3} \qquad \text{by (11.27) with m = 0, k = s+3,} \tag{16.6}$$

$$\sum_{i=0}^{s-1}\Gamma_{i}^{\left[\substack{s\\s}\right]\times(s+3)}\cdot 2^{-i} = 7\cdot 2^{2s-4} - 3\cdot 2^{s-3} \qquad \text{by (11.28) with m = 0, k = s+3,} \tag{16.7}$$

$$\Gamma_{s}^{\left[\substack{s\\s}\right]\times(s+3)} = 21\cdot 2^{3s-4} + 69\cdot 2^{2s-3} \qquad \text{by (13.2) with k = s+3,} \tag{16.8}$$

$$\Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(s+3)} = 21\cdot[2^{3s-1} + 3\cdot 2^{2s-1}] \qquad \text{by (13.4) with k = s+3.} \tag{16.9}$$

From (16.4), . . . , (16.9) with k = s+3 we obtain

(16.10)

$$\begin{aligned}&\Gamma_{s+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(s+3)}+\Gamma_{s+3}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(s+3)}\\&=2^{4s+4}-(3\cdot 2^{3s-4}-2^{2s-3})-(21\cdot 2^{3s-4}+69\cdot 2^{2s-3})-(21\cdot[2^{3s-1}+3\cdot 2^{2s-1}]),\end{aligned}$$

(16.11)

$$\begin{aligned}&\Gamma_{s+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(s+3)}\cdot 2^{-(s+2)}+\Gamma_{s+3}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(s+3)}\cdot 2^{-(s+3)}\\&=2^{3s+1}+2^{2s+4}-2^{s+1}-(7\cdot 2^{2s-4}-3\cdot 2^{s-3})-(21\cdot 2^{3s-4}+69\cdot 2^{2s-3})\cdot 2^{-s}\\&-(21\cdot[2^{3s-1}+3\cdot 2^{2s-1}])\cdot 2^{-(s+1)}.\end{aligned}$$

Hence by (16.10), (16.11) we deduce after some calculations

$$\Gamma_{s+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(s+3)}=21\cdot 2^{3s+2}-21\cdot 2^{2s+3}, \tag{16.12}$$

$$\Gamma_{s+3}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(s+3)}=2^{4s+4}-3\cdot 2^{3s+5}+2^{2s+7}. \tag{16.13}$$

By (12.32) with j = 2, $s\geq 3$ we get

$$\Gamma_{s+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(k+1)}-\Gamma_{s+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times k}=21\cdot 2^{k+s+2}\quad\text{if}\quad k>s+2. \tag{16.14}$$

From (16.14), (16.12) we deduce

$$\begin{aligned}&\sum_{j=s+3}^{k}\left(\Gamma_{s+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(j+1)}-\Gamma_{s+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times j}\right)=\sum_{j=s+3}^{k}21\cdot 2^{j+s+2}\\&\Leftrightarrow\sum_{j=s+4}^{k+1}\Gamma_{s+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times j}-\sum_{j=s+2}^{k}\Gamma_{s+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times j}=21\cdot 2^{k+s+3}-21\cdot 2^{2s+5}\\&\Leftrightarrow\Gamma_{s+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(k+1)}-\Gamma_{s+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(s+3)}=21\cdot 2^{k+s+3}-21\cdot 2^{2s+5}\\&\Leftrightarrow\Gamma_{s+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(k+1)}=21\cdot 2^{k+s+3}-21\cdot 2^{2s+5}+21\cdot(2^{3s+2}-21\cdot 2^{2s+3})\end{aligned}$$

$$\Leftrightarrow\Gamma_{s+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(k+1)}=21\cdot[2^{k+s+3}+2^{3s+2}-5\cdot 2^{2s+3}]\quad\text{if } k>s+2. \tag{16.15}$$

By (16.12) we see that (16.15) holds for k = s+ 2, then (16.15) holds for $k\geq s+2$.

From (13.4) (with $k\to k-2,\quad s\to s-1,\quad k-2>(s-1)+1$) and (16.15) we have

$$\begin{aligned}\Gamma_{s+2}^{\left[\begin{smallmatrix} s\\ s\end{smallmatrix}\right]\times k} &= 21\cdot[2^{k+s+2}+2^{3s+2}-5\cdot 2^{2s+3}],\\ 8^{2}\cdot\Gamma_{s}^{\left[\begin{smallmatrix} s-1\\ s-1\end{smallmatrix}\right]\times (k-2)} &= 8^{2}\cdot 21\cdot[2^{k+s-4}+2^{3s-4}-5\cdot 2^{2s-3}].\end{aligned}$$

□

Lemma 16.2. *We have for* $s\geq 2$

(16.16)
$$\Gamma_{s+j+1}^{\left[\begin{smallmatrix} s\\ s\end{smallmatrix}\right]\times k} = 8^{2j}\cdot\Gamma_{s-j+1}^{\left[\begin{smallmatrix} s-j\\ s-j\end{smallmatrix}\right]\times (k-2j)} \qquad \textit{if}\ \ 0\leq j\leq s-1,\ k\geq s+j+1,$$

(16.17)
$$\Gamma_{s-j+1}^{\left[\begin{smallmatrix} s-j\\ s-j\end{smallmatrix}\right]\times (s-j+1)} = 2^{4s-4j}-3\cdot 2^{3s-3j-1}+2^{2s-2j-1} \qquad \textit{if}\ \ 0\leq j\leq s-1,\ k=s+j+1\ ,$$

(16.18)
$$\Gamma_{s-j+1}^{\left[\begin{smallmatrix} s-j\\ s-j\end{smallmatrix}\right]\times (k-2j)} = 21\cdot[2^{k-3j+s-1}+2^{3s-3j-1}-5\cdot 2^{2s-2j-1}] \qquad \textit{if}\ \ 0\leq j\leq s-2,\ k>s+j+1,$$

(16.19)
$$\Gamma_{s-j+1}^{\left[\begin{smallmatrix} s-j\\ s-j\end{smallmatrix}\right]\times (k-2j)} = \Gamma_{2}^{\left[\begin{smallmatrix} 1\\ 1\end{smallmatrix}\right]\times (k-2(s-1))} = 2^{2k-4s+4}-3\cdot 2^{k-2s+2}+2 \qquad \textit{if}\ j=s-1, k>2s.$$

Proof. We proceed as in section 15 by induction on j.
Let l be a rational integer such that $2\leq l\leq s-2$.
Assume

$$(16.20)\quad (H_{l-1})\quad \Gamma_{s+j+1}^{\left[\begin{smallmatrix} s\\ s\end{smallmatrix}\right]\times k} = 8^{2j}\cdot\Gamma_{s-j+1}^{\left[\begin{smallmatrix} s-j\\ s-j\end{smallmatrix}\right]\times (k-2j)} \qquad \textit{for}\quad 0\leq j\leq l-1,\quad k\geq s+j+1.$$

We are going to show that (H_l) holds, that is

$$(16.21)\qquad \Gamma_{s+l+1}^{\left[\begin{smallmatrix} s\\ s\end{smallmatrix}\right]\times k} = 8^{2l}\cdot\Gamma_{s-l+1}^{\left[\begin{smallmatrix} s-l\\ s-l\end{smallmatrix}\right]\times (k-2l)} \qquad \textit{for}\ k\geq s+l+1.$$

By Lemma 16.1 (H_{l-1}) holds for $l=2$ (see (16.20)).

Proof of (16.21) with $k=s+l+1$

Consider the matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{s+l} & \alpha_{s+l+1} \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+l+1} & \alpha_{s+l+2} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s+2l-2} & \alpha_{2s+2l-1} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s+2l-1} & \alpha_{2s+2l} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{s+l} & \beta_{s+l+1} \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+l+1} & \beta_{s+l+2} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s+2l-2} & \beta_{2s+2l-1} \\
\beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s+2l-1} & \beta_{2s+2l}
\end{pmatrix}.$$

We have

(16.22)

$$\sum_{i=0}^{s+l+1}\Gamma_{i}^{\left[\substack{s\\ s}\right]\times(s+l+1)} = 2^{4s+2l} \qquad \text{by (11.1) with m} = 0,\ k = s+l+1,$$

(16.23)

$$\sum_{i=0}^{s-1}\Gamma_{i}^{\left[\substack{s\\ s}\right]\times(s+l+1)} = 3\cdot 2^{3s-4} - 2^{2s-3} \qquad \text{by (11.27) with m} = 0,\ k = s+l+1,$$

(16.24)

$$\Gamma_{s}^{\left[\substack{s\\ s}\right]\times(s+l+1)} = 3\cdot 2^{2s+l} + 21\cdot 2^{3s-4} - 27\cdot 2^{2s-3} \qquad \text{by (13.2) with } k = s+l+1,$$

(16.25)

$$\begin{aligned}
\Gamma_{s+j+1}^{\left[\substack{s\\ s}\right]\times(s+l+1)} &= 8^{2j}\cdot\Gamma_{s-j+1}^{\left[\substack{s-j\\ s-j}\right]\times(s+l+1-2j)} && \text{by (16.20) for } 0\leq j\leq l-1,\ k=s+l+1 \\
&= 2^{6j}\cdot[21\cdot(2^{(s+l+1-2j)+(s-j)-1} + 2^{3(s-j)-1} - 5\cdot 2^{2(s-j)-1})] && \text{by (13.4) with } s\to s-j,\ k\to s+l+1-2j > s-j+1 \\
&= 21\cdot[2^{2s+3j+l} + 2^{3s+3j-1} - 5\cdot 2^{2s+4j-1}].
\end{aligned}$$

From (16.22), (16.23), (16.24) and (16.25) we get after some calculations

(16.26)

$$\begin{aligned}
\Gamma_{s+l+1}^{\left[\substack{s\\ s}\right]\times(s+l+1)} &= 2^{4s+2l} - (3\cdot 2^{3s-4} - 2^{2s-3}) - (3\cdot 2^{2s+l} + 21\cdot 2^{3s-4} - 27\cdot 2^{2s-3}) \\
&\quad - \left(\sum_{j=0}^{l-1} 21\cdot[2^{2s+3j+l} + 2^{3s+3j-1} - 5\cdot 2^{2s+4j-1}]\right) \\
&= 2^{4s+2l} - 3\cdot 2^{3s+3l-1} + 2^{2s+4l-1}.
\end{aligned}$$

By (13.3) with $s\to s-l$ and (16.26) we obtain

$$\begin{aligned}\Gamma_{s+l+1}^{\left[\substack{s\\s}\right]\times(s+l+1)} &= 8^{2l}\cdot\Gamma_{s-l+1}^{\left[\substack{s-l\\s-l}\right]\times(s-l+1)} = 2^{6l}\cdot[2^{4(s-l)}-3\cdot 2^{3(s-l)-1}+2^{2(s-l)-1}]\\ &= 2^{4s+2l}-3\cdot 2^{3s+3l-1}+2^{2s+4l-1}. \qquad \square\end{aligned}$$

Proof of (16.21) with $k > s+l+1$

Consider the matrix

$$\begin{pmatrix}\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{s+l+1} & \alpha_{s+l+2}\\ \alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+l+2} & \alpha_{s+l+3}\\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s+l-1} & \alpha_{2s+l}\\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s+l} & \alpha_{2s+l+1}\\ \beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{s+l+1} & \beta_{s+l+2}\\ \beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+l+2} & \beta_{s+l+3}\\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\ \beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s+l-1} & \beta_{2s+l}\\ \beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s+l} & \beta_{2s+l+1}\end{pmatrix}.$$

We have

(16.27)

$$\sum_{i=0}^{s+l+2}\Gamma_i^{\left[\substack{s\\s}\right]\times(s+l+2)} = 2^{4s+2l+2} \qquad \text{by (11.1) with m} = 0,\ k = s+l+2,$$

(16.28)

$$\sum_{i=0}^{s+l+2}\Gamma_i^{\left[\substack{s\\s}\right]\times(s+l+2)}\cdot 2^{-i} = 2^{3s+l}+2^{2s+2l+2}-2^{s+l} \qquad \text{by (11.2) with m} = 0,\ k = s+l+2,$$

(16.29)

$$\sum_{i=0}^{s-1}\Gamma_i^{\left[\substack{s\\s}\right]\times(s+l+2)} = 3\cdot 2^{3s-4}-2^{2s-3} \qquad \text{by (11.27) with m} = 0,\ k = s+l+2,$$

(16.30)

$$\sum_{i=0}^{s-1}\Gamma_i^{\left[\substack{s\\s}\right]\times(s+l+2)}\cdot 2^{-i} = 7\cdot 2^{2s-4}-3\cdot 2^{s-3} \qquad \text{by (11.28) with m} = 0,\ k = s+l+2,$$

(16.31)

$$\Gamma_s^{\left[\substack{s\\s}\right]\times(s+l+2)} = 3\cdot 2^{2s+l+1}+21\cdot 2^{3s-4}-27\cdot 2^{2s-3} \qquad \text{by (13.2) with } k = s+l+2,$$

(16.32)

$$\Gamma_{s+j+1}^{\left[\substack{s\\s}\right]\times(s+l+2)} = 8^{2j}\cdot\Gamma_{s-j+1}^{\left[\substack{s-j\\s-j}\right]\times(s+l+2-2j)} \qquad \text{by (16.20) for } 0\leq j\leq l-1,\ k = s+l+2$$

$$\begin{aligned}&= 2^{6j}\cdot[21\cdot(2^{(s+l+2-2j)+(s-j)-1}+2^{3(s-j)-1}-5\cdot 2^{2(s-j)-1})] \quad \text{by (13.4) with } s\to s-j,\ k\to s+l+2-2j>s-j+1\\ &= 21\cdot[2^{2s+3j+l+1}+2^{3s+3j-1}-5\cdot 2^{2s+4j-1}].\end{aligned}$$

From $(16.27),\ldots,(16.32)$ with $k=s+l+2$ we obtain

(16.33)

$$\begin{aligned}&\Gamma_{s+l+1}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(s+l+2)}+\Gamma_{s+l+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(s+l+2)}\\ &= 2^{4s+2l+2}-(3\cdot2^{3s-4}-2^{2s-3})-(3\cdot2^{2s+l+1}+21\cdot2^{3s-4}-27\cdot2^{2s-3})\\ &-\sum_{j=0}^{l-1}21\cdot[2^{2s+3j+l+1}+2^{3s+3j-1}-5\cdot2^{2s+4j-1}],\end{aligned}$$

(16.34)

$$\begin{aligned}&\Gamma_{s+l+1}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(s+l+2)}\cdot2^{-(s+l+1)}+\Gamma_{s+l+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(s+l+2)}\cdot2^{-(s+l+2)}\\ &= 2^{3s+l}+2^{2s+2l+2}-2^{s+l}-(7\cdot2^{2s-4}-3\cdot2^{s-3})-(3\cdot2^{2s+l+1}+21\cdot2^{3s-4}-27\cdot2^{2s-3})\cdot2^{-s}\\ &-\sum_{j=0}^{l-1}21\cdot[2^{2s+3j+l+1}+2^{3s+3j-1}-5\cdot2^{2s+4j-1}]\cdot2^{-(s+j+1)}.\end{aligned}$$

Hence by (16.33), (16.34) we deduce after some calculations

$$\Gamma_{s+l+1}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(s+l+2)}=21\cdot2^{3s+3l-1}-21\cdot2^{2s+4l-1}, \tag{16.35}$$

$$\Gamma_{s+l+2}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(s+l+2)}=2^{4s+2l+2}-3\cdot2^{3s+3l+2}+2^{2s+4l+3}. \tag{16.36}$$

By (12.32) with $j=l+1,\ l+1\leq s-1$ we get

$$\Gamma_{s+l+1}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(k+1)}-\Gamma_{s+l+1}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times k}=21\cdot2^{k+s+3(l+1)-4}\quad\text{if}\quad k>s+l+1. \tag{16.37}$$

From (16.37), (16.35) we deduce

$$\begin{aligned}&\sum_{j=s+l+2}^{k}\left(\Gamma_{s+l+1}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(j+1)}-\Gamma_{s+l+1}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times j}\right)=\sum_{j=s+l+2}^{k}21\cdot2^{j+s+3l-1}\\ &\Leftrightarrow\sum_{j=s+l+3}^{k+1}\Gamma_{s+l+1}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times j}-\sum_{j=s+l+1}^{k}\Gamma_{s+l+1}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times j}=21\cdot2^{k+s+3l}-21\cdot2^{2s+4l+1}\\ &\Leftrightarrow\Gamma_{s+l+1}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(k+1)}-\Gamma_{s+l+1}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(s+l+2)}=21\cdot2^{k+s+3l}-21\cdot2^{2s+4l+1}\\ &\Leftrightarrow\Gamma_{s+l+1}^{\left[\begin{smallmatrix}s\\s\end{smallmatrix}\right]\times(k+1)}=21\cdot2^{k+s+3l}-21\cdot2^{2s+4l+1}+21\cdot(2^{3s+3l-1}-21\cdot2^{2s+4l-1})\end{aligned}$$

$$\Leftrightarrow \Gamma_{s+l+1}^{\left[\substack{s\\s}\right]\times(k+1)} = 21[2^{k+s+3l} + 2^{3s+3l-1} - 5\cdot 2^{2s+4l-1}] \quad \text{if } k > s+l+1. \tag{16.38}$$

By (16.35) we see that (16.38) holds for $k = s+l+1$, then (16.38) holds for $k \geq s+l+1$.

From (16.38) and (13.4) with $s \to s-l, k \to k-2l > s-l+1$ we get

$$\begin{aligned}\Gamma_{s+l+1}^{\left[\substack{s\\s}\right]\times k} &= 8^{2l}\cdot\Gamma_{s-l+1}^{\left[\substack{s-l\\s-l}\right]\times(k-2l)} = 2^{6l}\cdot[21\cdot(2^{k-2l+(s-l)-1} + 2^{3(s-l)-1} - 5\cdot 2^{2(s-l)-1})]\\ &= 21[2^{k+s+3l-1} + 2^{3s+3l-1} - 5\cdot 2^{2s+4l-1}].\end{aligned}$$

□

We have now established that

$$\Gamma_{s+j+1}^{\left[\substack{s\\s}\right]\times k} = 8^{2j}\cdot\Gamma_{s-j+1}^{\left[\substack{s-j\\s-j}\right]\times(k-2j)} \quad \text{if } 0\leq j\leq s-2,\ k\geq s+j+1. \tag{16.39}$$

Proof of (16.16) with $j = s-1,\ k \geq 2s$

We shall show that

$$\Gamma_{2s}^{\left[\substack{s\\s}\right]\times k} = 8^{2s-2}\cdot\Gamma_{2}^{\left[\substack{1\\1}\right]\times(k-2(s-1))} \quad \text{if } k\geq 2s. \tag{16.40}$$

If k =2s, (16.40) follows from (16.21) with $l = s-1$ and k = s +(s -1)+1 = 2s.
Let $k > 2s$.
Consider the matrix

$$\begin{pmatrix}\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{2s} & \alpha_{2s+1}\\ \alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{2s+1} & \alpha_{2s+2}\\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{3s-2} & \alpha_{3s-1}\\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{3s-1} & \alpha_{3s}\\ \beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{2s} & \beta_{2s+1}\\ \beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{2s+1} & \beta_{2s+2}\\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\ \beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{3s-2} & \beta_{3s-1}\\ \beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{3s-1} & \beta_{3s}\end{pmatrix}.$$

We have

(16.41)

$$\sum_{i=0}^{2s}\Gamma_{i}^{\left[\substack{s\\s}\right]\times(2s+1)} = 2^{6s} \qquad \text{by (11.1) with m = 0, k = 2s +1,}$$

(16.42)

$$\sum_{i=0}^{s-1}\Gamma_{i}^{\left[\substack{s\\s}\right]\times(2s+1)} = 3\cdot 2^{3s-4} - 2^{2s-3} \qquad \text{by (11.27) with m = 0, k = 2s +1,}$$

(16.43)

$$\Gamma_{s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(2s+1)} = 3\cdot 2^{3s}+21\cdot 2^{3s-4}-27\cdot 2^{2s-3} \qquad \text{by (13.2) with k = 2s +1,}$$

(16.44)

$$\begin{aligned}
\Gamma_{s+j+1}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(2s+1)} &= 8^{2j}\cdot\Gamma_{s-j+1}^{\left[\begin{smallmatrix} s-j \\ s-j \end{smallmatrix}\right]\times(2s+1-2j)} && \text{by (16.20) for } 0\leq j\leq s-2,\ k>2s+1 \\
&= 2^{6j}\cdot[21\cdot(2^{(2s+1-2j)+(s-j)-1}+2^{3(s-j)-1}-5\cdot 2^{2(s-j)-1})] && \text{by (13.4) with } s\rightarrow s-j,\ k\rightarrow 2s+1-2j>s-j+1 \\
&= 21\cdot[2^{3s+3j}+2^{3s+3j-1}-5\cdot 2^{2s+4j-1}].
\end{aligned}$$

From (16.41), (16.42), (16.43) and (16.44) we get after some calculations

$$\begin{aligned}
\Gamma_{2s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(2s+1)} &= 2^{6s}-(3\cdot 2^{3s-4}-2^{2s-3})-(3\cdot 2^{3s}+21\cdot 2^{3s-4}-27\cdot 2^{2s-3}) \\
&\quad -\sum_{j=0}^{s-2}21\cdot[2^{3s+3j}+2^{3s+3j-1}-5\cdot 2^{2s+4j-1}] \\
&= 21\cdot 2^{6s-5}.
\end{aligned} \tag{16.45}$$

By (12.32) with j = s, $k>2s$ we get

$$\Gamma_{2s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(k+1)} - \Gamma_{2s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times k} = 3\cdot 2^{2k+2s-2}-3\cdot 2^{k+4s-4} \quad \text{if} \quad k>2s. \tag{16.46}$$

From (16.46), (16.45) we deduce

$$\begin{aligned}
&\sum_{j=2s+1}^{k}\left(\Gamma_{2s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(j+1)} - \Gamma_{2s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times j}\right) = \sum_{j=2s+1}^{k}(3\cdot 2^{2j+2s-2}-3\cdot 2^{j+4s-4}) \\
&\Leftrightarrow \sum_{j=2s+2}^{k+1}\Gamma_{2s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times j} - \sum_{j=2s+1}^{k}\Gamma_{2s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times j} = 2^{2k+2s}-3\cdot 2^{k+4s-3}-5\cdot 2^{6s-3} \\
&\Leftrightarrow \Gamma_{2s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(k+1)} - \Gamma_{2s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(2s+1)} = 2^{2k+2s}-3\cdot 2^{k+4s-3}-5\cdot 2^{6s-3} \\
&\Leftrightarrow \Gamma_{2s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(k+1)} = 2^{2k+2s}-3\cdot 2^{k+4s-3}-5\cdot 2^{6s-3}+21\cdot 2^{6s-5}
\end{aligned}$$

$$\Leftrightarrow \Gamma_{2s}^{\left[\begin{smallmatrix} s \\ s \end{smallmatrix}\right]\times(k+1)} = 2^{2k+2s}-3\cdot 2^{k+4s-3}+2^{6s-5} \qquad \text{if } k>2s. \tag{16.47}$$

By (16.45) we see that (16.47) holds for k = 2s, then (16.47) holds for $k\geq 2s$.
Consider the matrix

$$\left(\begin{array}{cccccc} \alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{k-2s+1} & \alpha_{k-2s+2} \\ \beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{k-2s+1} & \beta_{k-2s+2} \end{array}\right).$$

We see easily that

$$\Gamma_{1}^{\left[\substack{1\\1}\right]\times(k-2(s-1))} = 3\cdot(2^{k-2s+2}-1).$$

Hence

$$\Gamma_{2}^{\left[\substack{1\\1}\right]\times(k-2(s-1))} = 2^{2k-4s+4} - 3\cdot(2^{k-2s+2}-1) - 1 = 2^{2k-4s+4} - 3\cdot 2^{k-2s+2} + 2. \tag{16.48}$$

From (16.48) and (16.47) with $k \geq 2s$ we get

$$\Gamma_{2s}^{\left[\substack{s\\s}\right]\times k} = 2^{2k+2s-2} - 3\cdot 2^{k+4s-4} + 2^{6s-5} = 8^{2(s-1)}\cdot\Gamma_{2}^{\left[\substack{1\\1}\right]\times(k-2(s-1))}.$$

□

17. A REDUCTION FORMULA FOR $\Gamma_{s+m+1+j}^{\left[\substack{s\\s+m}\right]\times k}$ IN THE CASE $1 \leq j \leq s-1,\ k \geq s+m+1+j$

In this section we prove the following reduction formula by induction on j

$$\Gamma_{s+m+1+j}^{\left[\substack{s\\s+m}\right]\times k} = 8^{2j+m}\cdot\Gamma_{s-j+1}^{\left[\substack{s-j\\s-j}\right]\times(k-m-2j)} \qquad \text{if}\quad 0 \leq j \leq s-1,\ k \geq s+m+1+j.$$

We recall again that the right hand side in the above equation has been computed in section 13.
In fact we have

$$\Gamma_{s-j+1}^{\left[\substack{s-j\\s-j}\right]\times(k-m-2j)} = \begin{cases} 2^{4s-4j} - 3\cdot 2^{3s-3j-1} + 2^{2s-2j-1} & \text{if}\quad 0 \leq j \leq s-1,\ k = s+m+j+1,\\ 21\cdot[2^{k-m+s-3j-1} + 2^{3s-3j-1} - 5\cdot 2^{2s-2j-1}] & \text{if}\quad 0 \leq j \leq s-2,\ k > s+m+j+1,\\ 2^{2k-2m-4s+4} - 3\cdot 2^{k-m-2s+2} + 2 & \text{if}\quad j = s-1,\ k > 2s+m. \end{cases}$$

Lemma 17.1. *We have*
(17.1)

$$\Gamma_{s+m+1}^{\left[\substack{s\\s+m}\right]\times k} = 8^{m}\cdot\Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(k-m)} \qquad \textit{if}\quad k \geq s+m+1,$$

(17.2)

$$\Gamma_{s+m+1}^{\left[\substack{s\\s+m}\right]\times(s+m+1)} = 2^{4s+3m} - 3\cdot 2^{3s+3m-1} + 2^{2s+3m-1} = 8^{m}\cdot\Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(s+1)} \qquad \textit{if}\quad k = s+m+1,$$

$$\Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(s+1)} = 2^{4s} - 3\cdot 2^{3s-1} + 2^{2s-1},$$

(17.3)

$$\Gamma_{s+m+1}^{\left[\substack{s\\s+m}\right]\times k} = 21\cdot(2^{k+s+2m-1} + 2^{3s+3m-1} - 5\cdot 2^{2s+3m-1}) = 8^{m}\cdot\Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(k-m)} \qquad \textit{if}\quad k > s+m+1,$$

$$\Gamma_{s+1}^{\left[\substack{s\\s}\right]\times(k-m)} = 21\cdot(2^{k-m+s-1} + 2^{3s-1} - 5\cdot 2^{2s-1}).$$

Proof. Proof of (17.2)

By (15.32) and (13.3) we get

$$\Gamma_{s+m+1}^{\left[\substack{s\\ s+m}\right]\times(s+m+1)} = 2^{4s+3m} - 3\cdot 2^{3s+3m-1} + 2^{2s+3m-1} = 8^{m}\cdot\Gamma_{s+1}^{\left[\substack{s\\ s}\right]\times(s+1)}.$$

Proof of (17.3)

Consider the matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{s+m+1} & \alpha_{s+m+2}\\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+m+2} & \alpha_{s+m+3}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s+m-1} & \alpha_{2s+m}\\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s+m} & \alpha_{2s+m+1}\\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{s+m+1} & \beta_{s+m+2}\\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+m+2} & \beta_{s+m+3}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s+m-1} & \beta_{2s+m}\\
\beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s+m} & \beta_{2s+m+1}\\
\beta_{s+1} & \beta_{s+2} & \beta_{s+3} & \ldots & \beta_{2s+m+1} & \beta_{2s+m+2}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{2s+2m} & \beta_{2s+2m+1}
\end{pmatrix}.$$

We have

(17.4)

$$\sum_{i=0}^{s+m+2}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(s+m+2)} = 2^{4s+3m+2} \qquad \text{by (11.1) with} \quad \text{k = s+ m +2,}$$

(17.5)

$$\sum_{i=0}^{s+m+2}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(s+m+2)}\cdot 2^{-i} = 2^{3s+m} + 2^{2s+2m+2} - 2^{s+m} \qquad \text{by (11.2) with} \quad \text{k = s+ m+2,}$$

(17.6)

$$\sum_{i=0}^{s-1}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(s+m+2)} = 3\cdot 2^{3s-4} - 2^{2s-3} \qquad \text{by (11.27) with} \quad \text{k = s+ m+2,}$$

(17.7)

$$\sum_{i=0}^{s-1}\Gamma_{i}^{\left[\substack{s\\ s+m}\right]\times(s+m+2)}\cdot 2^{-i} = 7\cdot 2^{2s-4} - 3\cdot 2^{s-3} \qquad \text{by (11.28) with} \quad \text{k = s+m+2,}$$

(17.8)

$$\Gamma_{s}^{\left[\substack{s\\ s+m}\right]\times(s+m+2)} = 2^{2s+m+1} + 21\cdot 2^{3s-4} - 11\cdot 2^{2s-3} \qquad \text{by (13.46) with k = s+ m+2,}$$

(17.9)

$$\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times(s+m+2)} = 8^{j-1}\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+(m-(j-1))}\right]\times(s+m+2-(j-1))} \qquad \text{by (15.40) for } 0\leq j\leq m-1$$

$$= 3\cdot 2^{2s+m+2j-1}+21\cdot 2^{3s+3j-4}-21\cdot 2^{2s+3j-4} \qquad \text{and} \quad \text{k = s+ m+2 ,}$$

(17.10)

$$\Gamma_{s+m}^{\left[\substack{s\\ s+m}\right]\times(s+m+2)} = 11\cdot 2^{(s+m+2)+2m+s-3}+21\cdot 2^{3s+3m-4}-53\cdot 2^{2s+3m-4} \qquad \text{by (15.41) for } k=s+m+2$$

$$= 21\cdot 2^{3s+3m-4}+35\cdot 2^{2s+3m-4}.$$

From (17.4), . . . , (17.10) with k = s+m+2 we obtain

(17.11)

$$\Gamma_{s+m+1}^{\left[\substack{s\\ s+m}\right]\times(s+m+2)} + \Gamma_{s+m+2}^{\left[\substack{s\\ s+m}\right]\times(s+m+2)}$$

$$= 2^{4s+3m+2}-(3\cdot 2^{3s-4}-2^{2s-3})-(2^{2s+m+1}+21\cdot 2^{3s-4}-11\cdot 2^{2s-3})$$

$$-\sum_{j=1}^{m-1}[3\cdot 2^{2s+m+2j-1}+21\cdot 2^{3s+3j-4}-21\cdot 2^{2s+3j-4}]-(21\cdot 2^{3s+3m-4}+35\cdot 2^{2s+3m-4})$$

$$= 2^{4s+3m+2}-3\cdot 2^{3s+3m-1}-5\cdot 2^{2s+3m-1},$$

(17.12)

$$\Gamma_{s+m+1}^{\left[\substack{s\\ s+m}\right]\times(s+m+2)}\cdot 2^{-(s+m+1)} + \Gamma_{s+m+2}^{\left[\substack{s\\ s+m}\right]\times(s+m+2)}\cdot 2^{-(s+m+2)}$$

$$= 2^{3s+m}+2^{2s+2m+2}-2^{s+m}-(7\cdot 2^{2s-4}-3\cdot 2^{s-3})-(2^{2s+m+1}+21\cdot 2^{3s-4}-11\cdot 2^{2s-3})\cdot 2^{-s}$$

$$-\sum_{j=1}^{m-1}[3\cdot 2^{2s+m+2j-1}+21\cdot 2^{3s+3j-4}-21\cdot 2^{2s+3j-4}]\cdot 2^{-(s+j)}$$

$$-(21\cdot 2^{3s+3m-4}+35\cdot 2^{2s+3m-4})\cdot 2^{-(s+m)}$$

$$= 2^{3s+2m}+9\cdot 2^{2s+2m-2}-13\cdot 2^{s+2m-2}.$$

Hence by (17.11), (17.12) we deduce after some calculations

$$\Gamma_{s+m+1}^{\left[\substack{s\\ s+m}\right]\times(s+m+2)} = 21\cdot 2^{3s+3m-1}-21\cdot 2^{2s+3m-1}, \tag{17.13}$$

$$\Gamma_{s+m+2}^{\left[\substack{s\\ s+m}\right]\times(s+m+2)} = 2^{4s+3m+2}-3\cdot 2^{3s+3m+2}+2^{2s+3m+3}. \tag{17.14}$$

By (12.35) with j = 1, $k>s+m+1$ we get

$$\Gamma_{s+m+1}^{\left[\substack{s\\ s+m}\right]\times(k+1)} - \Gamma_{s+m+1}^{\left[\substack{s\\ s+m}\right]\times k} = 21\cdot 2^{k+s+2m-1} \quad \text{if} \quad k>s+m+1. \tag{17.15}$$

From (17.15), (17.13) we deduce

$$\sum_{j=s+m+2}^{k}\left(\Gamma_{s+m+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(j+1)} - \Gamma_{s+m+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times j}\right) = \sum_{j=s+m+2}^{k} 21\cdot 2^{j+s+2m-1}$$

$$\Leftrightarrow \sum_{j=s+m+3}^{k+1}\Gamma_{s+m+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times j} - \sum_{j=s+m+1}^{k}\Gamma_{s+m+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times j} = 21\cdot 2^{k+s+2m} - 21\cdot 2^{2s+3m+1}$$

$$\Leftrightarrow \Gamma_{s+m+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+m+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+m+2)} = 21\cdot 2^{k+s+2m} - 21\cdot 2^{2s+3m+1}$$

$$\Leftrightarrow \Gamma_{s+m+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(k+1)} = 21\cdot 2^{k+s+2m} - 21\cdot 2^{2s+3m+1} + 21\cdot(2^{3s+3m-1} - 21\cdot 2^{2s+3m-1})$$

(17.16)

$$\Leftrightarrow \Gamma_{s+m+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(k+1)} = 21[2^{k+s+2m} + 2^{3s+3m-1} - 5\cdot 2^{2s+3m-1}] \quad \text{if } k > s+m+1.$$

By (17.13) we see that (17.16) holds for k = s+ m+1, then (17.16) holds for $k \geq s+m+1$.

From (17.16) and (13.4) with $k \to k-m > s+1$ we get

$$\begin{aligned}\Gamma_{s+m+1}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k} &= 8^{m}\cdot\Gamma_{s+1}^{\left[\begin{smallmatrix} s\\ s\end{smallmatrix}\right]\times(k-m)} = 2^{3m}\cdot[21\cdot(2^{k-m+s-1} + 2^{3s-1} - 5\cdot 2^{2s-1})]\\ &= 21[2^{k+s+2m-1} + 2^{3s+3m-1} - 5\cdot 2^{2s+3m-1}].\end{aligned}$$

□

Lemma 17.2. *We have*

(17.17)

$$\Gamma_{s+m+1+j}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k} = 8^{2j+m}\cdot\Gamma_{s-j+1}^{\left[\begin{smallmatrix} s-j\\ s-j\end{smallmatrix}\right]\times(k-m-2j)} \qquad \textit{if}\quad 0\leq j\leq s-1,\ k\geq s+m+1+j,$$

(17.18)

$$\begin{aligned}\Gamma_{s+m+1+j}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times(s+m+1+j)} &= 8^{2j+m}\cdot\Gamma_{s-j+1}^{\left[\begin{smallmatrix} s-j\\ s-j\end{smallmatrix}\right]\times(s-j+1)} \qquad \textit{if}\quad 0\leq j\leq s-1,\ k = s+m+1+j\\ &= 8^{2j+m}\cdot(2^{4s-4j} - 3\cdot 2^{3s-3j-1} + 2^{2s-2j-1})\\ &= 2^{4s+3m+2j} - 3\cdot 2^{3s+3m+3j-1} + 2^{2s+3m+4j-1},\end{aligned}$$

(17.19)

$$\begin{aligned}\Gamma_{s+m+1+j}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k} &= 8^{2j+m}\cdot\Gamma_{s-j+1}^{\left[\begin{smallmatrix} s-j\\ s-j\end{smallmatrix}\right]\times(k-m-2j)}\\ &= 8^{2j+m}\cdot 21\cdot[2^{k-m-3j+s-1} + 2^{3s-3j-1} - 5\cdot 2^{2s-2j-1}] \qquad \textit{if}\ 0\leq j\leq s-2,\ k > s+m+1+j\\ &= 21\cdot[2^{k+2m+3j+s-1} + 2^{3s+3m+3j-1} - 5\cdot 2^{2s+3m+4j-1}],\end{aligned}$$

(17.20)

$$\begin{aligned}\Gamma_{2s+m}^{\left[\begin{smallmatrix} s\\ s+m\end{smallmatrix}\right]\times k} &= 8^{2s+m-2}\cdot\Gamma_{2}^{\left[\begin{smallmatrix} 1\\ 1\end{smallmatrix}\right]\times(k-m-2s+2)} \qquad \textit{if}\quad j = s-1,\ k > 2s+m\\ &= 8^{2s+m-2}\cdot[2^{2(k-m)-4s+4} - 3\cdot 2^{k-m-2s+2} + 2]\end{aligned}$$

$$= 2^{2k+2s+m-2} - 3\cdot 2^{k+2m+4s-4} + 2^{6s+3m-5}.$$

Proof. We will do the proof by induction on j.
Let l be a rational integer such that $1 \leq l \leq s-2$.
Let (H_{l-1}) denote the following statement

$$(17.21)\quad \Gamma_{s+m+1+j}^{\left[\substack{s\\ s+m}\right]\times k} = 8^{2j+m}\cdot \Gamma_{s-j+1}^{\left[\substack{s-j\\ s-j}\right]\times (k-m-2j)} \quad \text{if} \quad 0\leq j\leq l-1,\ k\geq s+m+1+j.$$

By Lemma 17.1 (H_{l-1}) holds for $l=1$.
Assume that (H_{l-1}) holds.
We are going to show that (H_l) holds, that is

$$(17.22)\qquad \Gamma_{s+m+1+l}^{\left[\substack{s\\ s+m}\right]\times k} = 8^{2l+m}\cdot \Gamma_{s-l+1}^{\left[\substack{s-l\\ s-l}\right]\times (k-m-2l)} \qquad \text{if} \quad k\geq s+m+l+1.$$

The case $k = s+m+l+1$
Consider the matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{s+m+l} & \alpha_{s+m+l+1}\\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+m+l+1} & \alpha_{s+m+l+2}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s+m+l-2} & \alpha_{2s+m+l-1}\\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s+m+l-1} & \alpha_{2s+m+l}\\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{s+m+l} & \beta_{s+m+l+1}\\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+m+l+1} & \beta_{s+m+l+2}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s+m+l-2} & \beta_{2s+m+l-1}\\
\beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s+m+l-1} & \beta_{2s+m+l}\\
\beta_{s+1} & \beta_{s+2} & \beta_{s+3} & \ldots & \beta_{2s+m+l} & \beta_{2s+m+l+1}\\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots\\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{2s+2m+l-1} & \beta_{2s+2m+l}
\end{pmatrix}.$$

We have

(17.23)
$$\sum_{i=0}^{s+m+l+1}\Gamma_i^{\left[\substack{s\\ s+m}\right]\times (s+m+l+1)} = 2^{4s+3m+2l} \qquad \text{by (11.1) with} \quad k = s+m+l+1,$$

(17.24)
$$\sum_{i=0}^{s-1}\Gamma_i^{\left[\substack{s\\ s+m}\right]\times (s+m+l+1)} = 3\cdot 2^{3s-4} - 2^{2s-3} \qquad \text{by (11.27) with} \quad k = s+m+l+1,$$

(17.25)

$$\Gamma_{s}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right] \times (s+m+l+1)} = 2^{2s+m+l} + 21 \cdot 2^{3s-4} - 11 \cdot 2^{2s-3} \qquad \text{by (13.46) with } k = s+m+l+1,$$

(17.26)

$$\begin{aligned} &\Gamma_{s+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right] \times (s+m+l+1)} = 8^{j-1} \cdot \Gamma_{s+1}^{\left[\begin{smallmatrix} s \\ s+(m-(j-1)) \end{smallmatrix}\right] \times (s+m+l+1-(j-1))} && \text{by (15.40) for } 1 \leq j \leq m-1 \\ &= 3 \cdot 2^{2s+m+l+2j-2} + 21 \cdot 2^{3s+3j-4} - 21 \cdot 2^{2s+3j-4} && \text{and } \ k = s+m+l+1 \ , \end{aligned}$$

(17.27)

$$\begin{aligned} &\Gamma_{s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right] \times (s+m+l+1)} = 11 \cdot 2^{(s+m+l+1)+2m+s-3} + 21 \cdot 2^{3s+3m-4} - 53 \cdot 2^{2s+3m-4} && \text{by (15.41) for } k = s+m+2 \\ &= 11 \cdot 2^{2s+3m+l-2} + 21 \cdot 2^{3s+3m-4} - 53 \cdot 2^{2s+3m-4}, \end{aligned}$$

(17.28)

$$\begin{aligned} &\Gamma_{s+m+1+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right] \times s+m+l+1} = 8^{2j+m} \cdot \Gamma_{s-j+1}^{\left[\begin{smallmatrix} s-j \\ s-j \end{smallmatrix}\right] \times ((s+m+l+1)-m-2j)} && \text{by } (H_{l-1}) \text{ (see (17.21))} \\ &= 8^{2j+m} \cdot 21 \cdot [2^{(s+m+l+1)-m-3j+s-1} + 2^{3s-3j-1} - 5 \cdot 2^{2s-2j-1}] && \text{if } \ 0 \leq j \leq l-1, \ k = s+m+l+1 \\ &= 21 \cdot [2^{2s+3m+l+3j} + 2^{3s+3m+3j-1} - 5 \cdot 2^{2s+3m+4j-1}]. \end{aligned}$$

From $(17.23), \ldots, (17.28)$ *with* $k = s+m+l+1$ we obtain after some calculations

(17.29)

$$\begin{aligned} &\Gamma_{s+m+l+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right] \times (s+m+l+1)} = 2^{4s+3m+2l} - (3 \cdot 2^{3s-4} - 2^{2s-3}) - (2^{2s+m+l} + 21 \cdot 2^{3s-4} - 11 \cdot 2^{2s-3}) \\ &- \sum_{j=1}^{m-1} [3 \cdot 2^{2s+m+l+2j-2} + 21 \cdot 2^{3s+3j-4} - 21 \cdot 2^{2s+3j-4}] - (11 \cdot 2^{2s+3m+l-2} + 21 \cdot 2^{3s+3m-4} - 53 \cdot 2^{2s+3m-4}) \\ &- \sum_{j=0}^{l-1} 21 \cdot [2^{2s+3m+l+3j} + 2^{3s+3m+3j-1} - 5 \cdot 2^{2s+3m+4j-1}] \\ &= 2^{4s+3m+2l} - 3 \cdot 2^{3s+3m+3l-1} + 2^{2s+3m+4l-1}. \end{aligned}$$

By (13.3) with $s \to s-l$ and (17.29) we get

$$\Gamma_{s+m+l+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right] \times (s+m+l+1)} = 8^{2l+m} \cdot [2^{4(s-l)} - 3 \cdot 2^{3(s-l)-1} + 2^{2(s-l)-1}] = 2^{6l+3m} \cdot \Gamma_{s-l+1}^{\left[\begin{smallmatrix} s-l \\ s-l \end{smallmatrix}\right] \times (s-l+1)}.$$

The case $k > s+m+l+1$

Consider the matrix

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \alpha_3 & \ldots & \alpha_{s+m+l+1} & \alpha_{s+m+l+2} \\
\alpha_2 & \alpha_3 & \alpha_4 & \ldots & \alpha_{s+m+l+2} & \alpha_{s+m+l+3} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \alpha_{s+1} & \ldots & \alpha_{2s+m+l-1} & \alpha_{2s+m+l} \\
\alpha_s & \alpha_{s+1} & \alpha_{s+2} & \ldots & \alpha_{2s+m+l} & \alpha_{2s+m+l+1} \\
\beta_1 & \beta_2 & \beta_3 & \ldots & \beta_{s+m+l+1} & \beta_{s+m+l+2} \\
\beta_2 & \beta_3 & \beta_4 & \ldots & \beta_{s+m+l+2} & \beta_{s+m+l+3} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s-1} & \beta_s & \beta_{s+1} & \ldots & \beta_{2s+m+l-1} & \beta_{2s+m+l} \\
\beta_s & \beta_{s+1} & \beta_{s+2} & \ldots & \beta_{2s+m+l} & \beta_{2s+m+l+1} \\
\beta_{s+1} & \beta_{s+2} & \beta_{s+3} & \ldots & \beta_{2s+m+l+1} & \beta_{2s+m+l+2} \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \ldots & \beta_{2s+2m+l} & \beta_{2s+2m+l+1}
\end{pmatrix}.$$

We have

(17.30)
$$\sum_{i=0}^{s+m+l+2} \Gamma_i^{\left[\substack{s\\ s+m}\right]\times(s+m+l+2)} = 2^{4s+3m+2l+2} \qquad \text{by (11.1) with } k=s+m+l+2,$$

(17.31)
$$\sum_{i=0}^{s+m+l+2} \Gamma_i^{\left[\substack{s\\ s+m}\right]\times(s+m+l+2)} \cdot 2^{-i} = 2^{3s+2m+l} + 2^{2s+2m+2l+2} - 2^{s+m+l} \qquad \text{by (11.2) with } k=s+m+l+2,$$

(17.32)
$$\sum_{i=0}^{s-1} \Gamma_i^{\left[\substack{s\\ s+m}\right]\times(s+m+l+2)} \cdot 2^{-i} = 7\cdot 2^{2s-4} - 3\cdot 2^{s-3} \qquad \text{by (11.28) with } k=s+m+l+2,$$

(17.33)
$$\sum_{i=0}^{s-1} \Gamma_i^{\left[\substack{s\\ s+m}\right]\times(s+m+l+2)} = 3\cdot 2^{3s-4} - 2^{2s-3} \qquad \text{by (11.27) with } k=s+m+l+2,$$

(17.34)
$$\Gamma_s^{\left[\substack{s\\ s+m}\right]\times(s+m+l+2)} = 2^{2s+m+l+1} + 21\cdot 2^{3s-4} - 11\cdot 2^{2s-3} \qquad \text{by (13.46) with } k=s+m+l+2,$$

(17.35)
$$\Gamma_{s+j}^{\left[\substack{s\\ s+m}\right]\times(s+m+l+2)} = 8^{j-1}\cdot\Gamma_{s+1}^{\left[\substack{s\\ s+(m-(j-1))}\right]\times(s+m+l+2-(j-1))} \qquad \text{by (15.40) for } 1\leq j\leq m-1$$
$$= 3\cdot 2^{2s+m+l+2j-1} + 21\cdot 2^{3s+3j-4} - 21\cdot 2^{2s+3j-4} \qquad \text{and } k=s+m+l+2,$$

(17.36)

$$\Gamma_{s+m}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+m+l+2)} = 11\cdot 2^{(s+m+l+2)+2m+s-3}+21\cdot 2^{3s+3m-4}-53\cdot 2^{2s+3m-4} \quad \text{by (15.41) for } k=s+m+l+2$$
$$= 11\cdot 2^{2s+3m+l-1}+21\cdot 2^{3s+3m-4}-53\cdot 2^{2s+3m-4},$$

(17.37)

$$\Gamma_{s+m+1+j}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times s+m+l+2} = 8^{2j+m}\cdot\Gamma_{s-j+1}^{\left[\begin{smallmatrix} s-j \\ s-j \end{smallmatrix}\right]\times((s+m+l+2)-m-2j)} \quad \text{by } (H_{l-1}) \text{ (see (17.21))}$$
$$= 8^{2j+m}\cdot 21\cdot[2^{(s+m+l+2)-m-3j+s-1}+2^{3s-3j-1}-5\cdot 2^{2s-2j-1}] \quad \text{if } 0\leq j\leq l-1,\ k=s+m+l+2$$
$$= 21\cdot[2^{2s+3m+l+3j+1}+2^{3s+3m+3j-1}-5\cdot 2^{2s+3m+4j-1}].$$

From $(17.30),\ldots,(17.37)$ with $k=s+m+l+2$ we obtain after some calculations

(17.38)

$$\Gamma_{s+m+l+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+m+l+2)}+\Gamma_{s+m+l+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+m+l+2)}$$
$$= 2^{4s+3m+2l+2}-(3\cdot 2^{3s-4}-2^{2s-3})-(2^{2s+m+l+1}+21\cdot 2^{3s-4}-11\cdot 2^{2s-3})$$
$$-\sum_{j=1}^{m-1}[3\cdot 2^{2s+m+l+2j-1}+21\cdot 2^{3s+3j-4}-21\cdot 2^{2s+3j-4}]-(11\cdot 2^{2s+3m+l-1}+21\cdot 2^{3s+3m-4}-53\cdot 2^{2s+3m-4})$$
$$-\sum_{j=0}^{l-1}21\cdot[2^{2s+3m+l+3j+1}+2^{3s+3m+3j-1}-5\cdot 2^{2s+3m+4j-1}]$$
$$= 2^{4s+3m+2l+2}-3\cdot 2^{3s+3m+3l-1}-5\cdot 2^{2s+3m+4l-1},$$

(17.39)

$$\Gamma_{s+m+l+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+m+l+2)}\cdot 2^{-(s+m+l+1)}+\Gamma_{s+m+l+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+m+l+2)}\cdot 2^{-(s+m+l+2)}$$
$$= 2^{3s+2m+l}+2^{2s+2m+2l+2}-2^{s+m+l}-(7\cdot 2^{2s-4}-3\cdot 2^{s-3})-(2^{2s+m+l+1}+21\cdot 2^{3s-4}-11\cdot 2^{2s-3})\cdot 2^{-s}$$
$$-\sum_{j=1}^{m-1}[3\cdot 2^{2s+m+l+2j-1}+21\cdot 2^{3s+3j-4}-21\cdot 2^{2s+3j-4}]\cdot 2^{-(s+j)}$$
$$-(11\cdot 2^{2s+3m+l-1}+21\cdot 2^{3s+3m-4}-53\cdot 2^{2s+3m-4})\cdot 2^{-(s+m)}$$
$$-\sum_{j=0}^{l-1}21\cdot[2^{2s+3m+l+3j+1}+2^{3s+3m+3j-1}-5\cdot 2^{2s+3m+4j-1}]\cdot 2^{-(s+m+1+j)}$$
$$= 2^{3s+2m+l}+9\cdot 2^{2s+2m+2l-2}-13\cdot 2^{s+2m+3l-2}.$$

Hence by (17.38), (17.39) we deduce

$$\Gamma_{s+m+l+1}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+m+l+2)} = 21\cdot 2^{3s+3m+3l-1}-21\cdot 2^{2s+3m+4l-1}, \tag{17.40}$$

$$\Gamma_{s+m+l+2}^{\left[\begin{smallmatrix} s \\ s+m \end{smallmatrix}\right]\times(s+m+l+2)} = 2^{4s+3m+2l+2}-3\cdot 2^{3s+3m+3l+2}+2^{2s+3m+4l+3}. \tag{17.41}$$

By (12.35) with $j = l+1 (\leq s-1), k > s+m+l+1$ we get

$$\Gamma_{s+m+l+1}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+m+l+1}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times k} = 21\cdot 2^{k+s+2m+3l-1} \quad \text{if} \quad k > s+m+l+1. \tag{17.42}$$

From (17.42), (17.40) we deduce

$$\sum_{j=s+m+l+2}^{k}\left(\Gamma_{s+m+l+1}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times(j+1)} - \Gamma_{s+m+l+1}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times j}\right) = \sum_{j=s+m+l+2}^{k} 21\cdot 2^{j+s+2m+3l-1}$$

$$\Leftrightarrow \sum_{j=s+m+l+3}^{k+1}\Gamma_{s+m+1}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times j} - \sum_{j=s+m+l+2}^{k}\Gamma_{s+m+1}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times j} = 21\cdot 2^{k+s+2m+3l} - 21\cdot 2^{2s+3m+4l+1}$$

$$\Leftrightarrow \Gamma_{s+m+l+1}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times(k+1)} - \Gamma_{s+m+l+1}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times(s+m+l+2)} = 21\cdot 2^{k+s+2m+3l} - 21\cdot 2^{2s+3m+4l+1}$$

$$\Leftrightarrow \Gamma_{s+m+l+1}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times(k+1)} = 21\cdot 2^{k+s+2m+3l} - 21\cdot 2^{2s+3m+4l+1} + 21\cdot(2^{3s+3m+3l-1} - \cdot 2^{2s+3m+4l-1})$$

(17.43)

$$\Leftrightarrow \Gamma_{s+m+l+1}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times(k+1)} = 21[2^{k+s+2m+3l} + 2^{3s+3m+3l-1} - 5\cdot 2^{2s+3m+4l-1}] \quad \text{if } k > s+m+l+1.$$

By (17.40) we see that (17.43) holds for $k = s+m+l+1$, then (17.43) holds for $k \geq s+m+l+1$.
From (17.43) and (13.4) with $s \to s-l,\ k \to k-m-2l\ (> s-l+1)$ we get

$$\begin{aligned}\Gamma_{s+m+l+1}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times k} &= 8^{2l+m}\cdot\Gamma_{s-l+1}^{\left[\begin{smallmatrix}s-l\\ s-l\end{smallmatrix}\right]\times(k-m-2l)} = 2^{3m+6l}\cdot[21\cdot(2^{(k-m-2l)+(s-l)-1} + 2^{3(s-l)-1} - 5\cdot 2^{2(s-l)-1})] \\ &= 21[2^{k+s+2m+3l-1} + 2^{3s+3m+3l-1} - 5\cdot 2^{2s+3m+4l-1}] \quad \text{if } k > s+m+l+1.\end{aligned}$$

$\square$

We have now established that

$$\Gamma_{s+m+1+j}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times k} = 8^{2j+m}\cdot\Gamma_{s-j+1}^{\left[\begin{smallmatrix}s-j\\ s-j\end{smallmatrix}\right]\times(k-m-2j)} \quad \text{if} \quad 0 \leq j \leq s-2,\ k \geq s+m+1+j.$$

It remains to prove

$$\Gamma_{2s+m}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times k} = 8^{2s+m-2}\cdot\Gamma_{2}^{\left[\begin{smallmatrix}1\\ 1\end{smallmatrix}\right]\times(k-m-2(s-1))} \quad \text{if } k \geq 2s+m. \tag{17.44}$$

The case k = 2s + m
(17.41) holds for $l = s-2$, we then obtain by (16.48) with $k \to 2s$

$$8^{2(s-1)+m}\cdot\Gamma_{2}^{\left[\begin{smallmatrix}1\\ 1\end{smallmatrix}\right]\times 2} = 8^{2(s-1)+m}\cdot[2^4 - 3\cdot 2^2 + 2] = 2^{6s+3m-6}\cdot 6 = 3\cdot 2^{6s+3m-5} = \Gamma_{2s+m}^{\left[\begin{smallmatrix}s\\ s+m\end{smallmatrix}\right]\times(2s+m)}.$$

$\square$

The case $k > 2s+m$

We proceed as in the case $k > s+m+l+1$ with $l = s-1$.
From (17.38), with $l = s-1$ we get

(17.45)

$$\Gamma_{2s+m}^{\left[\substack{s\\s+m}\right]\times(2s+m+1)} = 2^{4s+3m+2(s-1)+2} - 3\cdot 2^{3s+3m+3(s-1)-1} - 5\cdot 2^{2s+3m+4(s-1)-1}$$
$$= 2^{6s+3m} - 3\cdot 2^{6s+3m-4} - 5\cdot 2^{6s+3m-5} = 21\cdot 2^{6s+3m-5}.$$

By (12.35) with $j = s, k > 2s+m$ we get

$$\text{(17.46)}\quad \Gamma_{2s+m}^{\left[\substack{s\\s+m}\right]\times(k+1)} - \Gamma_{2s+m}^{\left[\substack{s\\s+m}\right]\times k} = 3\cdot 2^{2k+2s+m-2} - 3\cdot 2^{k+4s+2m-4} \quad \text{if} \quad k > 2s+m$$

From (17.46), (17.45) we deduce

$$\sum_{j=2s+m+1}^{k}\left(\Gamma_{2s+m}^{\left[\substack{s\\s+m}\right]\times(j+1)} - \Gamma_{2s+m}^{\left[\substack{s\\s+m}\right]\times j}\right) = \sum_{j=2s+m+1}^{k} 3\cdot 2^{2j+2s+m-2} - 3\cdot 2^{j+4s+2m-4}$$
$$\Leftrightarrow \sum_{j=2s+m+2}^{k+1}\Gamma_{2s+m}^{\left[\substack{s\\s+m}\right]\times j} - \sum_{j=2s+m+1}^{k}\Gamma_{2s+m}^{\left[\substack{s\\s+m}\right]\times j} = 2^{2k+2s+m} - 3\cdot 2^{k+4s+2m-3} - 5\cdot 2^{6s+3m-3}$$
$$\Leftrightarrow \Gamma_{2s+m}^{\left[\substack{s\\s+m}\right]\times(k+1)} - \Gamma_{2s+m}^{\left[\substack{s\\s+m}\right]\times(2s+m+1)} = 2^{2k+2s+m} - 3\cdot 2^{k+4s+2m-3} - 5\cdot 2^{6s+3m-3}$$
$$\Leftrightarrow \Gamma_{2s+m}^{\left[\substack{s\\s+m}\right]\times(k+1)} = 2^{2k+2s+m} - 3\cdot 2^{k+4s+2m-3} - 5\cdot 2^{6s+3m-3} + 21\cdot 2^{6s+3m-5}$$

(17.47)

$$\Leftrightarrow \Gamma_{2s+m}^{\left[\substack{s\\s+m}\right]\times(k+1)} = 2^{2k+2s+m} - 3\cdot 2^{k+4s+2m-3} + 2^{6s+3m-5} \qquad \text{if } k > 2s+m.$$

By (17.45) we see that (17.47) holds for k = 2s+m, then (17.47) holds for $k \geq 2s+m$.
From (16.48)with $k \to k-m$, and (17.47) we obtain

$$8^{2s+m-2}\cdot\Gamma_{2}^{\left[\substack{1\\1}\right]\times(k-m-2(s-1))} = 2^{6s+3m-6}\cdot[2^{2(k-m)-4s+4} - 3\cdot 2^{k-m-2s+2} + 2]$$
$$= 2^{2k+m+2s-2} - 3\cdot 2^{k+2m+4s-4} + 2^{6s+3m-5}$$
$$= \Gamma_{2s+m}^{\left[\substack{s\\s+m}\right]\times k}.$$

$\square$

18. PROOF OF THEOREMS $3.1, 3.2, \ldots, 3.14, 3.15$

18.1. **PROOF OF THEOREM 3.1.** Follows from (4.1) and (4.19).

18.2. **PROOF OF THEOREM 3.2.** Follows from Lemma 5.14.

18.3. **PROOF OF THEOREM 3.3.** Follows from Lemma 8.3.

18.4. **PROOF OF THEOREM 3.4.** Follows from Lemma 8.4.

18.5. **PROOF OF THEOREM 3.5.** Follows from Lemma 8.5.

18.6. **PROOF OF THEOREM 3.6.** We have
(3.10) follows from (11.21),
(3.11) follows from (12.1),
(3.12) follows from (12.15),
(3.13) follows from (12.20),
(3.14) follows from (12.21).

18.7. **PROOF OF THEOREM 3.7.** Follows from Lemma 15.2.

18.8. **PROOF OF THEOREM 3.8.** We have
(3.19) follows from (17.17),
(3.20) follows from (16.17),
(3.21) follows from (16.18) with $k \to k - m$,
(3.22) follows from (17.20).

18.9. **PROOF OF THEOREM 3.9.** We have in the following cases :
From (11.21) we get $\Gamma_i^{\begin{bmatrix} s \\ s \end{bmatrix} \times k}$ in the case $1 \leq i \leq s-1,\ k > i$.
From (13.2) we get $\Gamma_i^{\begin{bmatrix} s \\ s \end{bmatrix} \times k}$ in the case $i = s,\ k > s$.
From (16.16), (16.18) and (16.19) we deduce $\Gamma_i^{\begin{bmatrix} s \\ s \end{bmatrix} \times k}$ in the case $s+1 \leq i \leq 2s,\ k > i$.

18.10. **PROOF OF THEOREM 3.10.** We have in the following two cases :
From (11.26) we get $\Gamma_i^{\begin{bmatrix} s \\ s \end{bmatrix} \times i}$ in the case $1 \leq i \leq s$.
From (16.16), (16.17) we deduce $\Gamma_i^{\begin{bmatrix} s \\ s \end{bmatrix} \times i}$ in the case $s+1 \leq i \leq 2s$.

18.11. **PROOF OF THEOREM 3.11.** We have in the following cases :
From (13.21) we get $\Gamma_i^{\begin{bmatrix} s \\ s+1 \end{bmatrix} \times k}$ in the case $1 \leq i \leq s-1,\ k > i$.
From (13.24) we get $\Gamma_i^{\begin{bmatrix} s \\ s+1 \end{bmatrix} \times k}$ in the case $i = s,\ k > s$.
From (13.26) we get $\Gamma_i^{\begin{bmatrix} s \\ s+1 \end{bmatrix} \times k}$ in the case $i = s+1,\ k > s+1$.
From (17.19) with j = i-s-2 we get $\Gamma_i^{\begin{bmatrix} s \\ s+1 \end{bmatrix} \times k}$ in the case $s+2 \leq i \leq 2s,\ k > i$.
From (17.20) with m = 1 we get $\Gamma_i^{\begin{bmatrix} s \\ s+1 \end{bmatrix} \times k}$ in the case $i = 2s+1,\ k > 2s+1$.

18.12. **PROOF OF THEOREM 3.12.** We have in the following two cases :
From (13.25) and (11.26) with m = 1 and $k \to i$ we get $\Gamma_i^{\begin{bmatrix} s \\ s+1 \end{bmatrix} \times i}$ in the case $1 \leq i \leq s+1$.
From (17.18) with m = 1 and j = i - s - 2 we deduce $\Gamma_i^{\begin{bmatrix} s \\ s+1 \end{bmatrix} \times i}$ in the case $s+2 \leq i \leq 2s+1$.

18.13. **PROOF OF THEOREM 3.13.** We have in the following cases :
From (13.21) we get $\Gamma_i^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times k}$ in the case $1 \leq i \leq s-1,\ k > i$.
From (13.46) we get $\Gamma_i^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times k}$ in the case $i = s,\ k > s$.

From (15.40) with $l = i-s$ we get $\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times k}$ in the case $s+1 \leq i \leq s+m-1,\ k > i.$

From (15.41) we get $\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times k}$ in the case $i = s+m,\ k > i.$

From (17.19) with j = i-s-m-1 we get $\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times k}$ in the case $s+m+1 \leq i \leq 2s+m-1,\ k > i.$

From (17.20) we get $\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times k}$ in the case $i = 2s+m,\ k > i.$

18.14. **PROOF OF THEOREM 3.14.** We have in the following three cases :

From (11.26) with $k \to i$ and (13.47) we get $\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times i}$ in the case $1 \leq i \leq s+1.$

From (15.39) with $l \to i$ and (17.2) we get $\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times i}$ in the case $s+2 \leq i \leq s+m+1.$

From (17.18) with j = i-s-m-1 we get $\Gamma_{i}^{\left[\substack{s \\ s+m}\right]\times i}$ in the case $s+m+2 \leq i \leq 2s+m.$

18.15. **PROOF OF THEOREM 3.15.** Follows from Lemma 4.13.

References

[1] Daykin David E, Distribution of Bordered Persymmetric Matrices in a finite field J. reine angew. Math, **203** (1960), 47-54.

[2] Hayes , D.R, The expression of a polynomial as a sum of three irreducibles Acta Arith., **11** (1966), 461-488.

[3] Landsberg, G Ueber eine Anzahlbestimmung und eine damit zusammenhangende Reihe, J. reine angew.Math, **111** (1893), 87-88..

[4] Cherly, J Exponential sums and rank of persymmetric matrices over F_2, arXiv: 0711.1306

Département de Mathématiques, Université de Brest, 29238 Brest cedex 3, France
E-mail address: `Jorgen.Cherly@univ-brest.fr`